

INTRALIPID[®] 10%, 20% & 30%

Sterile fat emulsions containing soya oil, egg lecithin and glycerol

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Intralipid. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Intralipid against any benefits they expect it will have for you.

Please read this leaflet carefully before using Intralipid 10%, 20% and 30%. If you have any questions or are unsure about anything, please ask your doctor or pharmacist.

Keep this leaflet with the medicine. You may need to read it again.

What is Intralipid used for and how does it work?

Intralipid is a sterile fat emulsion, which provides your body with energy and fatty acids. The active ingredient in Intralipid is soya oil. Soya oil consists of a mixture of mainly polyunsaturated fatty acids. It also contains egg lecithin as an emulsifier, which is isolated from egg yolk. It is available in 3 strengths namely 10%, 20% and 30%.

When the intake of nutrients or food into the mouth or directly into the gut is not possible, or it is not enough to supply the body's needs, then intravenous nutrients or foods can be given. This is especially important for people whose bodies are under physical

stress from illness or recent surgery. During illness or after surgery the body requires nutrition or food. It is usually given together with carbohydrates, amino acids, salts, trace elements and vitamins to provide a complete intravenous diet.

Before you are given Intralipid

You should not be given Intralipid if

- You have an allergy to soya oil, eggs, peanuts or any of the ingredients listed at the end of this leaflet.
- You have an inability to break down fats

If you are not sure whether any of these apply to you, check with your doctor.

You should tell your doctor BEFORE using Intralipid if the answer to any of the following questions is YES.

- Are you pregnant or trying to become pregnant?
- Are you breastfeeding?
- Are you allergic to soya-, egg- or peanut protein?
- Do you have liver or kidney disease?
- Are you a diabetic?
- Do you have a disease of the pancreas?
- Do you have a disorder of the thyroid gland?
- Are you taking anticoagulants (medicines for preventing blood clotting)?
- Are you taking any other medicines including any that

you buy without a prescription from your pharmacy, supermarket or healthfood shop. These medicines may affect the action of Intralipid or may affect how well Intralipid works.

If you have not told your doctors about any of the above, tell them before you are given Intralipid.

How is Intralipid given?

The dose of Intralipid which you will require will be determined by your doctor or pharmacist. Your doctor will supervise your treatment with Intralipid.

Side Effects

Intralipid, like all other nutrient solutions which are given intravenously, may cause unwanted effects in some people. All the unwanted effects associated with Intralipid may not yet have been detected.

Intralipid may cause a rise in your body temperature (this occurs in less than 3% of people who use it). Shivering, chills and nausea and vomiting occur less often (in less than 1% of people who use it). Reports of other unwanted effects caused by Intralipid are extremely rare.

Unwanted effects which have been reported to occur immediately or soon after Intralipid has been given include: allergic reactions (skin rash and hives); breathing difficulties (rapid breathing, shortness of

CMI for Intralipid

breath); effects on blood pressure; abdominal pain; tiredness; headaches; flushing; slight pressure over the eyes and dizziness.

If you have these or any other unwanted effects during treatment, tell your doctor.

If you are given too much (overdose)

This rarely happens as Intralipid is usually administered under the care of a trained professional in a hospital or clinic setting.

However, if you are given Intralipid too quickly or too much, you may experience the following side effects: fever or convulsions.

Your doctor has information on how to recognise and treat an overdose. Ask your doctor if you have any concerns.

Storage

The expiry date of Intralipid is on the label of the pack. Intralipid should not be used if the expiry date has passed.

Intralipid should be stored below 25°C, but not frozen. Do not use Intralipid if it has been frozen.

The contents of each bottle of Intralipid are for single infusion only. Any unused Intralipid should be discarded. Do not use Intralipid if it is discoloured. Do not use if the product does not look quite right.

List of other ingredients

Intralipid also contains other ingredients such as glycerol, egg lecithin, sodium hydroxide and water for injections.

Intralipid does not contain any preservative.

Product Description***What it looks like***

Intralipid is a milky white emulsion which is supplied in glass bottles or plastic bags.

The active ingredient is soya oil. Intralipid comes in three strengths: 10%, 20% and 30%. It also contains egg lecithin, glycerol, sodium hydroxide and water for injections.

1000mL of Intralipid supplies the following calories:

10%	4600 kJ
20%	8400 kJ
30%	12600kJ.

Intralipid does not contain gluten, lactose, sucrose, tartrazine or any other azo dyes.

Intralipid comes in different pack sizes and can be identified by AUST R numbers:

Bottle

Intralipid 10%, 500 mL	AUST R 14471
Intralipid 20%, 100 mL	AUST R 14472
Intralipid 20%, 500 mL	AUST R 48245
Intralipid 20%, 1000 mL	AUST R 48246
Intralipid 30%, 250 mL	AUST R 53538
Intralipid 30%, 333 mL	AUST R 53539
Intralipid 30%, 500 mL	AUST R 53540
Intralipid 30%, 1000 mL	AUST R 53541

Bags

Intralipid 10%, 500 mL	AUST R 77744
Intralipid 20%, 100 mL	AUST R 77745
Intralipid 20%, 500 mL	AUST R 77746
Intralipid 30%, 250 mL	AUST R 77747

Intralipid 30%, 333 mL AUST R 77748

In New Zealand, the following pack sizes are available:

Bottles

Intralipid 10%: 100ml, 250ml, 500 mL
 Intralipid 20%: 100 mL, 250ml, 500 mL, 1000 mL
 Intralipid 30%: 250 mL, 333 mL, 500 mL, 1000 mL

Bags

Intralipid 10%: 500 mL
 Intralipid 20%: 100 mL, 500 mL
 Intralipid 30%: 250 mL, 333 mL

It may be supplied from the pharmacy as a mixture of amino acids, glucose, and vitamins. In this case it would be a milky white mixture in a plastic bag.

Further Information

More detailed information is available from your doctor or pharmacist. Therefore, if you have any concerns about the information or about Intralipid ask your doctor or pharmacist.

Sponsor

Australia:
 Fresenius Kabi Australia Pty Limited
 964 Pacific Highway
 Pymble NSW 2073
 Telephone: (02) 9391 5555

New Zealand:
 Fresenius Kabi New Zealand Limited
 60 Pavilion Drive
 Airport Oaks, Auckland 2022
 New Zealand
 Freecall: 0800 144 892

® = Registered Trademark

Date of Information

This leaflet was prepared in May, 2010.