

ACCUPRIL[®]

Quinapril hydrochloride

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about ACCUPRIL.

It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking ACCUPRIL against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with your ACCUPRIL tablets.

You may need to read it again.

What ACCUPRIL is used for

ACCUPRIL is used to lower high blood pressure (hypertension). It is also used to treat heart failure.

Hypertension

ACCUPRIL is used to lower high blood pressure (hypertension). Everyone has blood pressure. This pressure helps get your blood all around your body. Your blood pressure may be different at different times of the day, depending on how busy or worried you are. You have hypertension (high blood pressure) when your blood pressure stays higher than is needed, even when you are calm and relaxed.

There are usually no symptoms of hypertension. The only way of knowing that you have hypertension is to have your blood pressure checked on a regular basis. If high blood pressure is not treated it can lead to serious health problems, including stroke, heart disease and kidney failure.

Heart Failure

Heart failure means that the heart muscle is weak and cannot pump blood strongly enough to supply all the blood needed throughout the body. Heart failure is not the same as heart attack and does not mean that the heart stops. Heart failure may start off with no symptoms, but as the condition progresses, patients may feel short of breath or may get tired easily after light physical activity such as walking. Some patients may wake up short of breath at night. Fluid may collect in different parts of the body, often first noticed as swollen ankles and feet.

How ACCUPRIL works

ACCUPRIL works by widening your blood vessels, which reduces pressure in the vessels, making it easier for your heart to pump blood around your body. This helps increase the supply of oxygen to your heart, so that when you place extra demands on your heart, such as during exercise, your heart may cope better and you may not get short of breath as easily.

ACCUPRIL belongs to a group of medicines called angiotensin converting enzyme (ACE) inhibitors.

Your doctor may have prescribed ACCUPRIL for another reason. Ask your doctor if you have any questions about why ACCUPRIL has been prescribed for you.

ACCUPRIL is not addictive.

This medicine is available only with a doctor's prescription.

Before you take ACCUPRIL

When you must not take it

Do not take ACCUPRIL if:

- **you have an allergy to ACCUPRIL or any other medicine containing quinapril, or any of the ingredients listed at the end of this leaflet.**

Symptoms of an allergic reaction to ACCUPRIL may include skin rash, itchiness, shortness of breath, swelling of the face, lips or tongue, muscle pain or tenderness or joint pain.

- **you have taken any other 'ACE inhibitor' medicine before, which caused your face, lips, tongue, throat, hands or feet to swell up, or made it hard for you to breathe.**

If you have had an allergic reaction to an ACE inhibitor before, you may be allergic to ACCUPRIL.

- **you or your family have a history of swelling of the face, lips, tongue, throat, hands or feet for no apparent reason**

- you have kidney problems or a condition called 'renal artery stenosis'
- you have regular dialysis for blood filtration

You may experience an allergic reaction.

Do not take ACCUPRIL if you are pregnant or breastfeeding.

ACCUPRIL may enter your womb or it may pass into the breast milk and there is the possibility that your baby may be affected.

Do not take ACCUPRIL after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If you are not sure whether you should start taking ACCUPRIL, talk to your doctor.

Before you start to take it

Tell your doctor if you have allergies to:

- any other medicines
- any other substances, such as foods, preservatives or dyes.

Tell your doctor if you have or have had any medical conditions, especially the following:

- kidney problems, or are having dialysis
- liver problems
- heart problems
- low blood pressure, which you may notice as dizziness or light-headedness
- diabetes
- high levels of potassium in your blood.

You must also tell your doctor if you:

- are following a very low salt diet
- are about to receive desensitisation therapy for an allergy
- are about to undergo dialysis or lipoprotein apheresis

- are about to have surgery or a general anaesthetic
- plan to become pregnant or breastfeed.

If you have not told your doctor about any of the above, tell them before you start taking ACCUPRIL.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and ACCUPRIL may interfere with each other. These include:

- other medicines used to treat high blood pressure
- other medicines that work in a similar fashion to ACE Inhibitors, such as Angiotensin Receptor Blockers (these are used to treat high blood pressure and/or heart failure).
- diuretics, also known as fluid or water tablets
- non-steroidal anti-inflammatory drugs (NSAIDs) or Cox 2 inhibiting medicines used to relieve pain, swelling and other symptoms of inflammation, including arthritis
- potassium supplements or potassium-containing salt substitutes
- lithium, a medicine used to treat mood swings and some types of depression
- tetracycline antibiotics
- certain medicines used to treat bacterial and fungal infections

These medicines may be affected by ACCUPRIL, or may affect how well it works. You may need different amounts of your medicine, or you may need to take different medicines. Your doctor will advise you.

Your doctor and pharmacist may have more information on medicines to be careful with or avoid while taking ACCUPRIL.

Use in children

The safety and effectiveness of ACCUPRIL in children have not been established.

How to take ACCUPRIL

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet.

Your doctor or pharmacist will tell you how many tablets you will need to take each day. This depends on your condition and whether or not you are taking any other medicines.

If you do not understand the instructions on the box, ask your doctor or pharmacist for help.

How much to take

For high blood pressure

For most patients, not on diuretics, the usual starting dose is 5 to 10 mg taken once a day. The dose may need to be increased depending on your blood pressure at an interval of 4 weeks. Most patients take between 10 and 40 mg each day.

This dose may be taken once a day or divided into two equal doses per day.

For heart failure

The usual starting dose is 5 mg taken once a day. In most patients, effective doses are between 10 and 20 mg a day. Your doctor will advise whether the dose is to be taken as a single dose or as two separate doses.

How to take it

Swallow ACCUPRIL whole with a full glass of water.

Do not chew the tablets.

When to take it

Take ACCUPRIL at about the same time each day.

Taking your tablets at the same time each day will have the best effect. It will also help you remember when to take the tablets.

Take ACCUPRIL before meals.

Food with a high fat content may interfere with the absorption of ACCUPRIL.

How long to take it

ACCUPRIL helps control your condition, but does not cure it. Therefore you must take ACCUPRIL every day. Continue taking your medicine for as long as your doctor tells you.

If you forget to take it

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as soon as you remember and then go back to taking your medicine as you would normally.

Do not take a double dose to make up for the dose that you missed.

This may increase the chance of you getting an unwanted side effect.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or the Australian Poisons Information Centre (telephone 13 11 26), or the New Zealand National Poisons Information Centre (telephone 0800 POISON or 0800 764 766), or go to Accident and Emergency at the nearest

hospital, if you think that you or anyone else may have taken too much ACCUPRIL. Do this even if there are no signs of discomfort or poisoning.

If you take too much ACCUPRIL, you may feel light-headed, dizzy or you may faint.

While you are taking ACCUPRIL

Things you must do

If you are about to be started on any new medicine, tell your doctor, dentist or pharmacist that you are taking ACCUPRIL.

Make sure you drink enough water during exercise and hot weather when you are taking ACCUPRIL, especially if you sweat a lot.

If you do not drink enough water while taking ACCUPRIL, you may feel faint, light-headed or sick. This is because your blood pressure is dropping suddenly. If you continue to feel unwell, tell your doctor.

If you have excess vomiting or diarrhoea while taking ACCUPRIL, tell your doctor.

You may lose too much water and salt and your blood pressure may drop too much.

If you feel light-headed or dizzy after taking your first dose of ACCUPRIL, or when your dose is increased, tell your doctor immediately.

This is especially important if you are taking ACCUPRIL for heart failure.

If you plan to have surgery that needs a general anaesthetic, tell your doctor or dentist that you are taking ACCUPRIL.

Your blood pressure may drop suddenly.

If you become pregnant while taking ACCUPRIL, tell your doctor immediately.

If you are about to have any blood tests, tell your doctor that you are taking ACCUPRIL.

ACCUPRIL may interfere with the results of some tests.

Have your blood pressure checked when your doctor says, to make sure ACCUPRIL is working.

Go to your doctor regularly for a check-up.

Your doctor may occasionally do a blood test to check your potassium levels and see how your kidneys are working.

Things you must not do

Do not give ACCUPRIL to anyone else, even if they have the same condition as you.

Do not take ACCUPRIL to treat any other complaints unless your doctor or pharmacist tells you to.

Do not stop taking ACCUPRIL, or lower the dosage, without checking with your doctor.

Things to be careful of

If you feel light-headed, dizzy or faint when getting out of bed or standing up, get up slowly.

Standing up slowly, especially when you get up from bed or chairs, will help your body get used to the change in position and blood pressure. If this problem continues or gets worse, talk to your doctor.

Be careful driving or operating machinery until you know how ACCUPRIL affects you.

As with other ACE inhibitor medicines, ACCUPRIL may cause dizziness, light-headedness or tiredness in some people. Make sure you know how you react to ACCUPRIL before you drive a car, operate machinery, or do anything else that could be dangerous if you are dizzy or light-headed. If this occurs do not drive. If you drink alcohol, dizziness or light-headedness may be worse.

Things that would be helpful for your blood pressure or heart failure

Some self help measures suggested below may help your condition.

Talk to your doctor or pharmacist about these measures and for more information.

- **Alcohol**

your doctor may advise you to limit your alcohol intake.

- **Weight**

your doctor may suggest losing some weight to help lower your blood pressure and help lessen the amount of work your heart has to do. Some people may need a dietician's help to lose weight

- **Diet**

eat a healthy diet which includes plenty of fresh vegetables, fruit, bread (preferably wholegrain), cereals and fish. Also eat less sugar and fat (especially saturated fat) which includes sausages, fatty meats, full cream dairy products, biscuits, cakes, pastries, chocolates, chips and coconut. Monounsaturated and polyunsaturated fats from olive oil, canola oil, avocado and nuts are beneficial in small quantities.

- **Salt**

your doctor may advise you to watch the amount of salt in your diet. To reduce your salt intake you should avoid using salt in cooking or at the table and avoid cooked or processed foods containing high sodium (salt) levels.

- **Exercise**

regular exercise, maintained over the long term, helps to reduce blood pressure and helps get the heart fitter. Regular exercise also improves your blood cholesterol levels, helps reduce your weight and stress levels, and improves your sleep, mood and ability to concentrate. However, it is important not to overdo it.

Before starting any exercise, ask your doctor about the best kind of programme for you.

- **Smoking**

your doctor may advise you to stop smoking or at least cut down. Ask your doctor or pharmacist for further information and advice.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking ACCUPRIL.

ACCUPRIL helps most people with high blood pressure and heart failure, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

If you are over 65 years of age you may have an increased chance of getting side effects.

Ask your doctor or pharmacist to answer any questions you may have.

Following is a list of possible side effects. Do not be alarmed by this list. You may not experience any of them.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- feeling light-headed, dizzy or faint because your blood pressure is too low
- dry cough
- headache
- feeling sick (nausea) or vomiting
- stomach pain
- diarrhoea
- constipation
- aching, tender or weak joints or muscles not caused by exercise

- unusual tiredness or weakness, fatigue
- feeling drowsy or sleepy during the day
- feelings of deep sadness and unworthiness (depression)
- hair loss or thinning
- dry mouth or throat
- taste disturbances or loss of taste
- confusion or nervousness
- back pain
- rash
- difficulty in getting or maintaining an erection.

These side effects are usually mild.

Tell your doctor as soon as possible if you notice any of the following:

- disturbed vision
- symptoms of sunburn (such as redness, itching, swelling, blistering) which may occur more quickly than normal
- itchy, raised or red skin rash
- fast or irregular heart beat
- shortness of breath or tightness in the chest
- signs of worrying or frequent infections such as fever, severe chills, sore throat or mouth ulcers
- severe upper stomach pain, often with nausea and vomiting
- passing little or no urine
- bleeding or bruising more easily than normal.

These may be serious side effects. You may need medical attention. Serious side effects are rare.

If any of the following happen, stop taking ACCUPRIL and either tell your doctor immediately or go to Accident and Emergency at your nearest hospital:

- fainting within a few hours of taking a dose
- swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing

- sudden onset of stomach pains or cramps with or without nausea or vomiting
- severe flaking or peeling of the skin
- severe blisters and bleeding in the lips, eyes, mouth, nose and genitals
- chest pain.

These are very serious side effects. You may need urgent medical attention or hospitalisation. These side effects are very rare.

Other side effects not listed above may occur in some patients. Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

After taking ACCUPRIL

Storage

Keep your tablets in the box until it is time to take them.

If you take the tablets out of the box they will not keep well.

Keep your tablets in a cool dry place where the temperature stays below 25 degrees C.

Do not store ACCUPRIL or any other medicine in the bathroom or near a sink.

Do not leave it on a windowsill or in the car on hot days.

Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor or pharmacist tells you to stop taking ACCUPRIL or the tablets have passed their expiry date, ask your pharmacist what to do with any that are left over.

Product description

What it looks like

ACCUPRIL tablets come in three strengths and each has a different appearance:

- ACCUPRIL 5 mg - Reddish-brown, oval, biconvex, film-coated tablet with bisecting score on both sides and debossing "5" on both sides in opposite direction.
- ACCUPRIL 10 mg - Reddish-brown, triangular, biconvex film-coated tablet with bisecting score on both sides and debossing "10" on one side.
- ACCUPRIL 20 mg - Reddish-brown, round, biconvex film-coated tablet, with bisecting score on both sides and debossing "20" on one side.

A box of ACCUPRIL contains 30 tablets.

Ingredients

Active ingredient

- ACCUPRIL 5 mg - 5 mg quinapril (as the hydrochloride)
- ACCUPRIL 10 mg - 10 mg quinapril (as the hydrochloride)
- ACCUPRIL 20 mg - 20 mg quinapril (as the hydrochloride)

Other ingredients:

- magnesium carbonate
- lactose
- gelatin
- crospovidone
- magnesium stearate
- candelilla wax
- iron oxide red
- titanium dioxide

ACCUPRIL does not contain sucrose, gluten, tartrazine or any other azo dyes.

Supplier

ACCUPRIL is supplied in Australia by:

Pfizer Australia Pty Ltd

ABN 50 008 422 348

38-42 Wharf Road

West Ryde NSW 2114

Australia

Toll Free Number: 1800 675 229

ACCUPRIL is supplied in New Zealand by:

Pfizer New Zealand Ltd

PO Box 3998

Auckland

New Zealand

Toll Free Number: 0800 736 363

Australian Registration Numbers

ACCUPRIL 5 mg - AUST R 40926

ACCUPRIL 10 mg - AUST R 40928

ACCUPRIL 20 mg - AUST R 40930

This leaflet was prepared in January 2011

© Pfizer Australia Pty Ltd

®Registered Trademark.