

Curam[®] Duo 500/125

tablets

amoxicillin trihydrate/potassium clavulanate equivalent to 500 mg amoxicillin/125 mg clavulanic acid

Consumer Medicine Information (CMI)

What is in this leaflet

This leaflet answers some common questions about Curam Duo 500/125.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

Please read this leaflet before you start taking Curam Duo 500/125. If you are helping someone else take Curam Duo 500/125, please read this leaflet before you give the first dose.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking this medicine against the benefits they expect it will have.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What Curam Duo 500/125 is used for

This medicine is used for the short term treatment of a wide range of infections caused by bacteria. These infections may affect:

- the chest (bronchitis or pneumonia)
- the bladder (cystitis)
- the sinuses (sinusitis)
- the ears (otitis media)
- the bones and joints (osteomyelitis)
- or the skin.

Curam Duo 500/125 will not work against infections caused by viruses such as colds or the flu.

Curam Duo 500/125 is an antibiotic that belongs to a group of medicines called penicillins. It works by killing the bacteria that cause these infections.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

There is no evidence that this medicine is addictive.

This medicine is available only with a doctor's prescription.

Before you take Curam Duo 500/125

When you must not take it

Do not give this medicine if you have an allergy to:

- amoxicillin trihydrate or potassium clavulanate, the active ingredients, or to any of the other ingredients listed at the end of this leaflet under Product description
- any other similar medicines (such as penicillins, cephalosporins, carbapenems, monobactams or any other types of antibiotics).

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body

- rash, itching or hives on the skin.

If you have ever had an allergic reaction to cephalosporins (another type of antibiotic). You may have an increased chance of being allergic to Curam Duo 500/125 if you are allergic to cephalosporins.

Do not take this medicine if you have previously experienced liver problems or yellowing of the skin and/or eyes, also called jaundice, after taking Curam Duo 500/125 or any other medicines.

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should be given this medicine, talk to your doctor.

Before you start taking it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have had any of the following medical conditions:

- an allergic reaction (such as a rash) to any antibiotics in the past
- glandular fever (mononucleosis) or a blood disorder
- liver or kidney problems. The dosage of Curam Duo 500/125 may need to be changed or you may need to be given an alternative medicine
- you are taking warfarin or other medicines used to prevent blood clots

- pregnant or intend to become pregnant. Your doctor will discuss with you the possible risks and benefits of using Curam Duo 500/125 during pregnancy
- breastfeeding or wish to breastfeed. Your doctor will discuss with you the possible risks and benefits of using Curam Duo 500/125 during breastfeeding.
- passing less urine
- you are taking any other medicines or herbal remedies.

If you have not told your doctor about any of the above, tell him/her before you take Curam Duo 500/125.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Curam Duo 500/125 may interfere with each other. These include:

- probenecid
- allopurinol
- warfarin or other medicines used to prevent blood clots
- birth control pills ("the pill", oral contraceptives). Birth control pills may not work as well in preventing pregnancy when Curam Duo 500/125 is used. Talk to your doctor about the need for an additional method of contraception while taking Curam Duo 500/125.
- methotrexate, a medicine used to treat arthritis and some types of cancer
- mycophenolate mofetil, a medicine used to suppress your immune system
- other antibiotics. These may interfere with the actions of Curam Duo 500/125.

These medicines may be affected by Curam Duo 500/125 or may affect how well it works. You may need different amounts of their medicines, or may need to take different medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

How to take Curam Duo 500/125

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions, ask your doctor or pharmacist for help.

How much to take

Take Curam Duo 500/125 as directed by your doctor or pharmacist.

Usual dose - one Curam Duo 500/125 tablet every 12 hours.

These dosages may vary depending on your conditions and the severity of your illness.

Children weighing 40 kg and more should be dosed according to adult recommendations.

Curam Duo tablets are not recommended for children weighing less than 40 kg.

Ask your doctor or pharmacist if you are unsure of the correct dose for your child.

They will tell you exactly how much to give.

Follow the instructions they give you.

If you give the wrong dose, Curam Duo 500/125 may not work as well and your condition may not improve.

How and when to take it

Curam Duo tablets should be taken immediately before or with the first mouthful of food.

If you take it on an empty stomach, it may cause stomach upset.

How long to take Curam Duo 500/125

Keep taking this medicine for the full time of treatment, even if you begin to feel better after a few days.

If you do not complete the full course prescribed by your doctor, the infection may not clear completely or your symptoms may return.

Curam Duo 500/125 is not usually used for longer than 14 days without another check-up by the doctor.

If you forget to take it

Take your dose as soon as you remember, and continue to take it as you would normally.

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not give a double dose to make up for the dose that you missed.

This may increase the chance of you getting an unwanted side effect.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering to give the medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or the Poisons Information Centre (telephone 0800 POISON or 0800 764766) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much Curam Duo 500/125. Do this even if there are no signs of discomfort or poisoning.

Symptoms of overdose include mild to severe nausea, vomiting, cramps and diarrhoea.

You may need urgent medical attention.

While you are taking Curam Duo 500/125

Things you must do

If the symptoms of your infection do not improve within a few days, or if they become worse, tell your doctor.

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking Curam Duo 500/125.

Tell any other doctors, dentists, and pharmacists who are treating you that you are taking this medicine.

If you are about to have any blood tests, tell your doctor that you are taking this medicine.

It may interfere with the results of some tests.

Keep all of your doctor's appointments so that your progress can be checked.

Your doctor may do some tests from time to time to make sure the medicine is working and to prevent unwanted side effects.

Tell your doctor if, for any reason, you have not taken the medicine exactly as directed.

Otherwise, your doctor may think that it was not working as it should and change your treatment unnecessarily.

If you develop itching, swelling, difficulty breathing or a skin rash while taking Curam Duo 500/125, do not take any more and tell the doctor immediately.

If you develop severe diarrhoea or stomach cramps, either while taking Curam Duo 500/125 or within several weeks after treatment, tell the

doctor as soon as possible.

Diarrhoea or stomach cramps may mean that you have a serious condition affecting your bowel. You may need urgent medical care. Do not take any diarrhoea medicine without first checking with your doctor.

If you get a sore white mouth or tongue while taking or soon after stopping Curam Duo 500/125, tell your doctor. Also tell your doctor if you get vaginal itching or discharge. This may mean that you have a fungal infection called thrush.

Tell your doctor if you become pregnant while you are taking Curam Duo 500/125.

Drink plenty of water, juice or other fluids (e.g. 6 to 8 glasses per day).

Things you must not do

Do not take Curam Duo 500/125 to treat any other complaints unless your doctor tells you to.

Do not give this medicine to anyone else, even if they have the same condition as you.

Do not stop giving Curam Duo 500/125 or lower the dosage without checking with your doctor. If the full course prescribed by your doctor is not completed, all of the bacteria causing the infection may not be killed. These bacteria may continue to grow and multiply so that the infection may not clear completely or it may return.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while taking Curam Duo 500/125.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if he/she gets some of the side effects.

Do not be alarmed by the following lists of side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor or pharmacist if you or your child notices any of the following and they worry you:

- diarrhoea (several loose bowel movements per day), indigestion, pain in the stomach, feeling sick or being sick
- soreness of the mouth or tongue, abnormal taste, hairy tongue
- headache, dizziness, tiredness, hot flushes
- oral thrush (white, furry sore tongue and mouth)
- vaginal thrush (sore and itchy vagina, vaginal discharge)
- unusually active (hyperactive).

The above list includes the more common side effects of your medicine.

Tell your doctor as soon as possible if you or your child notices any of the following:

- nausea, vomiting, loss of appetite, generally feeling unwell
- itching or burning skin
- rash
- yellowing of the skin or eyes
- dark urine or pale stools
- severe abdominal cramps or stomach cramps
- severe diarrhoea
- unusual bleeding or bruising
- symptoms such as fever, severe chills, sore throat or mouth ulcers.

The above list includes serious side effects that may require medical attention. Serious side effects are rare.

Stop taking Curam Duo 500/125 if any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:

- wheezing, swelling of the lips, mouth, tongue or other parts of the body, difficulty in breathing or swallowing, hay fever, lumpy rash (hives) or fainting, joint discomfort or swelling, swollen lymph glands, nausea and vomiting. These could be symptoms of an allergic reaction.
- severe abdominal cramps or stomach cramps
- watery and severe diarrhoea, which may also be bloody
- fever, in combination with one or both of the above
- chest pain.

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation. These side effects are rare.

Rare events that have been reported with Curam Duo 500/125 include:

- inflammation of the bowel (colitis)
- inflammation of the liver or kidney
- blood disorders
- crystals in the urine (crystalluria)
- seizures, dizziness, headache
- change in colour of teeth
- change in colour of the tongue with a furry appearance
- flu like symptoms with a rash, fever, swollen glands and abnormal blood test results (including increased white blood cells (eosinophilia) and liver enzymes) (Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)).

Tell your doctor or pharmacist if you notice anything else that is making you feel unwell.

Other side effects not listed above may also occur in some people.

After giving Curam Duo 500/125

Storage

Keep this medicine in the original container.

If you take the tablets out of its original container it may not keep well.

Keep the medicine in a cool dry place where the temperature stays below 25°C.

Do not store Curam Duo or any other medicine in the bathroom or near a sink.

Do not leave it on a window sill or in the car. Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking the tablets, or the tablets have passed their expiry date, ask your pharmacist what to do with any that are left over.

Product description

What it looks like

Curam Duo 500/125 - Off-white, oval, biconvex, film-coated tablet, scored on both sides.

Available in strip packs or blister packs of 10 tablets.

Ingredients

Active ingredients:

- Curam Duo 500/125 tablets - 500mg amoxicillin (as trihydrate) and 125 mg clavulanic acid (as potassium clavulanate).

Inactive ingredients:

- magnesium stearate

- purified talc
- povidone
- croscarmellose sodium
- microcrystalline cellulose
- triethyl citrate
- ethylcellulose
- sodium lauryl sulfate
- cetyl alcohol
- hypromellose
- titanium dioxide

This medicine does not contain sucrose, gluten, tartrazine or any other azo dyes.

Supplier

Sandoz New Zealand Limited
12 Madden Street
Auckland 1010
New Zealand

Tel: 0800 726 369

This leaflet was revised in March 2023.

®: CURAM is a registered trademark