

Heparinised Saline Injection

CONSUMER MEDICINE INFORMATION

What is in this leaflet

This leaflet answers some common questions about Heparinised Saline Injection. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have benefits and risks. Your doctor has weighed the risks of you taking Heparinised Saline Injection against the benefits this medicine is expected to have for you.

This medicine is likely to be used while you are at the clinic or in hospital. If possible, please read this leaflet carefully before this medicine is given to you. In some cases this leaflet may be given to you after the medicine has been used.

If you have any concerns about taking this medicine, ask your doctor, nurse or pharmacist.

Keep this leaflet.

You may need to read it again.

This medicine is available only with a doctor's prescription.

What Heparinised Saline Injection is used for

Heparinised Saline Injection belongs to a group of medicines known as anticoagulants. Anticoagulants work by decreasing the clotting ability of your blood and help stop clots

forming in the blood vessels.

Anticoagulants are sometimes called blood thinners, although they do not actually thin the blood. Heparin will not dissolve blood clots that have already formed, but it may prevent any clots that have already formed from becoming larger and causing serious problems.

Heparinised Saline Injection is used to prevent the blocking of injection equipment often caused by blood clots.

Heparinised Saline Injection may be used for the treatment of other conditions that are not mentioned above. Your doctor will be able to tell you about the specific condition for which you have been prescribed Heparinised Saline Injection.

Before you are given Heparinised Saline Injection

When you must not be given it

Do not use Heparinised Saline Injection if you:

- have an allergy to heparin
- have, or may have, a bleeding disease or a problem with your blood vessels
- have low blood platelet count

If you are not sure whether any of these apply to you, check with your doctor.

Before you are given it

Tell your doctor if:

1. you have any allergies to:

- any other medicine
- any other substances, such as foods, preservatives or dyes

2. you are pregnant or intend to become pregnant

your doctor will discuss the possible risks and benefits of being given heparin during pregnancy.

3. you are breastfeeding or plan to breast feed

your doctor will discuss the possible risks and benefits of being given heparin whilst breastfeeding.

4. you have or have had any medical conditions, especially the following:

- heart problems or high blood pressure
- blood disease or bleeding problems
- heavy or unusual periods
- medical or dental surgery
- stomach ulcers
- liver disease

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and heparin may interfere with each other. These include:

- pain relieving medicines such as aspirin and ibuprofen
- medicines for heart conditions such as digitalis and dipyridamole
- antibiotics such as tetracycline and ciprofloxacin
- medicines for rheumatoid arthritis such as hydroxychloroquine
- anti-inflammatory medicines such as indomethacin and phenobutazone
- medicines for hay fever such as antihistamines
- nicotine

Some medicines and Heparinised Saline Injection may be incompatible in solution. These include:

- anticancer drugs such as doxorubicin and mitozantrone
- anticlotting medicines such as aprotinin
- sedatives such as diazepam
- antipsychotics such as some phenothiazines

Your doctor will advise you about continuing to take other medicines while you are receiving Heparinised Saline Injection.

How Heparinised Saline Injection is given

Heparinised Saline Injection is put into the injection device that is positioned in the arm and then it is injected into the vein. This will help to prevent blood clots from forming and blocking the device. Heparinised Saline Injection must only be given by a doctor or nurse.

How much is given

Your doctor will decide what dose, how often and how long you will receive Heparinised Saline Injection.

This depends on your condition and other factors, such as age, blood tests, method it is being given and whether or not other medicines are being given at the same time.

If you are given too much (overdose)

This rarely happens as Heparinised Saline Injection is administered under the care of a highly trained doctor.

However, if you are given too much heparin, you may experience some of the effects listed under "Side effects" below.

Your doctor has information on how to recognise and treat an overdose. Ask your doctor if you have any concerns.

In the case of an overdose, immediately tell your doctor or telephone the Poisons Information Centre (in Australia, phone 13 11 26; in New Zealand, phone 0800 POISON or 0800 764 766) for advice, or go to Accident and Emergency at your nearest hospital if you have side effects after being given Heparinised Saline Injection. You may need urgent medical attention.

Side effects

Like other medicines, heparin can cause some side effects. If they occur, most are likely to be minor or temporary. However, some may be serious and need medical attention.

Ask your doctor or nurse to answer any questions that you may have.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

Tell your doctor or nurse immediately if you experience any of the following:

- bleeding or bruising more easily than normal e.g. unexplained

nosebleeds, bleeding from gums when brushing teeth, red or dark brown urine, bloody or black stools

- signs of allergy such as a rash, itching, hives on the skin, swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or trouble breathing

These side effects are serious. You may need urgent medical attention

Tell your doctor or nurse as soon as possible if you notice the following:

- change in skin colour or pain around the injection site
- fever
- chills
- runny nose
- watering eyes
- vomiting
- nausea
- itchy soles of the feet

These are also side effects of heparin and are mostly mild and short-lived.

Other side effects not listed above may also occur in some patients. If you notice any other effects, check with your doctor. Some side effects may only be seen by your doctor.

Storage

Heparinised Saline Injection will be stored in the pharmacy or on the ward. The injection is kept in a cool dry place, where the temperature stays below 25°C.

Product Description

What it looks like

Heparinised Saline Injection is a clear, colourless solution in a plastic ampoule.

Heparinised Saline Injection can be identified by an Australian

Registration Number, which is found on the packaging:

AUST R 66684: Heparinised Saline Injection 50IU/5mL (sterile) Steriluer[®](50s)

Ingredients

Heparinised Saline Injection contains Heparin Sodium BP (porcine) 50IU/5mL, Sodium Chloride BP in Water for Injections BP. It does not contain a preservative.

Manufacturer/Sponsor

:

Manufacturer:

Pfizer (Perth) Pty Limited
ABN 32 051 824 956
15 Brodie Hall Drive
Bentley WA 6102 Australia

Sponsor and Distributor in Australia:

Pfizer Australia Pty Ltd
ABN 50 008 422 348
38-42 Wharf Road
West Ryde NSW 2114 Australia

Distributor in New Zealand:

Pfizer New Zealand Limited
PO Box 3998
Auckland New Zealand

This Consumer Medicine Information was written in March 2001.

Date of most recent amendment:
24 January 2011.