

RYTMONORM®

Propafenone Film coated tablet 150 mg



What is in this leaflet

Please read this leaflet carefully before you start taking RYTMONORM.

This leaflet answers some common questions about RYTMONORM.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking RYTMONORM against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine. You may need to read it again.

What RYTMONORM is used for

RYTMONORM is used to treat arrhythmias (abnormal heart rhythms).

RYTMONORM contains the active ingredient propafenone hydrochloride. It belongs to a group of medicines called anti-arrhythmic agents.

It works by slowing down the heart rate and helping to regulate the heartbeat.

Your doctor may have prescribed it for another reason.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

This medicine is available only with a doctor's prescription.

There is not enough information to recommend the use of this medicine for children.

Before you take RYTMONORM

When you must not take it

Do not take RYTMONORM if you have an allergy to:

- any medicine containing propafenone hydrochloride
- any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include: shortness of breath; wheezing or difficulty breathing; swelling of the face, lips, tongue or other parts of the body; rash, itching or hives on the skin.

Do not take RYTMONORM if you:

- have a heart condition called Brugada Syndrome
- have had a heart attack within the previous 3 months
- have or have had significant heart disease such as heart failure, severe symptomatic slow heart beat, changes in heart rate in the absence of an artificial pacemaker, severe low blood pressure or any heart problems other than abnormal heart rhythm
- have difficulty in breathing, such as asthma or chronic bronchitis or emphysema
- manifest electrolyte imbalance (e.g. potassium metabolism disorders)
- have severe lung disease
- have a condition known as myasthenia gravis (disease of the muscles causing

drooping eyelids, double vision, difficulty in speaking and swallowing and sometimes muscle weakness in the arms or legs)

- are currently taking ritonavir (a medicine used to treat HIV/AIDS)

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your doctor.

Before you start to take it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have had any of the following medical conditions:

- a heart condition called Brugada Syndrome
- change in heart rate
- difficulty of breathing that is not due to an allergy
- heart failure
- heart rhythm problems
- have a pacemaker
- liver problems
- kidney problems
- autoimmune disorders.

Tell your doctor or pharmacist if you are pregnant or plan to become pregnant or are breast-feeding.

Your doctor or pharmacist can discuss with you the risks and benefits involved.

If you have not told your doctor about any of the above, tell them

before you start taking RYTMONORM.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including medicines that you buy without a prescription from your pharmacy, supermarket or health food shop.

You should also tell any health professional who is prescribing a new medication for you that you are taking RYTMONORM.

Some medicines may interfere with RYTMONORM. These include:

- medicines that affect the activity of the heart (e.g. amiodarone, digoxin, quinidine)
- local anaesthetics such as lidocaine (e.g. used for surgery or dental work)
- medicines used to prevent blood clots (e.g. warfarin or phenprocoumon)
- antibiotics used to treat infections (e.g. erythromycin or rifampicin)
- any of the group of medicines known as beta-blockers, used to treat high blood pressure (e.g. propranolol, metoprolol)
- any antiviral agents (e.g. ritonavir)
- medicines used to treat depression (e.g. venlafaxine, fluoxetine, paroxetine, desipramine)
- medicine used to treat reflux and ulcers (e.g. cimetidine)
- medicine used after transplant operations, or in the treatment of arthritis or psoriasis (e.g. ciclosporin)
- medicines used to treat asthma (e.g. theophylline)
- medicines used to treat fungal infections (e.g. ketoconazole)
- medicines used to treat epilepsy (e.g. phenobarbital)
- grapefruit juice as it may interfere with the way this medicine works in your body.

These medicines may be affected by RYTMONORM or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

How to take RYTMONORM

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the box, ask your doctor or pharmacist for help.

How much to take

Follow your doctor's directions about when and how to take your tablets and look at the label on the carton. Your pharmacist will also help if you are not sure. The number of tablets that you will need to take will be decided by your doctor.

This may be:

- 1 - 2 film coated tablets of RYTMONORM 150 mg up to 3 times a day

You may need a lower dose of RYTMONORM if you are elderly, if you have problems with your kidneys or liver, or if you have a low bodyweight.

How to take it

Swallow the tablets whole, with a full glass of water. Do not take with grapefruit juice.

When to take it

Take your medicine at about the same time(s) each day.

Taking it at the same time(s) each day will have the best effect. It will also help you remember when to take it.

How long to take it

Continue taking your medicine for as long as your doctor tells you.

This medicine helps to control your condition, but does not cure it. It is important to keep taking your medicine even if you feel well.

If you forget to take it

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as soon as you remember, and then go back to taking your medicine as you would normally.

Do not take a double dose to make up for the dose that you missed.

This may increase the chance of you getting an unwanted side effect.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

While you are taking RYTMONORM

Things you must do

If you are about to be started on any new medicine, tell your doctor and pharmacist that you are taking RYTMONORM.

Tell any other doctors, dentists, and pharmacists who treat you that you are taking this medicine.

If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine. It may affect other medicines used during surgery.

If you become pregnant while taking this medicine, tell your doctor immediately.

If you are about to have any blood tests, tell your doctor that you are taking this medicine. It may interfere with the results of some tests.

Keep all of your doctor's appointments so that your progress can be checked.

Your doctor may do some tests (e.g. to check your heart function) from time to time to make sure the medicine is working and to prevent unwanted side effects.

Things you must not do

Do not take RYTMONORM to treat any other complaints unless your doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not stop taking your medicine or lower the dosage without checking with your doctor.

Things to be careful of

Be careful driving or operating machinery until you know how RYTMONORM affects you.

This medicine may cause blurred vision, dizziness, tiredness and low blood pressure in some people. If you have any of these symptoms, do not drive, operate machinery or do anything else that could be dangerous.

In case of overdose

If you take too much (overdose)

Immediately telephone your doctor or the National Poisons Centre (telephone 0800 POISON or 0800 764 766), or go to Accident and Emergency at your nearest hospital, if you think that you or anyone else may have taken too much RYTMONORM. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Symptoms of an overdose, may include:

- low blood pressure
- somnolence
- irregular heart beat
- abnormally fast heart beat
- heart attack
- abnormally fast or slow breathing
- headache

- blurred vision
- dizziness
- shaking
- nausea
- tingling or numbness of the hands and feet
- constipation
- dry mouth
- convulsions.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking RYTMONORM.

This medicine helps most people with irregular heartbeats, but it may have unwanted side effects in a few people.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- taste disturbance or loss of taste
- smell disturbance
- ringing in the ears
- vision disturbance
- dizziness, light-headedness
- fainting
- restlessness
- headache
- constipation
- nausea and/or vomiting
- abdominal pain
- diarrhoea
- flatulence
- heart palpitations (being aware of your heart beat)
- dry mouth
- sleep disorders
- nightmares
- confusion
- decreased or loss of appetite
- disorders of balance, walking, speech or movement
- tingling or numbness of hands or feet

- loss of memory
- changes in mental condition such as depression, anxiety or over-excitement
- hot flushes
- unusual hair loss or thinning (muscle) weakness
- inability to have or maintain an erection
- decreased sperm count
- signs of high blood glucose levels such as excessive thirst, increased appetite with weight loss, passing large amounts of urine.

The above list includes the less serious side effects of your medicine.

If any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:

- a rash, itching, swelling of the lips, face, tongue or other parts of the body, skin reddening, hives or other signs of an allergic reaction such as shortness of breath, wheezing or difficulty breathing.
- yellowing of the skin and/or eyes, light colored bowl motions or dark colored urine, as these may be a sign of liver problems
- unusual bleeding or bruising
- signs of infection such as sore throat, chills, mouth ulcers or fever
- tiredness, headaches, being short of breath when exercising, dizziness and looking pale
- convulsions, seizures or fits
- difficulty breathing
- coma
- unusually slow, fast or arrhythmic heart beat
- chest pain.

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation.

Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

Other side effects not listed above may also occur in some people. Tell

your doctor if you notice any other effects.

Some side effects (e.g. disturbance in your immune response, heart function, kidney function or liver function) can only be found when your doctor does tests from time to time to check your progress.

Do not be alarmed by these lists of possible side effects. You may not experience any of them.

After taking RYTMONORM

Storage

Keep your tablets in the pack until it is time to take them.

If you take the tablets out of the pack they may not keep well.

Keep your tablets in a cool dry place where the temperature stays below 25°C.

Do not store RYTMONORM or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car. Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Product description

What it looks like

RYTMONORM 150 mg

White to off-white, round, biconvex film coated tablets embossed on one face with "150". The other face is unmarked.

Ingredients

Active ingredient:
RYTMONORM tablets contain propafenone hydrochloride equivalent to 150 mg of propafenone as the active ingredient.

Inactive ingredients:

RYTMONORM also contains:

- microcrystalline cellulose
- starch - pregelatinised
- hypromellose
- sodium croscarmellose
- macrogol
- titanium dioxide
- magnesium stearate.

If you want to know more

Should you have any questions regarding this product, please contact your pharmacist or doctor.

Who supplies this medicine

RYTMONORM is supplied in New Zealand by:

Viatrix Ltd,
PO Box 11-183,
Ellerslie,
AUCKLAND
www.viatrix.co.nz
Telephone 0800 168 169

Date of Information

21 January 2022
(Based on datasheet dated 21 January 2022)

Rytmonorm® is a Viatrix company trade mark.