



New Zealand Data Sheet

APO-AMOXI

Amoxicillin Trihydrate equivalent to Amoxicillin 250mg and 500mg Capsules USP

Presentation

APO-AMOXI 250mg capsules are red and gold (size 2 capsules), imprinted 'APO 250'. Each capsule contains amoxicillin trihydrate equivalent to 250mg of amoxicillin, and typically weighs 376mg.

APO-AMOXI 500mg capsules are red and gold (size 0 capsules), imprinted 'APO 500'. Each capsule contains amoxicillin trihydrate equivalent to 500mg of amoxicillin, and typically weighs 735mg.

Uses

Actions

Amoxicillin is a bactericidal penicillin active against both Gram-positive and Gram-negative organisms and is subject to the hydrolytic activity of penicillinase. Amoxicillin differs *in vitro* from benzylpenicillin in the Gram-negative spectrum.

In vitro most strains of *Haemophilus influenzae*, *Neisseria gonorrhoeae*, *N. meningitidis*, *Proteus mirabilis*, *Amoxicillin*, Salmonellae, alpha- and beta-haemolytic Streptococci, *Diplococcus pneumoniae*, nonpenicillinase producing Staphylococci and *Streptococcus faecalis*, are sensitive to amoxicillin at serum concentrations which may be expected following the recommended doses. However, some of the organisms are sensitive to amoxicillin only at concentrations achieved in the urine (see Indications).

Escherichia coli isolates are becoming increasingly resistant to amoxicillin *in vitro* due to the presence of penicillinase producing strains.

Strains of gonococci that are relatively resistant to benzylpenicillin may be sensitive to amoxicillin.

Amoxicillin is not effective against penicillinase producing bacteria, particularly resistant Staphylococci, which now have a high prevalence.

All strains of *Pseudomonas*, *Klebsiella*, *Enterobacter*, indole positive *Proteus*, *Serratia marcescens*, *Citrobacter*, penicillinase producing *N. gonorrhoeae* and penicillinase producing *H. influenzae* are also resistant.

Combining Amoxicillin with a beta-lactamase inhibitor such as clavulanic acid effectively broadens its spectrum of activity to include many strains of beta-lactamase producing bacteria.

Like benzylpenicillin, amoxicillin is bactericidal against sensitive organisms during the stage of active multiplication. Amoxicillin exerts its mode of action by interfering with bacterial cell wall synthesis by acylating the enzyme transpeptidase, thus rendering it unable to cross-link muramic acid containing peptidoglycan strands. This inhibition of the biosynthesis of dipeptidoglycan, a substance necessary for cell wall strength and rigidity, results in a defective cell wall.

Pharmacokinetics

Amoxicillin is stable at gastric pH. It is completely absorbed in the upper gastrointestinal tract. Absorption is independent of meals.

APO-AMOXI

Amoxicillin Trihydrate equivalent to Amoxicillin 250mg and 500mg Capsules USP

Amoxicillin diffuses rapidly into most body tissues and fluids except for the brain and spinal fluid unless the meninges are inflamed. Amoxicillin has been shown to diffuse into saliva and sputum and is excreted mainly via the urine where it exists in high concentration. The amount to be found in bile is variable depending on normal biliary secretory function.

Amoxicillin is not highly protein bound, with only 17% protein bound in serum.

The serum half-life of amoxicillin is 61.3 minutes but in the absence of renal function it is 7 to 10 hours. The half-life may be longer in neonates and the elderly.

Oral doses of 250mg and 500mg of amoxicillin give average peak serum levels in one to two hours or 5micrograms/mL and 6.6 to 10.8micrograms /mL respectively. Detectable serum levels of amoxicillin are present 8 hours after the ingestion of a single dose.

Amoxicillin is primarily excreted by the renal system and approximately 75% of a 1g dose is excreted within 6 hours in the presence of normal renal function. It is excreted in the urine both unchanged (60%) and as penicilloic acid (15%). However, only about 32% of a 3g dose is excreted via the urine as the biologically active component in 8 hours (by which time most of the urinary excretion is complete). This proportional difference in the amount excreted from the different doses reflects a lack of linearity between doses and extent of absorption with a levelling off at higher doses or oral amoxicillin. Excretion of amoxicillin can be delayed by concurrent administration of probenecid thus prolonging its therapeutic effect.

Indications

Amoxicillin is indicated for the treatment of the following infections due to susceptible strains of sensitive organisms:

Upper Respiratory Infections:

Otitis media, pharyngitis, sinusitis and tonsillitis.

Lower Respiratory Infections:

Bronchitis, bronchopneumonia and lobar pneumonia.

Urinary Tract Infections:

Cystitis, cysto-pyelitis, urethritis, and gonococcal urethritis.

Prophylaxis:

against α -haemolytic (viridans group) and β -haemolytic Streptococci before dental, oral or upper respiratory tract surgery or instrumentation.

Prophylaxis:

of bacterial endocarditis in patients with any of the following conditions: congenital cardiac malformations, rheumatic and other acquired valvular lesions, prosthetic heart valves, previous history of bacterial endocarditis, hypertrophic cardiomyopathy, surgically constructed systemic-pulmonary shunts, mitral valve prolapse with valvular regurgitation or mitral valve prolapse without valvular regurgitation but associated with thickening and/or redundancy of the valve leaflets.

Amoxicillin is further indicated for the treatment of cutaneous infections. In emergency cases where the causative organism is not yet identified, therapy may be initiated with amoxicillin on the

APO-AMOXI

Amoxicillin Trihydrate equivalent to Amoxicillin 250mg and 500mg Capsules USP

basis of clinical judgement, while awaiting the results of bacteriologic studies to determine its antimicrobial sensitivity.

Dosage and Administration

Amoxicillin can be administered orally independent of meals. Therapy should be maintained for a minimum for 5 days. Treatment should be continued for a minimum of 96 hours beyond the time the patient becomes asymptomatic or after evidence of bacterial eradication is observed. At least 10 days treatment is recommended for any infection caused by beta-haemolytic streptococci to prevent the occurrence of acute rheumatic fever or glomerulonephritis.

Infections of the upper respiratory or genitourinary tracts, skin and soft tissues

due to susceptible strains of the causative organism:

Adults: 250 mg every 8 hours.

Children < 20 kg: 25 mg/kg/day in divided doses every 8 hours.

In severe infections

or infections associated with organisms where sensitivity determinations require higher blood concentrations:

Adults: 500 mg every 8 hours.

Children < 20 kg: 50 mg/kg/day in divided doses every 8 hours; this dosage should not exceed the recommended adult dosage.

Infections of the lower respiratory tract

due to susceptible strains of the causative organism, and acute otitis media:

Adults: 500 mg every 8 hours.

Children < 20 kg: 50 mg/kg/day in divided doses every 8 hours; this dosage should not exceed the recommended adult dosage.

High Dosage Therapy:

An adult dosage of 3g twice daily is recommended in appropriate cases for the treatment of severe or recurrent purulent infection of the respiratory tract.

Urethritis

due to non-penicillinase producing *N. Gonorrhoea* acquired in area with active monitoring for resistance to penicillin and where the percentage of penicillin-resistant isolates is <3.0%:

Adults and children > 45 kg: 3 g as a single oral dose; 1 g of oral probenecid should be administered concomitantly as well as appropriate therapy for presumptive or proven infection with *C. Trachomatis*.

Children < 45 kg: a single 50 mg/kg dose (maximum 3 g) of amoxicillin given with a single 25 mg/kg (up to 1 g) dose of probenecid. However, probenecid is not recommended in children under 2 years of age. Appropriate therapy of presumptive or proven infection with *C. Trachomatis* should be included as well.

Before prescribing amoxicillin, dark field examinations should be carried out in those cases where syphilis is also suspected and monthly serologic tests should be carried out for a minimum of 4 months.

Acute, uncomplicated lower urinary tract infections:

Adults: 3g as a single dose.

In the treatment of chronic urinary tract infections, frequent bacteriologic and clinical evaluations are necessary. Doses smaller than those recommended above should not be used. In stubborn

APO-AMOXI

Amoxicillin Trihydrate equivalent to Amoxicillin 250mg and 500mg Capsules USP

infections, therapy may be required for several weeks, sometimes at doses higher than those recommended. Concurrent bacteriologic sensitivity monitoring is recommended. It may be necessary to continue clinical and/or bacteriologic follow-up for several months after cessation of therapy.

For prevention of endocarditis:

Patient not having a general anaesthetic:

Adults: 3 g orally one hour before the procedure. A second dose may be given 6 hours later if considered necessary.

Children: For children under 5 administer quarter the adult dose and for children aged 5 to 10 administer half the adult dose.

Patient having a general anaesthetic – oral antibiotics appropriate:

Adults: 3g orally 4 hours prior to anaesthesia followed by 3g orally 6 hours after the initial dose.

Children: For children under 5 administer quarter the adult dose and for children aged 5 to 10 administer half the adult dose.

Note: The children's dose is intended for individuals whose weight will not cause dosage to be calculated greater than that recommended for adults. Children weighing more than 20 kg should be dosed according to the adult recommendations.

Dosage in renal failure:

The relative dose interval for amoxicillin is 4 hours; thus, in patients with renal failure in whom the half life is 6 hours the dosage interval is 24 hours and it may be necessary to reduce the total daily dosage. In patients with a creatinine clearance of more than 30mL/min no dosage adjustment is required. For patients with a creatinine clearance between 10 and 30mL/min the maximum recommended dose is 500mg twice daily. For patients with a creatinine clearance of less than 10mL/min the maximum recommended dose is 500mg per day. In patients receiving peritoneal dialysis the maximum recommended daily dose is 500mg. Amoxicillin may be removed from the system by haemodialysis.

Contraindications

Patients with a history of hypersensitivity to beta-lactum antibiotics (e.g. penicillins and cephalosporins).

Warnings and Precautions

Amoxicillin is not effective against beta-lactamase-producing (penicillin resistant) staphylococci.

Serious and occasionally fatal hypersensitivity (anaphylactoid) reactions have been reported in patients on parenteral and oral penicillin therapy. These reactions, however, are more likely to occur in individuals with a history of sensitivity to multiple allergens. Therefore, before initiating therapy with amoxicillin or any penicillin, careful inquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins or other allergens. If an allergic reaction occurs, amoxicillin should be discontinued and the appropriate measures instituted. Serious anaphylactoid reactions require immediate emergency treatment with epinephrine, oxygen, intravenous steroids and airway management, as indicated.

As with other penicillins, amoxicillin should be administered with caution in the presence of renal or hepatic insufficiency. Periodic assessment of renal, hepatic and haematopoietic functions should be made during prolonged amoxicillin therapy. Since amoxicillin is excreted mainly by the

APO-AMOXI

Amoxicillin Trihydrate equivalent to Amoxicillin 250mg and 500mg Capsules USP

kidney, the dosage for patients with renal impairment should be reduced when creatinine clearance is <30 mL/min (see Dosage and Administration).

The possibility of superinfections with mycotic organisms or bacterial pathogens should be kept in mind during amoxicillin therapy. If superinfections occur (usually involving *Aerobacter*, *Candida* or *Pseudomonas*), the drug should be discontinued and appropriate measures instituted.

Amoxicillin should not be used if infectious mononucleosis or lymphatic leukaemia is suspected since the occurrence of a morbilliform rash in patients with infectious mononucleosis following the use of amoxicillin is well documented. These patients are also very susceptible to ampicillin-induced skin rashes.

Following single dose therapy of acute lower urinary tract infections, the urine should be cultured. A positive culture may be evidence of a complicated or upper urinary tract infection and call for a longer or larger course of therapy.

Adequate fluid intake and urinary output must be maintained in patients receiving high doses of amoxicillin.

In patients with reduced urine output crystalluria has been observed very rarely, predominantly with parental therapy. During the administration of high doses of amoxicillin, it is advisable to maintain adequate fluid intake and urinary output in order to reduce the possibility of amoxicillin crystalluria.

Prolongation of prothrombin time has been reported rarely in patients receiving amoxicillin. Appropriate monitoring should be undertaken when anticoagulants are prescribed concurrently.

Use in Pregnancy

Category A.

Animal studies with amoxicillin have not shown any teratological effects. The product has been used extensively for more than twenty years and its suitability in human pregnancy has been well documented. When antibiotic therapy is required during pregnancy, amoxicillin may be considered appropriate.

Use in Lactation

Amoxicillin is distributed into breast milk in low concentrations and should be used with caution in nursing women.

Adverse Effects

The following convention has been utilised for the classification of undesirable effects:- Very common (more than 1/10), common (more than 1/100, less than 1/10), uncommon (more than 1/1000, less than 1/100), rare (more than 1/10,000, less than 1/1000), very rare (less than 1/10,000).

The majority of the side-effects listed below are not unique to amoxicillin and may occur when using other penicillins.

Unless otherwise stated, the frequency of adverse events (AE's) has been derived from more than 30 years of post-marketing reports.

Blood and lymphatic system disorders

APO-AMOXI

Amoxicillin Trihydrate equivalent to Amoxicillin 250mg and 500mg Capsules USP

Very rare: Reversible leucopenia (including severe neutropenia or agranulocytosis), reversible thrombocytopenia and haemolytic anaemia. Prolongation of bleeding time and prothrombin time.

Immune system disorders

Very rare: As with other antibiotics, severe allergic reactions, including angioneurotic oedema, anaphylaxis, serum sickness and hypersensitivity vasculitis. If a hypersensitivity reaction is reported, the treatment must be discontinued.

Nervous system disorders

Very rare: Hyperkinesia, dizziness, and convulsions. Convulsions may occur in patients with impaired renal function or in those receiving high doses.

Infections and Infestations

Very rare: Mucocutaneous candidiasis.

Gastrointestinal disorders

Common: Diarrhoea and nausea.

Uncommon: Vomiting.

Very rare: Antibiotic associated colitis (including pseudomembranous colitis and haemorrhagic colitis). Black hairy tongue. Superficial tooth discolouration has been reported in children. Good oral hygiene may help to prevent tooth discolouration as it can usually be removed by brushing.

Hepato-biliary disorders:

Very rare: Hepatitis and cholestatic jaundice. A moderate rise in AST and/or ALT. The significance of a rise in AST and/or ALT is unclear.

Skin and subcutaneous tissue disorders:

Common: Skin rash.

Uncommon: Urticaria and pruritus.

Very rare: Skin reactions such as erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis, bullous and exfoliative dermatitis and acute generalised exanthematous pustulosis (AGEP).

Renal and Urinary tract disorders:

Very rare: Interstitial nephritis, crystalluria.

The incidence of these AEs was derived from clinical studies involving a total of approximately 6,000 adults and paediatric patients taking amoxicillin.

Interactions

Probenecid decreases the renal tubular secretion of amoxicillin, and concurrent use may result in increased and prolonged blood levels of amoxicillin.

There is an increased risk of skin rashes in patients taking amoxicillin and allopurinol concurrently

Amoxicillin may reduce the efficiency of oral contraceptives and patients should be warned according.

Tetracyclines and other bacteriostatic drugs may interfere with the bactericidal effects of amoxicillin.



APO-AMOXI

Amoxicillin Trihydrate equivalent to Amoxicillin 250mg and 500mg Capsules USP

Overdosage

Serious toxicity is unlikely to occur. Acute ingestion of large doses may cause nausea, vomiting, diarrhoea and abdominal pain and should be treated symptomatically. Amoxicillin is removable from the circulation by haemodialysis. For all cases of suspected overdose appropriate supportive measures and symptomatic treatment are recommended.

Pharmaceutical Precautions

Store below 25°C.
Protect from heat, light and moisture.

Medicine Classification

Prescription-Only Medicine

Package Quantities

APO-AMOXI 250 mg capsules:
Bottles of 28, 100 and 500 capsules.
APO-AMOXI 500 mg capsules:
Bottles of 28, 100 and 500 capsules.

Further Information

Nil

Name and Address

Apotex NZ Ltd
32 Hillside Road
Glenfield, Auckland
Private Bag 102-995
North Shore Mail Centre
Tel: (09) 444-2073
Fax: (09) 444-2951

Date of Preparation

03 November 2011