



DIAMIDE

1. Product Name

Diamide 2 mg capsules

2. Qualitative and Quantitative Composition

Each capsule contains 2 mg of loperamide hydrochloride.

Diamide capsules contain lactose. For the full list of excipients, see section 6.1.

3. Pharmaceutical Form

Size 4 capsule, with mauve opaque body, dark green opaque cap, marked "LOPERA-MIDE 2" on the cap in white ink.

4. Clinical Particulars

4.1 *Therapeutic indications*

DIAMIDE is indicated for the symptomatic control of acute and chronic diarrhoea. In patients with an ileostomy it can be used to reduce the number and volume of stools and to harden their consistency.

4.2 *Dose and method of administration*

Dose

Adults and children over 12 years of age

Acute diarrhoea: the initial dose is 2 capsules; followed by 1 capsule after every subsequent loose stool.

Chronic diarrhoea: the initial dose is 2 capsules daily; this initial dose will be adjusted until 1-2 solid stools a day are obtained, which is usually achieved with a maintenance dose of 1-6 capsules daily.

The maximum dose for acute and chronic diarrhoea is 8 capsules daily.

4.3 *Contraindications*

DIAMIDE is contraindicated in patients with known hypersensitivity to loperamide or to any of the excipients (see section 6.1).

DIAMIDE should not be used as the primary therapy:

- in patients with acute dysentery, which is characterised by blood in stools and high fever;
- in patients with acute ulcerative colitis;

- in patients with bacterial enterocolitis caused by invasive organisms including *Salmonella*, *Shigella* and *Campylobacter*;
- in patients with pseudomembranous colitis associated with the use of broad-spectrum antibiotics.

In general, DIAMIDE should not be used when inhibition of peristalsis is to be avoided due to the possible risk of significant sequelae including ileus, megacolon and toxic megacolon. DIAMIDE must be discontinued promptly when constipation, abdominal distension or ileus develop.

Treatment of diarrhoea with DIAMIDE is only symptomatic. Whenever an underlying etiology can be determined, specific treatment should be given when appropriate.

Use in children

DIAMIDE is contraindicated in children under the age of 12 years.

4.4 Special warnings and precautions for use

Treatment of diarrhoea with DIAMIDE is only symptomatic. Whenever an underlying etiology can be determined, specific treatment should be given when appropriate. The priority in acute diarrhoea is the prevention or reversal of fluid and electrolyte depletion. This is particularly important in young children and in frail and elderly patients with acute diarrhoea. Use of this medicine does not preclude the administration of appropriate fluid and electrolyte replacement therapy.

Since persistent diarrhoea can be an indicator of potentially more serious conditions, this medicine should not be used for prolonged periods until the underlying cause of the diarrhoea has been investigated. In acute diarrhoea, if clinical improvement is not observed within 48 hours, the administration of DIAMIDE should be discontinued and patients should be advised to consult their physician.

Cardiac events

Cardiac events including QT prolongation and torsades de pointes have been reported in association with overdose. Some cases had a fatal outcome (see section 4.9). Patients should not exceed the recommended dose and/or the recommended duration of treatment.

Abuse and dependence

Physical dependence to loperamide in humans has not been observed. However, studies in monkeys demonstrated that loperamide at high dose produced symptoms of physical dependence of the morphine type.

Use in patients with hepatic impairment

Although no pharmacokinetic data are available in patients with hepatic impairment, DIAMIDE should be used with caution in such patients because of reduced first pass metabolism, as it may result in a relative overdose leading to central nervous system (CNS) toxicity.

Use in patients with renal impairment

Since the majority of the drug is metabolised, and the metabolites or the unchanged drug is excreted in the faeces, dose adjustments in patients with a kidney disorder are not required.

Use in patients with AIDS

Use with caution in patients with AIDS. Patients with AIDS treated with loperamide for diarrhoea should have therapy stopped at the earliest signs of abdominal distension. There have been isolated reports of obstipation with an increased risk for toxic megacolon in AIDS patients with infectious colitis from both viral and bacterial pathogens treated with loperamide.

Patients with lactose intolerance

Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine because it contains lactose.

Anticholinergic effects

In vitro studies have demonstrated anti-cholinergic properties. Hence, caution should be used in patients with glaucoma, urinary bladder neck obstruction, pyloric obstruction, significant gastric retention, or intestinal stasis.

4.5 Interaction with other medicines and other forms of interaction

Effect of loperamide on other drugs

Although the pharmacological effect of loperamide is not associated with a central action, patients with concomitant administration of tranquillisers or alcohol should be carefully observed.

Other drugs that affect loperamide

Consideration should always be given with new drugs as to possible interaction with monoamine oxidase inhibitors. Theoretically, the combination of loperamide with monoamine oxidase inhibitors, which are also inhibitors of liver microsomal enzymes, may potentiate the action of loperamide by blocking its metabolic pathway.

Non-clinical data have shown that loperamide is a P-glycoprotein substrate. Concomitant administration of loperamide (16 mg single dose) with quinidine, or ritonavir, which are both P-glycoprotein inhibitors, resulted in a 2 to 3 fold increase in loperamide plasma levels. The clinical relevance of this pharmacokinetic interaction with P-glycoprotein inhibitors, when loperamide is given at recommended dosages, is unknown.

The concomitant administration of loperamide (4 mg single dose) and itraconazole, an inhibitor of CYP3A4 and P-glycoprotein, resulted in a 3 to 4-fold increase in loperamide plasma concentrations. In the same study a CYP2C8 inhibitor, gemfibrozil, increased loperamide by approximately 2-fold. The combination of itraconazole and gemfibrozil resulted in a 4-fold increase in peak plasma levels of loperamide and a 13-fold increase in total plasma exposure. These increases were not associated with central nervous system (CNS) effects as measured by psychomotor tests (i.e. subjective drowsiness and the Digit Symbol Substitution Test).

The concomitant administration of loperamide (16 mg single dose) and ketoconazole, an inhibitor of CYP3A4 and P-glycoprotein, resulted in a 5-fold increase in loperamide plasma concentrations. This increase was not associated with increased pharmacodynamic effects as measured by pupillometry.

Concomitant treatment with oral desmopressin resulted in a 3-fold increase of desmopressin plasma concentrations, presumably due to slower gastrointestinal motility.

It is expected that drugs with similar pharmacological properties may potentiate loperamide's effect and that drugs that accelerate gastrointestinal transit may decrease its effect.

4.6 Fertility, pregnancy and lactation

Pregnancy

Safety in human pregnancy has not been established, although from animal studies there are no indications that loperamide possesses any teratogenic or embryotoxic properties. As with other drugs, it is not advisable to administer this medicine in pregnancy, especially during the first trimester.

Women who are pregnant should therefore be advised to consult their doctor for appropriate treatment.

Breast-feeding

Small amounts of loperamide may appear in human breast milk. Therefore, this medicine is not recommended during breastfeeding.

Women who are breastfeeding infants should therefore be advised to consult their doctor for appropriate treatment.

Fertility

No data available.

4.7 Effects on ability to drive and use machines

Loss of consciousness, depressed level of consciousness, tiredness, dizziness or drowsiness may occur when diarrhoea is treated with loperamide. Therefore, it is advisable to use caution when driving a car or operating machinery.

4.8 Undesirable effects

Adults and children aged ≥ 12 years

The safety of loperamide was evaluated in 2755 adults and children aged ≥ 12 years who participated in 26 controlled and uncontrolled clinical trials of loperamide used for the treatment of acute diarrhoea.

The most commonly reported (i.e. $\geq 1\%$ incidence) adverse drug reactions (ADRs) in clinical trials with loperamide in acute diarrhoea were: constipation (2.7%), flatulence (1.7%), headache (1.2%) and nausea (1.1%).

Table 1 displays ADRs that have been reported with the use of loperamide from either clinical trial (acute diarrhoea) or post-marketing experience.

The frequency categories use the following convention: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); and very rare ($< 1/10,000$).

Table 1: Adverse drug reactions

System organ class	Indication		
	Common	Uncommon	Rare
Immune system disorders			Hypersensitivity reaction ^a Anaphylactic reaction (including Anaphylactic shock) ^a Anaphylactoid reaction ^a
Nervous system disorders	Headache	Dizziness Somnolence ^a	Loss of consciousness ^a Stupor ^a Depressed level of consciousness ^a Hypertonia ^a Coordination abnormality ^a
Eye disorders			Miosis ^a
Gastrointestinal disorders	Constipation Nausea Flatulence	Abdominal pain Abdominal discomfort Dry mouth Abdominal pain upper Vomiting Dyspepsia ^a	Ileus ^a (including paralytic ileus) Megacolon ^a (including toxic megacolon ^b) Abdominal distension
Skin and subcutaneous tissue disorders		Rash	Bullous eruption ^a (including Stevens-Johnson syndrome, Toxic epidermal necrolysis and Erythema multiforme) Angioedema ^a Urticaria ^a Pruritus ^a
Renal and urinary disorders			Urinary retention ^a

General disorders and administration site conditions			Fatigue ^a
<p>a: Inclusion of this term is based on post-marketing reports for loperamide. As the process for determining post marketing ADRs did not differentiate between chronic and acute indications or adults and children, the frequency is estimated from all clinical trials with loperamide (acute and chronic), including trials in children ≤ 12 years (N=3683).</p> <p>b: See Warnings and Precautions.</p>			

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare professionals are asked to report any suspected adverse reactions <https://nzphvc.otago.ac.nz/reporting/>.

4.9 Overdose

Symptoms

In case of overdosage (including relative overdosage due to hepatic dysfunction), central nervous system depression (stupor, coordination abnormality, somnolence, miosis, muscular hypertonia, respiratory depression), constipation, urinary retention and paralytic ileus may occur. Children and patients with hepatic dysfunction may be more sensitive to CNS effects.

In individuals who have ingested overdoses of loperamide HCl, cardiac events such as QT interval prolongation, torsades de pointes, other serious ventricular arrhythmias, cardiac arrest and syncope have been observed (see section 4.4). Fatal cases have also been reported.

In clinical trials using loperamide, an adult took three 20 mg doses within a 24-hour period, was nauseated after the second, and vomited after the third dose.

Treatment

If vomiting has occurred spontaneously, a slurry of 100 g of activated charcoal should be administered orally as soon as fluids can be maintained.

If vomiting has not occurred, gastric lavage should be performed, followed by administration of 100 g of activated charcoal slurry through gastric tube. In the case of overdosage, patient should be monitored for signs of CNS depression and/or respiratory depression for at least 24 hours. If CNS depression is observed, naloxone may be administered. If responsive to naloxone, vital signs must be monitored carefully for recurrence of symptoms of drug overdosage for at least 24 hours after the last dose of naloxone. In view of the prolonged action of loperamide and the short duration (one to three hours) of naloxone, the patient must be monitored closely and treated repeatedly with naloxone as indicated. Based on the fact that relatively little loperamide is excreted in urine, forced diuresis is not expected to be effective for loperamide overdosage.

For further advice on management of overdose please contact the National Poisons Information Centre (0800 POISON or 0800 764 766).

5. Pharmacological Properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic Group: Antipropulsives: ACT code: A07DA03.

Mechanism of action

Loperamide binds to the opiate receptor in the gut wall, reducing propulsive peristalsis, increasing intestinal transit time and enhancing resorption of water and electrolytes. Loperamide increases the tone of the anal sphincter, which helps reduce faecal incontinence and urgency.

Clinical efficacy and safety

In a double blind randomised clinical trial in 56 patients with acute diarrhoea receiving loperamide, onset of anti-diarrhoeal action was observed within one hour following a single 4 mg dose. Clinical comparisons with other antidiarrhoeal drugs confirmed this exceptionally rapid onset of action of loperamide.

Cardiovascular effects

In human volunteers, analysis of electrocardiograms obtained pre-therapy, and then two and six hours after administration of loperamide (16 mg), revealed no evidence of cardiovascular toxicity.

5.2 Pharmacokinetic properties

Absorption

Most ingested loperamide is absorbed from the gut, but as a result of significant first pass metabolism, systemic bioavailability is only approximately 0.3%.

Distribution

Studies on distribution in rats show a high affinity for the gut wall with a preference for binding to receptors of the longitudinal muscle layer. The plasma protein binding of loperamide is 95%, mainly to albumin. Non-clinical data have shown that loperamide is a P-glycoprotein substrate.

Biotransformation

Loperamide is almost completely extracted by the liver, where it is predominantly metabolized, conjugated and excreted via the bile. Oxidative N-demethylation is the main metabolic pathway for loperamide, and is mediated mainly through CYP3A4 and CYP2C8. Due to this very high first pass effect, plasma concentrations of unchanged drug remain extremely low.

Elimination

The half-life of loperamide in man is about 11 hours with a range of 9-14 hours. Excretion of the unchanged loperamide and the metabolites mainly occurs through the faeces.

5.3 Preclinical safety data

Acute and chronic studies on loperamide showed no specific toxicity. Results of *in vivo* and *in vitro* studies carried out indicated that loperamide is not genotoxic. In reproduction studies, very high doses (40 mg/kg/day - 240 times the maximum human use level) loperamide impaired fertility and foetal survival in association with maternal toxicity in rats. Lower doses had no effects on maternal or foetal health and did not affect peri- and post-natal development.

Non-clinical *in vitro* and *in vivo* evaluation of loperamide indicates no significant cardiac electrophysiological effects within its therapeutically relevant concentration range and at significant multiples of this range (up to 47-fold). However, at extremely high concentrations associated with overdoses (see section 4.4), loperamide has cardiac electrophysiological actions consisting of inhibition of potassium (hERG) and sodium currents, and arrhythmias.

CNS activity

Animal studies indicate that loperamide is devoid of analgesic properties (2-16 mg/kg). Studies in morphine-dependent monkeys demonstrated that loperamide in high S.C. doses prevented signs of morphine withdrawal. However, in humans the naloxone challenge pupil test which when positive indicated opiate-like effects, was negative when performed after a single high dose or after more than two years of therapeutic use (mean dose 4 mg/day) of loperamide.

6. Pharmaceutical Particulars

6.1 List of excipients

Lactose, magnesium stearate, maize starch. The capsule shell contains gelatin, titanium dioxide (E171), erythrosine (E127), indigo carmine (E132), black iron oxide (E172) quinoline yellow (E104), and printing ink.

Diamide does not contain gluten.

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

5 years.

6.4 Special precautions for storage

Store at or below 25°C.

6.5 Nature and contents of container

Blister strip pack containing 10 or 20 capsules.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

Not applicable.

7. Medicines Schedule

Pharmacy only Medicine

8. Sponsor Details

Mylan New Zealand Ltd
PO Box 11183
Ellerslie
AUCKLAND
Telephone 09-579-2792

9. Date of First Approval

16 September 2004

10. Date of Revision of the Text

27 July 2017 Added information on serious cardiac events with high doses to sections 4.4, 4.9 and 5.3.