

VELCADE[®]

DATA SHEET

NAME OF THE DRUG

Bortezomib


DESCRIPTION

VELCADE (bortezomib) is an antineoplastic agent for intravenous injection (IV) use only. Each single dose vial contains 1.0 mg or 3.5mg of bortezomib as a sterile lyophilised powder. After reconstitution, 1 mL of solution contains 1 mg bortezomib. Inactive ingredients: 35mg mannitol and nitrogen qs.

Bortezomib is a modified dipeptidyl boronic acid. The product is provided as a mannitol boronic ester which, in reconstituted form, consists of the mannitol ester in equilibrium with its hydrolysis product, the monomeric boronic acid.

The chemical name for bortezomib, the monomeric boronic acid, is [(1R)-3-methyl-1-[[[(2S)-1-oxo-3-phenyl-2-[(pyrazinylcarbonyl)amino]propyl]amino]butyl] boronic acid.

Bortezomib has the following chemical structure:



$C_{19}H_{25}BN_4O_4$

MW: 384.24

CAS Registry No. 179324-69-7

The solubility of bortezomib, as the monomeric boronic acid, in water is: 3.3 – 3.8 mg/mL in a pH range of 2 – 6.5.

PHARMACOLOGY

Pharmacodynamics

Mechanism of Action

Bortezomib is a reversible inhibitor of the chymotrypsin-like activity of the 26S proteasome in mammalian cells. The 26S proteasome is a large protein complex that degrades ubiquitinated proteins. The ubiquitin-proteasome pathway plays an essential role in regulating the intracellular concentration of specific proteins, thereby maintaining homeostasis within cells. Inhibition of the 26S proteasome prevents this targeted proteolysis which can affect multiple signalling cascades within the cell. This disruption of normal homeostatic mechanisms can lead to cell death. Experiments have demonstrated that bortezomib is cytotoxic to a variety of cancer cell types *in vitro*. Bortezomib causes a delay in tumour growth *in vivo* in nonclinical tumour models, including multiple myeloma.

Data from *in vitro*, *ex-vivo*, and animal models with bortezomib suggest that it increases osteoblast differentiation and activity and inhibits osteoclast function. These effects have been observed in patients with multiple myeloma affected by an advanced osteolytic disease and treated with bortezomib.

Pharmacokinetics Following intravenous bolus administration of a 1.0 mg/m² and 1.3 mg/m² dose to eleven patients with multiple myeloma, the mean first-dose maximum plasma concentrations of bortezomib were 57 and 112 ng/mL, respectively. In subsequent doses, mean maximum observed plasma concentrations ranged from 67 to 106ng/mL for the 1.0mg/m² dose and 89 to 120ng/mL for the 1.3mg/m² dose. The mean elimination half-life of bortezomib upon multiple dosing ranged from 40-193 hours. Bortezomib is eliminated more rapidly following the first dose compared to subsequent doses. Mean total body clearances were 102 and 112 L/h following the first dose for doses of 1.0mg/m² and 1.3mg/m², respectively, and ranged from 15 to 32 L/h following subsequent doses of 1.0mg/m² and 1.3mg/m², respectively. The mean distribution volume of bortezomib ranged from 1659 litres to 3294 litres (489 to 1884L/m²) following single- or repeat-dose administration of 1.0mg/m² or 1.3mg/m² to patients with multiple myeloma. This suggests that bortezomib distributes widely to peripheral tissues.

Protein Binding: Over a bortezomib concentration range of 10 to 1000 ng/mL, the *in vitro* protein binding averaged 83% in human plasma. The percent of bortezomib bound to plasma proteins was not concentration dependent.

Metabolism: *In vitro* studies with human liver microsomes and human cDNA-expressed cytochrome P450 isozymes indicate that bortezomib is primarily oxidatively metabolised via cytochrome P450 enzymes, 3A4, 2C19, 2D6, 2C9, and 1A2. The major metabolic pathway is deboronation, with the two main metabolites formed undergoing subsequent hydroxylation. One of the two main deboronated metabolites was shown to be inactive as a 26S proteasome inhibitor. Pooled plasma data from 8 patients at 10 min and 30 min after dosing indicate that the plasma levels of metabolites are low compared to the parent drug.

Elimination: The elimination pathways of bortezomib have not been evaluated *in vivo*.

Renal Impairment

A pharmacokinetic study was conducted in patients with various degrees of renal impairment who were classified according to their creatinine clearance values (CrCL) into the following groups: Normal (CrCL \geq 60 mL/min/1.73 m², n=12), Mild (CrCL=40-59 mL/min/1.73 m², n=10), Moderate (CrCL=20-39 mL/min/1.73 m², n=9), and Severe (CrCL < 20 mL/min/1.73 m², n=3). A group of dialysis patients who were dosed after dialysis was also included in the study (n=8). Patients were administered intravenous doses of 0.7 to 1.3 mg/m² of bortezomib twice weekly. Exposure of bortezomib (dose-normalized AUC and C_{max}) was comparable among all the groups. (see **DOSAGE AND ADMINISTRATION**)

Hepatic Impairment: The effect of hepatic impairment (see Table 13 for definition of hepatic impairment) on the pharmacokinetics of bortezomib was assessed in 51 cancer patients at bortezomib doses ranging 0.5 to 1.3 mg/m². When compared to patients with normal hepatic function, mild hepatic impairment did not alter dose-randomisation bortezomib AUC. However, the dose-randomised mean AUC values were increased by approximately 60% in patients with moderate or severe hepatic impairment. A lower starting dose is recommended in patients with moderate or severe hepatic impairment, and those patients should be monitored closely (see **Table 13**).

CLINICAL TRIALS

All response and progression data listed below for both previously untreated multiple myeloma in non-transplant eligible patients and relapsed / refractory multiple myeloma were assessed using the European Group for Blood and Marrow Transplantation (EBMT) criteria. The response and progression data for transplant-eligible multiple myeloma patients were assessed using the International Myeloma Working Group (IMWG) criteria.

Previously Untreated Multiple Myeloma

Transplant Eligible

The safety and efficacy of VELCADE, as induction therapy prior to stem cell transplantation in previously untreated multiple myeloma patients, has been assessed in multiple Phase III and Phase II trials.

A Phase III, randomised (1:1), open-label, multi-centre study conducted by the Italian Myeloma Network - GIMEMA, randomised 480 transplant-eligible patients under the age of 65 to receive three 3-week cycles of VELCADE (1.3 mg/m², days 1, 4, 8, 11) in combination with thalidomide (100 mg, days 1-14 in cycle 1, then 200 mg daily) and dexamethasone (40 mg, days 1, 2, 4, 5, 8, 9, 11, 12) (Vc-TD), or thalidomide and dexamethasone (TD) prior to tandem autologous transplant. Three months following transplant, patients received two cycles of consolidation treatment; patients randomized to receive Vc-TD induction received two 35-day cycles of VELCADE (1.3 mg/m², days 1, 8, 15, 22), thalidomide (100 mg daily) and dexamethasone (40 mg, days 1, 2, 8, 9, 15, 16, 22, 23) consolidation; patients randomized to receive thalidomide-dexamethasone induction received two 35-day cycles of thalidomide-dexamethasone consolidation. The primary endpoint of the study was response rate \geq nCR following induction therapy.

Patients randomized to Vc-TD arm achieved significantly higher rates of complete plus near complete response and very good partial response or better, compared to the thalidomide-dexamethasone arm following induction treatment. This difference was maintained following both transplant and consolidation therapy. Response rates are presented in Table 1.

Table 1: Summary of Response Rates by IMWG criteria in the GIMEMA study

Response Rate n (%)	Vc-TD n=236	TD n=238	p-value
Post-induction Therapy*			
CR	44 (19)	11 (5)	<0.0001
CR+nCR**	73 (31)	27 (11)	<0.0001
≥VGPR	146 (62)	66 (28)	<0.0001
≥PR	220 (93)	187 (79)	<0.0001
MR/SD	16 (7)	39 (16)	0.0011
PD	0	12 (5)	0.0005
Post-first ASCT			
CR	89 (38)	54 (23)	0.0004
CR+nCR	123 (52)	74 (31)	<0.0001
≥VGPR	186 (79)	137 (58)	<0.0001
≥PR	220 (93)	201 (84)	0.0025
MR/SD	15 (6)	20 (8)	0.3941
PD	1 (0)	17 (7)	0.0001
Post-second ASCT			
CR	98 (42)	72 (30)	0.0105
CR+nCR	130 (55)	98 (41)	0.0024
≥VGPR	193 (82)	152 (64)	<0.0001
≥PR	220 (93)	199 (84)	0.0011
MR/SD	14 (6)	19 (8)	0.3804
PD	2 (1)	20 (8)	0.0001
Post-consolidation			
CR	116 (49)	82 (35)	0.0012
CR+nCR	147 (62)	108 (45)	0.0002
≥VGPR	201 (85)	162 (68)	<0.0001
≥PR	218 (92)	201 (84)	0.0071
MR/SD	12 (5)	16 (7)	0.4495
PD	6 (3)	21 (9)	0.0032
Best overall response			
CR	136 (58)	97 (41)	0.0001
CR+nCR	168 (71)	128 (54)	<0.0001
≥VGPR	210 (89)	175 (73.5)	<0.0001
≥PR	227 (96)	212 (89)	0.0074


* Similar differences in post-induction response rates were reported by study investigators (CR+nCR: 32% vs. 13%, $p<0.0001$). Differences in RR following transplantation and consolidation by investigator assessment were also similar to those centrally assessed.

** These significant differences in CR+nCR rates between arms were maintained following cyclophosphamide to collect peripheral blood stem cells (42% vs 21%, $p<0.0001$).


ASCT: autologous stem cell transplantation; CR: complete response; MR: minimal response; nCR: near-complete response; PD: progressive disease; PR: partial disease; SD: stable disease; TD = thalidomide-dexamethasone; VGPR: very good partial response; Vc-TD: VELCADE-thalidomide-dexamethasone

In addition, compared with the TD arm, Progression Free Survival (PFS) was also significantly longer for patients randomized to the Vc-TD arm (HR, 0.629 [CI: 0.451-0.878], $p=0.0061$). The estimated 3-year PFS rate was 68% in the VTD arm and 56% in TD ($p=0.0057$) (see Figure 1). 58 (24.5%) and 86 (36%) patients progressed or died, respectively. The estimated 3-year probability of progression or relapse was 29% in the Vc-TD versus 39% in the TD arm (HR, 0.609 [CI: 0.425-0.873], $p=0.0073$; $p=0.0061$ by Kaplan-Meier analysis) (see Figure 2).

**Figure 1: Progression-Free Survival
Study GIMEMA: All Randomised Subjects Analysis Set)**



**Figure 2: Time to Disease Progression
(Study GIMEMA: All Randomised Subjects Analysis Set)**



The IFM-2005, Phase III, randomised (1:1:1:1), multi-centre, open-label study was conducted to compare the efficacy and safety of VELCADE-dexamethasone (Vc-Dex) and vincristine-doxorubicin-dexamethasone (VAD) as induction therapy prior to HDT-ASCT, and to evaluate the impact of post-induction consolidation therapy. Patients in this study were randomised to receive VAD plus no consolidation (arm A1), VAD plus dexamethasone, cyclophosphamide, etoposide, cis-platin (DCEP) consolidation (arm A2), Vc-Dex plus no consolidation (arm B1), or Vc-Dex plus DCEP consolidation (arm B2).

A total of 482 patients aged ≤65 years were randomised; 240 patients received four 3-week cycles of VELCADE (1.3 mg/m²), days 1, 4, 8 and 11 plus dexamethasone (40 mg) days 1-4 (all cycles) and days 9-12 (cycles 1 and 2), while 242 patients received four 4-week cycles of VAD. The primary endpoint of this study was the CR/nCR rate post-induction.

Patients randomized to the Vc-Dex arm achieved significantly higher rates of complete plus near complete response and very good partial response or better, compared to the VAD arm following induction treatment. Based on an intention to treat analysis, response rates were similar regardless of whether patients received DCEP consolidation or not. Efficacy results are presented in Table 2:

Table 2: Response to induction therapy (overall) in the IFM2005 study*

	VAD (A1+A2) N=242	Vc-Dex (B1+B2) N=240	p-value
Evaluable population, N	218	223	
ORR (≥PR), n (%)	137 (62.8)	175 (78.5)	<0.001
≥VGPR	33 (15.1)	84 (37.7)	<0.001
CR/nCR	14 (6.4)	33 (14.8)	0.004
CR	3 (1.4)	13 (5.8)	0.012
MR+SD	58 (26.6)	28 (12.6)	
PD	9 (4.1)	10 (4.5)	
Death	6 (2.8)	1 (0.5)	
Not assessable	8 (3.7)	9 (4.0)	

A total of 184/218 (84.4%) and 197/223 (88.3%) evaluable patients who received VAD and Vc-Dex induction, respectively, underwent autologous stem cell transplantation. The number of patients who received a second transplantation was 41 (20.8%) in the Vc-Dex arm, compared to 50 (27.2%) for patients in the VAD arm. Post-transplant response rates are shown in Table 3.

Table 3: Response rates post-transplantation*

	VAD (A1+A2) N=218	Vc-Dex (B1+B2) N=223	p-value
Response to first transplant			
ORR (≥PR), n (%)	168 (77.1)	179 (80.3)	0.401
≥VGPR	81 (37.2)	121 (54.3)	<0.001
CR/nCR	40 (18.4)	78 (35.0)	<0.001
CR	19 (8.7)	36 (16.1)	0.016
MR+SD+PD	8 (3.7)	6 (2.7)	
Death	2 (0.9)	1 (0.5)	
No transplantation	34 (15.6)	26 (11.7)	
Overall, including second transplantation			
≥VGPR	102 (46.7)	151 (67.7)	<0.001
CR/nCR	49 (22.5)	88 (39.5)	<0.001

* All response assessments were confirmed by an Independent Review Committee.

CR: complete response; MR: minimal response; nCR: near-complete response; ORR: overall response rate; PD: progressive disease; PR: partial response; SD: stable disease; VGPR: very good partial response.

In addition, the median PFS was 29.7 months among patients who received VAD versus 36.0 months among patients who received Vc-Dex induction, with 128 (52.9%) of 242 and 110 (45.8%) of 240 patients, respectively, having progressed ($p < .064$, or $p < .057$ if adjusted for initial stratification factors) after median follow-up of 31.2 months.

The efficacy and safety of VELCADE as induction therapy in newly diagnosed multiple myeloma patients were also assessed in various Phase I/II open-label studies. The results from these studies showed that the addition of VELCADE to the various combination chemotherapy regimens resulted in an overall response rates between 66% and 100%.

Non-Transplant Eligible

The VISTA study is a prospective phase III, international, randomized (1:1), open-label clinical study of 682 patients, conducted to determine whether VELCADE (1.3 mg/m²) in combination with melphalan (9 mg/m²) and prednisone (60 mg/m²) resulted in improvement in time to progression (TTP) when compared to melphalan (9 mg/m²) and prednisone (60 mg/m²) in patients with previously untreated multiple myeloma. Treatment was administered for a maximum of 9 cycles (approximately 54 weeks) and was discontinued early for disease progression or unacceptable toxicity. Baseline demographics and patient characteristics are summarized in Table 4.

Table 4: Summary of Baseline Patient and Disease Characteristics in the VISTA Study

Patient Characteristics	VcMP N=344	MP N=338
Median age in years (range)	71.0 (57, 90)	71.0 (48, 91)
Gender: male/female	51% / 49%	49% / 51%
Race: Caucasian/asian/black/other	88% / 10% / 1% / 1%	87% / 11% / 2% / 0%
Karnofsky performance status score ≤70	35%	33%
Hemoglobin <100 g/L	37%	36%
Platelet count <75 x 10 ⁹ /L	<1%	1%
Disease Characteristics		
Type of myeloma (%): IgG/IgA/Light chain	64% / 24% / 8%	62% / 26% / 8%
Median β ₂ -microglobulin (mg/L)	4.2	4.3
Median albumin (g/L)	33.0	33.0
Creatinine clearance ≤30 mL/min [n (%)]	20 (6%)	16 (5%)

VcMP = VELCADE + melphalan + prednisone; MP = melphalan + prednisone

At the time of a pre-specified interim analysis, the primary endpoint, time to progression, was met and patients in the MP arm were offered VcMP treatment. Median follow-up was 16.3 months. A statistically significant survival benefit in favour of the VcMP treatment group was observed (HR=0.65; *p*=0.00084) despite subsequent therapies that included VELCADE based regimens. While the median survival in MP treatment group has now been estimated at 43.1 months, the median survival on the VcMP treatment group has not been reached. Efficacy results are presented in Table 5 and Figures 3 and 4.

Table 5: Summary of Efficacy Analyses in the VISTA study

Efficacy Endpoint	VcMP n=344	MP n=338
Time to Progression – Events n (%)	101 (29)	152 (45)
Median ^a (95% CI)	20.7 mo (17.6, 24.7)	15.0 mo (14.1, 17.9)
Hazard ratio ^b (95% CI)	0.54 (0.42, 0.70)	
p-value ^c	0.000002	
Progression-free Survival Events n (%)	135 (39)	190 (56)
Median ^a (95% CI)	18.3 mo (16.6, 21.7)	14.0 mo (11.1, 15.0)
Hazard ratio ^b (95% CI)	0.61 (0.49, 0.76)	
p-value ^c	0.00001	
Overall Survival Events (deaths) n (%)	45 (13)	76 (23)
Hazard ratio ^b (95% CI)	0.61 (0.42, 0.88)	
p-value ^c	0.00782	
Response Rate population ^e n = 668	n=337	n=331
CR ^f n (%)	102 (30)	12 (4)
PR ^f n (%)	136 (40)	103 (31)
nCR n (%)	5 (1)	0
CR + PR ^f n (%)	238 (71)	115 (35)
p-value ^d	<10 ⁻¹⁰	
Reduction in Serum M-protein population ^g n=667	n=336	n=331
>=90% n (%)	151 (45)	34 (10)
Time to First Response in CR + PR		
Median	1.4 mo	4.2 mo
Median^a Response Duration		
CR ^f	24.0 mo	12.8 mo
CR + PR ^f	19.9 mo	13.1 mo
Time to Next Therapy Events n (%)	73 (21)	127 (38)
Median ^a (95% CI)	NE (26.1, NE)	20.8 mo (18.3, 28.5)
Hazard ratio ^b (95% CI)	0.52 (0.39, 0.70)	
p-value ^c	0.000009	

^a Kaplan-Meier estimate.

^b Hazard ratio estimate is based on a Cox proportional-hazard model adjusted for stratification factors: beta2-microglobulin, albumin, and region. A hazard ratio less than 1 indicates an advantage for VMP

^c p-value based on the stratified log-rank test adjusted for stratification factors: beta2-microglobulin, albumin, and region

^d p-value for Response Rate (CR + PR) from the Cochran-Mantel-Haenszel chi-square test adjusted for the stratification factors

^e Response population includes patients who had measurable disease at baseline


^f EBMT criteria

^g All randomized patients with secretory disease

NE: Not estimable


The time to progression (TTP) was significantly longer on the VELCADE arm (see Figure 3)

Figure 3: Time to Disease Progression
(Study 26866138-MMY-3002 Update: All Randomised Subjects Analysis Set)



A significant survival advantage is shown with VELCADE (see Figure 4)

Figure 4: Overall Survival
(Study 26866138-MMY-3002 Update: All Randomised Subjects Analysis Set)



Relapsed / Refractory Multiple Myeloma

The safety and efficacy of VELCADE were evaluated in 2 studies at the recommended dose of 1.3 mg/m²: The APEX study - a phase III randomised, stratified, open-label, comparative study, versus Dexamethasone (Dex), of 669 patients with relapsed or refractory multiple myeloma who had received 1-3 prior lines of therapy, and a phase II single-arm study of 202 patients with relapsed and refractory multiple myeloma, who had received at least 2 prior lines of treatment and who were progressing on their most recent treatment (see Tables 6 and 7).

Table 6: Dosing regimens in the APEX and Phase II studies

Phase/arm	Drug Schedule	Dose	Regimen
II	VELCADE: Day 1,4,8,11, (rest Day 12-21)	1.3 mg/m ² (IV bolus)	Q3 weeks x 8 cycles (extension**)
III (APEX)	VELCADE* a) Days 1,4,8,11, (Rest Day 12-21) b) Days 1,8,15,22 (Rest Day 23-35)	1.3 mg/m ² (IV bolus)	a) Q3 weeks x 8, then b) Q5 weeks x 3
III (APEX)	DEXAMETHASONE a) Days 1-4, 9-12, 17-20 Days 1-4	40 mg (PO)	a) Q5 weeks x 4 b) Q4 weeks x 5
II	Add DEXAMETHASONE***	20 mg (PO) (Days 1,2,4,5,8,9, 11,12)	Q3 weeks

* a) is the initial treatment, a) and b) represent a full course of treatment

** An extension study authorised patients benefiting from treatment to continue receiving VELCADE

*** If after 2 or 4 cycles of VELCADE, the patients had progressive disease or stable disease, respectively, they could receive dexamethasone

Table 7: Patient characteristics in the Phase II* and APEX studies

	Phase II study VELCADE N=202	APEX study VELCADE N=333	APEX study DEX. N=336
Patient characteristics			
Median age in years (range)	59(34-84)	62.0 (33-84)	61.0 (27-86)
Gender: male/female	60% / 40%	56% / 44%	60% / 40%
Karnofsky Performance Status score ≤ 70	20%	13%	17%
Haemoglobin <100 g/L	44%	32%	28%
Platelet count <75 x 10 ⁹ /L	21%	6%	4%
Disease Characteristics			
Type of myeloma (%): IgG/IgA/Light chain	60%/24%/14%	60%/23%/12%	59%/24%/13%
Median β2-microglobulin (mg/L)	3.5	3.7	3.6
Median creatinine clearance (mL/min)	73.9	73.3	75.3
Abnormal cytogenetics	35%		
Chromosome 13 abnormalities	15%	25.7%	25.0%
Median Duration of Multiple Myeloma Since Diagnosis in Years	4.0	3.5	3.1
Previous Therapy			
Number of Prior Therapeutic Lines of Treatment			
Median (range)**	6 (2-15)	2 (1-7)	2 (1-8)
1 prior line	0	40%	35%
>1 prior line		60%	65%
All patients			
Any prior steroids, e.g., dexamethasone, VAD	99%	98%	99%
Any prior alkylating agents, e.g., MP, VBMCP	92%	91%	92%
Any prior anthracyclines, e.g., VAD, mitoxantrone	81%	77%	76%
Any prior thalidomide therapy	83%	48%	50%
Any prior stem cell transplant/other high-dose therapy	64%	67%	68%
Prior experimental or other types of therapy	44%	3%	2%
*Based on number of patients with baseline data available			
**Including steroids, alkylating agents, anthracyclines, thalidomide and stem cell transplants			

APEX Study (Phase III)

In the APEX study described above, patients considered to be refractory to prior high-dose dexamethasone were excluded as were those with baseline grade ≥ 2 peripheral neuropathy or platelet counts $<50,000/\mu\text{L}$. A total of 627 patients were evaluable for response. Stratification factors were based on the number of lines of prior therapy the patient had previously received (1 previous line versus more than 1 line of therapy), time of progression relative to prior treatment (progression during or within 6 months of stopping their most recent therapy versus relapse >6 months after receiving their most recent therapy), and screening β_2 -microglobulin levels (≤ 2.5 mg/L versus >2.5 mg/L).

Following a preplanned interim analysis of time to progression, the dexamethasone arm was halted and all patients randomized to dexamethasone were offered VELCADE, regardless of disease status. At this time of study termination, a final statistical analysis was performed. Due to this early termination of the study, the median duration of follow-up for surviving patients (n=534) is limited to 8.3 months. The time to event analyses and response rates from the APEX trial are presented in Table 8.

Table 8: Summary of Efficacy Analyses in the APEX Study

Efficacy Endpoint	All Patients		1 Prior Line of Therapy		>1 Prior Line of Therapy	
	VELCADE n=333	Dex n=336	VELCADE n=132	Dex n=119	VELCADE n=200	Dex n=217
Time to Progression – Events n (%)	147(44)	196(58)	55(42)	64(54)	92(46)	132(61)
Median ^a (95% CI)	6.2 mo (4.9, 6.9)	3.5 mo (2.9, 4.2)	7.0 (6.2, 8.8)	5.6 (3.4, 6.3)	4.9 (4.2, 6.3)	2.9 (2.8, 3.5)
Hazard ratio ^b (95% CI)	0.55 (0.44, 0.69)		0.55 (0.38, 0.81)		0.54 (0.41, 0.72)	
p-value ^c	<0.0001		0.0019		<0.0001	
Overall survival Events (deaths) n (%)	51(15)	84(25)	12(9)	24(20)	39(20)	60(28)
Hazard ratio ^b (95% CI)	0.57 (0.40, 0.81)		0.39 (0.19, 0.81)		0.65 (0.43, 0.97)	
p-value ^{c, d}	<0.05		<0.05		<0.05	
Response Rate population^e n=627	n=315	n=312	n=128	n=110	n=187	n=202
CR ^f n(%)	20(6)	2(<1)	8(6)	2(2)	12(6)	0(0)
PR ^f n(%)	101(32)	54(17)	49(38)	27(25)	52(28)	27(13)
nCR ^{f, g} n(%)	21(7)	3(<1)	8(6)	2(2)	13(7)	1(<1)
CR + PR ^f n(%)	121(38)	56(18)	57(45)	29(26)	64(34)	27(13)
p-value ^h	<0.0001		0.0035		<0.0001	
Median Response Duration						
CR ^f	9.9 mo	NE ⁱ	9.9 mo	NE	6.3 mo	NA ^j
nCR ^f	11.5 mo	9.2 mo	NE	NE	11.5 mo	9.2 mo
CR + PR ^f	8.0 mo	5.6 mo	8.1 mo	6.2 mo	7.8 mo	4.1 mo

^a Kaplan-Meier estimate

^b Hazard ratio is based on Cox proportional-hazard model with the treatment as single independent variable. A hazard ratio less than 1 indicates an advantage for VELCADE.

^c p-value based on the stratified log-rank test including randomisation stratification factors.

^d Precise p-value cannot be rendered

^e Response population includes patients who had measurable disease at baseline and received at least 1 dose of study dose

^f EBMT criteria; nCR meets all EBMT criteria for CR but has positive IF. Under EBMT criteria, nCR in the PR category.

^g In 2 patients, the IF was unknown.

^h p-value for Response Rate (CR + PR) from the Chochran-Mantel-Haenszel chi-square test adjusted for the stratification factors;

ⁱ Not Estimable.

^j Not Applicable, no patients in category.


For the 121 patients achieving a response (CR or PR) on the VELCADE arm, the median duration was 8.0 months (95% CI: 6.9, 11.5 months) compared to 5.6 months (95% CI: 4.8, 9.2 months) for the 56 responders on the dexamethasone arm.

Treatment with VELCADE led to a significantly longer TTP, a significantly prolonged survival and a significantly higher response rate, compared to treatment with dexamethasone in patients who have received more than one prior therapy as well as in patients who have received only one prior line of therapy.

Both in patients who were refractory to their last prior therapy and those who were not refractory, overall survival was significantly longer and response rate was significantly higher on the VELCADE arm. Of the 669 patients enrolled, 245 (37%) were 65 years of age or older. Response parameters as well as TTP remained significantly better for VELCADE independently of age. Regardless of β 2- microglobulin levels at baseline, all efficacy parameters (time to progression and overall survival, as well as response rate) were significantly improved on the VELCADE arm.


The time to progression (TTP) was significantly longer on the VELCADE arm (see Figure 5).

**Figure 5: Time to progression
Bortezomib vs Dexamethasone**



As shown in Figure 6, VELCADE had a significant survival advantage relative to dexamethasone (p<0.05). The median follow-up was 8.3 months.

**Figure 6: Overall Survival
Bortezomib vs Dexamethasone**



Phase II studies

The safety and efficacy of VELCADE were evaluated in an open-label, single-arm, multi-centre study of 202 patients who had received at least 2 prior therapies and demonstrated disease progression on their most recent therapy. The median number of prior therapies was six. Dosing regimens and baseline patient and disease characteristics are summarised in Table 6 and Table 7. The study employed dose modifications for toxicity (see **DOSAGE AND ADMINISTRATION**). Responses to VELCADE alone in the phase II study are shown in Table 9.

In general, patients who had confirmed Complete Response received 2 additional cycles of VELCADE treatment beyond confirmation. The median time to response was 38 days (range 30 to 127 days). The median survival of all patients enrolled was 16 months (range <1 to 18+ months). The response rate to VELCADE was independent of the number and types of prior therapies.

Table 9: Summary of disease outcomes in Phase II study

Response Analyses (VELCADE monotherapy) N=188	N (%)	(95% CI)
Overall Response Rate (CR + PR)	52 (27.7%)	(21, 35)
Complete Response (CR) ¹	5 (2.7%)	(1, 6)
Partial Response (PR) ²	47 (25%)	(19, 32)
Clinical Remission (SWOG) ³	33 (17.6%)	(12, 24)
Kaplan-Meier Estimated Median Duration of Response (95% CI)	365 Days	(224, NE)

¹**Complete Response** required 100% disappearance of the original monoclonal protein from blood and urine on at least 2 determinations at least 6 weeks apart by immunofixation, and <5% plasma cells in the bone marrow on at least two determinations for a minimum of six weeks, stable bone disease and calcium.

²**Partial Response** required $\geq 50\%$ reduction in serum myeloma protein and $\geq 90\%$ reduction of urine myeloma protein on at least 2 occasions for a minimum of at least 6 weeks, stable bone disease and calcium.

³**Clinical remission (SWOG)** required $\geq 75\%$ reduction in serum myeloma protein and/or $\geq 90\%$ reduction of urine myeloma protein on at least 2 occasions for a minimum of at least 6 weeks, stable bone disease and calcium.

Patients who did not obtain an optimal response to therapy with VELCADE alone were able to receive high-dose dexamethasone in conjunction with VELCADE (i.e., 40 mg dexamethasone with each dose of VELCADE administered orally as 20 mg on the day of and 20 mg the day after VELCADE administration, (i.e., Days 1, 2, 4, 5, 8, 9, 11, and 12), thus 160mg over 3 weeks. Eighteen percent (13/74) of patients achieved or had an improved response (CR 11% or PR 7%) with combination treatment.

A small dose-response study was performed in 54 patients with multiple myeloma who received a 1.0 mg/m²/dose or a 1.3 mg/m²/dose twice weekly for two out of three weeks. A single complete response was seen at each dose, and there were overall (CR + PR) response rates of 30% (8/27) at 1.0 mg/m² and 38% (10/26) at 1.3 mg/m².

Patients with previously treated light-chain (AL) Amyloidosis

A Phase 1/2 study was conducted to determine the safety and efficacy of VELCADE in patients with previously treated light-chain (AL) Amyloidosis. No new safety concerns were observed during the study, and in particular VELCADE did not exacerbate target organ damage (heart, kidney and liver). In 49 evaluable patients treated at 1.6 mg/m² weekly or 1.3 mg/m² twice-weekly, a 67.3% response rate (including a 28.6% CR rate) as measured by haematological response (M-protein) was reported. For these dose cohorts, the combined 1-year survival rate was 88.1%.

Paediatric Use

The safety and effectiveness of VELCADE in children has not been established.

INDICATIONS

VELCADE in combination with melphalan and prednisone, is indicated for the treatment of patients with previously untreated multiple myeloma, who are not suitable for high dose chemotherapy.

VELCADE, as part of combination therapy, is indicated for induction therapy prior to high dose chemotherapy with autologous stem cell rescue for patients under 65 years of age with previously untreated multiple myeloma

VELCADE is also indicated for the treatment of multiple myeloma patients who have received at least one prior therapy, and who have progressive disease.

CONTRAINDICATIONS

VELCADE is contraindicated in patients with hypersensitivity to bortezomib, boron or mannitol.

PRECAUTIONS

VELCADE should be administered under the supervision of a physician experienced in the use of antineoplastic therapy.

There have been fatal cases of inadvertent intrathecal administration of VELCADE.

VELCADE is for IV use only. **DO NOT ADMINISTER VELCADE INTRATHECALLY.**

Overall, the safety profile of patients treated with VELCADE in monotherapy was similar to that observed in patients treated with VELCADE in combination with melphalan and prednisone.

Peripheral Neuropathy

VELCADE treatment causes a peripheral neuropathy that is predominantly sensory. However, cases of severe motor neuropathy with or without sensory peripheral neuropathy have been reported. Patients with pre-existing symptoms (numbness, pain or burning feeling

in the feet or hands) and/or signs of peripheral neuropathy may experience worsening (including \geq Grade 3) during treatment with VELCADE. Patients should be monitored for symptoms of neuropathy, such as a burning sensation, hyperaesthesia, hypoesthesia, paraesthesia, discomfort, neuropathic pain or weakness. Patients experiencing new or worsening peripheral neuropathy may require change in the dose and schedule of VELCADE (see **DOSAGE AND ADMINISTRATION**).

Following dose adjustments, improvement in or resolution of peripheral neuropathy was reported in 51% of patients with \geq Grade 2 peripheral neuropathy in the APEX study. Improvement in or resolution of peripheral neuropathy was reported in 73% of patients who discontinued due to Grade 2 neuropathy or who had \geq Grade 3 peripheral neuropathy in the phase II studies (see **ADVERSE REACTIONS**).

In addition to peripheral neuropathy, there may be a contribution of autonomic neuropathy to some adverse reactions such as postural hypotension and severe constipation with ileus. Information on autonomic neuropathy and its contribution to these undesirable effects is limited.

Hypotension

Patients developing orthostatic hypotension on VELCADE did not have evidence of orthostatic hypotension prior to treatment with VELCADE. Most patients required treatment for their orthostatic hypotension. A minority of patients with orthostatic hypotension experienced syncopal events. Orthostatic/postural hypotension was not acutely related to bolus infusion of VELCADE.

In phase II studies and the APEX study, the incidence of hypotension (postural, orthostatic and hypotension not otherwise specified) was 11% to 12%. These events are observed throughout therapy. Caution should be used when treating patients with a history of syncope receiving medications known to be associated with hypotension and with patients who are dehydrated. Management of orthostatic/postural hypotension may include adjustment of antihypertensive medications, hydration, or administration of mineralocorticoids and/or sympathomimetics (see **ADVERSE REACTIONS**).

Cardiac Disorders

Acute development or exacerbation of congestive heart failure, and/or new onset of decreased left ventricular ejection fraction has been reported, including reports in patients with few or no risk factors for decreased left ventricular ejection fraction. Patients with risk factors for, or an existing heart disease should be closely monitored. In the APEX study, the incidence of any treatment-emergent cardiac disorder was 15% and 13% in the VELCADE and dexamethasone groups, respectively. The incidence of heart failure events (acute pulmonary edema, cardiac failure, congestive cardiac failure, cardiogenic shock, pulmonary edema) was similar in the VELCADE and dexamethasone groups, 5% and 4%, respectively. There have been isolated cases of QT-interval prolongation in clinical studies; causality has not been established.

Pulmonary Disorders

There have been rare reports of acute diffuse infiltrative pulmonary disease of unknown etiology such as pneumonitis, interstitial pneumonia, lung infiltration and Acute Respiratory Distress Syndrome (ARDS) in patients receiving VELCADE. Some of these events have been fatal. A higher proportion of these events have been reported in Japan. In the event of new or worsening pulmonary symptoms, a prompt diagnostic evaluation should be performed and patients treated appropriately.

In a clinical trial, two patients given high-dose cytarabine ($2\text{g}/\text{m}^2$ per day) by continuous infusion with daunorubicin and VELCADE for relapsed acute myelogenous leukaemia died of ARDS early in the course of therapy.

Reversible Posterior Leukoencephalopathy Syndrome (RPLS)

There have been reports of RPLS in patients receiving VELCADE. RPLS is a rare, reversible, neurological disorder which can present with seizure, hypertension, headache, lethargy, confusion, blindness, and other visual and neurological disturbances. Brain imaging, preferably

MRI (Magnetic Resonance Imaging), is used to confirm the diagnosis. In patients developing RPLS, discontinue VELCADE. The safety of reinitiating VELCADE therapy in patients previously experiencing RPLS is not known.

Seizures

Seizures have been uncommonly reported in patients without previous history of seizures or epilepsy. Special care is required when treating patients with any risk factors for seizures.

Laboratory Tests

Complete blood counts (CBC) should be frequently monitored throughout treatment with VELCADE.

Thrombocytopenia: VELCADE treatment is associated with thrombocytopenia (see **ADVERSE REACTIONS**). Platelet counts were lowest at Day 11 of each cycle of VELCADE treatment and typically recovered to baseline by the next cycle. On average, the pattern of platelet count decrease and recovery remained consistent over the 8 cycles of twice weekly dosing, and there was no evidence of cumulative thrombocytopenia. The mean platelet count nadir measured was approximately 40% of baseline. The severity of thrombocytopenia related to pre-treatment platelet count is shown in Table 10 for the APEX study. In the APEX study, the incidence of significant bleeding events (\geq Grade 3) was similar on both the VELCADE (4%) and dexamethasone (5%) arms. Platelet counts should be monitored prior to each dose of VELCADE. VELCADE therapy should be held when the platelet count is $<25,000/\mu\text{L}$ and reinitiated at a reduced dose after resolution (see **DOSAGE AND ADMINISTRATION** and **ADVERSE REACTIONS**). Transfusions may be used at the discretion of the physician. There have been reports of gastrointestinal and intracerebral haemorrhage in association with VELCADE.

Table 10: The Severity of Thrombocytopenia Related to Pre-treatment Platelet Count in the APEX study

Pre-treatment Platelet Count*	Number of Patients (N= 331)**	Number (%) of Patients with Platelet Count $< 10,000/\mu\text{L}$	Number (%) of Patients with Platelet Count $10,000/\mu\text{L} - 25,000\mu\text{L}$
$\geq 75,000/\mu\text{L}$	309	8 (3%)	36 (12%)
$\geq 50,000/\mu\text{L} - <75,000/\mu\text{L}$	14	2 (14%)	11 (79%)
$\geq 10,000/\mu\text{L} - <50,000/\mu\text{L}$	7	1(14%)	5 (71%)

*A baseline platelet count of $50,000/\mu\text{L}$ was required for study eligibility.

**Data for one patient was missing at baseline

Thrombocytopenia was reported in 43% of patients in the phase II studies.

Gastrointestinal Adverse Events

VELCADE treatment can cause nausea, diarrhoea, constipation and vomiting (see **ADVERSE REACTIONS**) sometimes requiring use of antiemetics and antidiarrhoeals. Fluid and electrolyte replacement should be administered to prevent dehydration. Since patients receiving VELCADE therapy may experience vomiting and/or diarrhoea, patients should be advised regarding appropriate measures to avoid dehydration. Patients should be instructed to seek medical advice if they experience symptoms of dizziness, light headedness or fainting spells.

Tumour Lysis Syndrome

Because VELCADE is a cytotoxic agent and can rapidly kill malignant cells the complications of tumour lysis syndrome may occur. The patients at risk of tumour lysis syndrome are those with high tumour burden prior to treatment. These patients should be monitored closely and appropriate precautions taken.

Hepatic Events

Rare cases of acute liver failure have been reported in patients receiving multiple concomitant medications and with serious underlying medical conditions. Other reported hepatic events include increases in liver enzymes, hyperbilirubinemia, and hepatitis. Such changes may be reversible upon discontinuation of VELCADE. There is limited re-challenge information in these patients.

Carcinogenicity, Mutagenicity and Impairment of Fertility

Carcinogenicity studies have not been conducted with bortezomib.

Bortezomib showed clastogenic activity at a high concentration (3 µg/mL) in an *in vitro* chromosomal aberration assay using Chinese hamster ovary cells. Clastogenic activity was not observed *in vivo* in a mouse micronucleus test using intravenous doses of up to 3 mg/m². Bortezomib was not genotoxic in *in vitro* tests for bacterial gene mutation.

Fertility studies with bortezomib were not performed but degenerative changes seen in the testes and ovary in a rat general toxicity study suggest that VELCADE may affect male and female fertility.

Use in Pregnancy

Category C

Women of child bearing potential should avoid becoming pregnant while being treated with VELCADE. The placental transfer of bortezomib is unknown, but any occurrence may disrupt cycling in the developing fetus, although teratogenicity was not observed in rats and rabbits at maximum tolerated doses.

Bortezomib was not teratogenic in nonclinical developmental toxicity studies in rats and rabbits at the highest dose tested (approximately 0.5 mg/m²/day) when administered during organogenesis. These dosages are approximately half the clinical dose of 1.3 mg/m² based on body surface area and calculated on a single-dose basis. Increased post-implantation loss and reduced foetal weights were seen in rabbits at the highest dose tested, which was a maternally toxic dose. Litter values were unaffected by a non-maternotoxic dose (approximately 0.3 mg/m²/day).

No placental transfer studies have been conducted with bortezomib. There are no adequate and well-controlled studies in pregnant women. If VELCADE is used during pregnancy, or if the patient becomes pregnant while receiving this drug, the patient should be informed of the potential hazard to the foetus.

Patients should be advised to use effective contraceptive measures to prevent pregnancy.

Use in Lactation

It is not known whether bortezomib or its metabolites are excreted in animal or human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in breast-fed infants from VELCADE, women should be advised against breast-feeding while being treated with VELCADE.

Interactions with Other Drugs

In vitro and animal *ex vivo* studies indicate that bortezomib is a weak inhibitor of cytochrome P450 (CYP) isoenzymes, 1A2, 2C9, 2C19, 2D6, and 3A4. Based on the limited contribution (7%) of CYP2D6 to the metabolism of bortezomib, the CYP2D6 poor metabolizer phenotype is not expected to affect the overall disposition of bortezomib.

A drug-drug interaction study assessing the effect of ketoconazole, a potent CYP3A inhibitor, on the pharmacokinetics of VELCADE, showed a bortezomib AUC mean increase of 35%, based on data from 12 patients. Therefore, patients should be closely monitored when given bortezomib in combination with potent CYP3A4-inhibitors (e.g., ketoconazole, ritonavir).

In a drug-drug interaction study assessing the effect of omeprazole, a potent inhibitor of CYP2C19, on the pharmacokinetics of VELCADE, there was no significant effect on the pharmacokinetics of bortezomib, based on data from 17 patients.

A drug-drug interaction study assessing the effect of rifampicin, a potent CYP3A4 inducer, on the pharmacokinetics of VELCADE showed a mean bortezomib AUC reduction of 45% based

on data from 6 patients. The concomitant use of VELCADE with strong CYP3A4 inducers is not recommended, as efficacy may be reduced. Examples of CYP-3A4 inducers are rifampicin, carbamazepine, phenytoin, phenobarbital and St. John's Wort. IN the same drug-drug interaction study, the effect of dexamethasone, a weaker CYP3A4 inducer was assessed. There was no significant effect on bortezomib pharmacokinetics based on data from 7 patients.

A drug-drug interaction study assessing the effect of melphalan-prednisone on VELCADE showed a 17% increase in mean bortezomib AUC based on data from 21 patients. This is not considered clinically relevant.

During clinical trials, hypoglycaemia and hyperglycaemia were reported in diabetic patients receiving oral hypoglycaemics. Patients on oral antidiabetic agents receiving VELCADE treatment may require close monitoring of their blood glucose levels and adjustment of the dose of their antidiabetic medication.

Patients should be cautioned about the use of concomitant medications that may be associated with peripheral neuropathy (such as amiodarone, anti-virals, isoniazid, nitrofurantoin, or statins), or with a decrease in blood pressure.

Effects on Laboratory Tests

None known.

Effect on Ability to Drive or Operate Machinery

VELCADE may cause tiredness, dizziness, fainting or blurred vision. Patients should be advised not to drive or operate machinery if they experience these symptoms.

ADVERSE REACTIONS

Adverse events

Summary of Clinical Trials in patients with previously untreated multiple myeloma:

Results from the GIMEMA and IFM2005 studies

The following table describes the safety data from the GIMEMA and IFM2005 studies in patients with previously untreated multiple myeloma who were eligible for autologous stem cell transplantation, and received VELCADE (1.3 mg/m²) in combination with thalidomide (100 mg, then 200 mg) and dexamethasone (40 mg) in the GIMEMA study, or dexamethasone (40 mg) in the IFM2005 study.

Table 11: Adverse events following induction in randomised, controlled studies GIMEMA and IFM2005

Adverse event, n (%)	GIMEMA		IFM2005	
	VcTD n=236	TD n=238	VcD n=239	VAD n=239
Any adverse event	nr	nr	231 (96.7)*	219 (91.6)*
Any serious adverse event	31 (13.1)	30 (12.6)	65 (27.2)	81 (33.9)
Any grade 3 or 4 adverse event	132 (55.9)	79 (33.1)	112 (46.9)	110 (46.0)
Any grade 3 or 4 non-haematologic adverse event	120 (50.8)	73 (30.6)	nr	nr
Skin rash	24 (10.1)	4 (1.6)	0 (0)	0 (0)
Peripheral neuropathy	23 (9.7)	5 (2.1)	17 (7.1)	5 (2.1)
Deep vein thrombosis	8 (3.3)	12 (5.0)	nr	nr
Constipation	10 (4.2)	7 (2.9)	nr	nr
Infections	nr	nr	21 (8.8)	29 (12.1)
Infections excluding herpes zoster	7 (2.9)	11 (4.6)	nr	nr
Herpes zoster (all grades)	nr	nr	22 (9.2)	5 (2.1)
Gastrointestinal events (excluding constipation where individually reported)	5 (2.1)	1 (0.4)	0 (0)	0 (0)
Cardiac toxicity	5 (2.1)	5 (2.1)	0 (0)	0 (0)
Liver toxicity	4 (1.6)	7 (2.9)	nr	nr

Fatigue	nr	nr	0 (0)	0 (0)
Pneumonia	nr	nr	nr	nr
Any grade 3 or 4 haematologic adverse event	nr	nr	nr	nr
Anaemia	nr	nr	10 (4.2)*	21 (8.8)*
Neutropaenia	nr	nr	12 (5.0)*	24 (10.0)*
Thrombocytopenia	nr	nr	7 (2.9)	3 (1.3)
Thrombosis	nr	nr	4 (1.7)*	13 (5.4)*
Discontinued during or after induction therapy	13 (5.5)	26 (10.9)	44 (18.4)	32 (13.4)
Adverse event leading to death	1 (0.4)	0 (0)	0 (0)*	7 (2.9)*

* $p < 0.05$ for comparison of AE rate between VcD and VADVcTD: VELCADE-thalidomide-dexamethasone; TD: thalidomide-dexamethasone; VcD: VELCADE-dexamethasone; VAD: vincristine-doxorubicine-dexamethasone.

During consolidation therapy of the GIMEMA study, grade 3-4 adverse events were similar to those reported during induction, although rates were much lower. Notably, the rate of grade 3-4 peripheral neuropathy was 1.2% with VcTD consolidation.

Results from the VISTA study

The following table describes safety data from the VISTA study in 340 patients with previously untreated multiple myeloma who received VELCADE (1.3 mg/m²) in combination with melphalan (9 mg/m²) and prednisone (60 mg/m²) in the VISTA study.

Table 12: Treatment Emergent Drug-Related Adverse Events reported in $\geq 10\%$ of patients treated with VELCADE in combination with melphalan and prednisone

MedDRA System Organ Class Preferred Term	----- VcMP ----- (n=340)			----- MP ----- (n=337)		
	Total n (%)	Toxicity Grade, n (%) 3 ≥ 4		Total n (%)	Toxicity Grade, n (%) 3 ≥ 4	
Blood and Lymphatic System Disorders						
Thrombocytopenia	164 (48)	60 (18)	57 (17)	140 (42)	48 (14)	39 (12)
Neutropenia	160 (47)	101 (30)	33 (10)	143 (42)	77 (23)	42 (12)
Anaemia	109 (32)	41 (12)	4 (1)	156 (46)	61 (18)	18 (5)
Leukopenia	108 (32)	64 (19)	8 (2)	93 (28)	53 (16)	11 (3)
Lymphopenia	78 (23)	46 (14)	17 (5)	51 (15)	26 (8)	7 (2)
Gastrointestinal Disorders						
Nausea	134 (39)	10 (3)	0	70 (21)	1 (<1)	0
Diarrhoea	119 (35)	19 (6)	2 (1)	20 (6)	1 (<1)	0
Vomiting	87 (26)	13 (4)	0	41 (12)	2 (1)	0
Constipation	77 (23)	2 (1)	0	14 (4)	0	0
Abdominal Pain Upper	34 (10)	1 (<1)	0	20 (6)	0	0
Nervous System Disorders						
Peripheral Neuropathy	156 (46)	42 (12)	2 (1)	4 (1)	0	0
Neuralgia	117 (34)	27 (8)	2 (1)	1 (<1)	0	0
Paraesthesia	42 (12)	6 (2)	0	4 (1)	0	0
General Disorders and Administration Site Conditions						
Fatigue	85 (25)	19 (6)	2 (1)	48 (14)	4 (1)	0
Asthenia	54 (16)	18 (5)	0	23 (7)	3 (1)	0
Pyrexia	53 (16)	4 (1)	0	19 (6)	1 (<1)	1 (<1)
Infections and Infestations						
Herpes Zoster	39 (11)	11 (3)	0	9 (3)	4 (1)	0
Metabolism and Nutrition Disorders						
Anorexia	64 (19)	6 (2)	0	19 (6)	0	0
Skin and Subcutaneous Tissue Disorders						
Rash	38 (11)	2 (1)	0	7 (2)	0	0
Psychiatric Disorders						
Insomnia	35 (10)	1 (<1)	0	21 (6)	0	0

Herpes zoster virus reactivation

Physicians should consider using antiviral prophylaxis in patients being treated with VELCADE. In the VISTA study in patients with previously untreated multiple myeloma, the overall incidence of herpes zoster reactivation was more common in patients treated with VcMP compared with MP (14% vs 4% respectively). Antiviral prophylaxis was administered to 26% of the patients in the VcMP arm. The incidence of herpes zoster among patients in the VcMP treatment group was 17% for patients not administered antiviral prophylaxis compared to 3% for patients administered antiviral prophylaxis. Similar results were observed during the IFM2005 study; herpes zoster was more common in patients treated with VELCADE-based regimen compared to control regimen. During consolidation, the GIMEMA study reported similar rates (0.6%) of grade 3-4 incidences of herpes zoster between the two study arms ($p=1.0000$).

Summary of Clinical Trials in patients with relapsed/refractory multiple myeloma:

The adverse events most commonly reported, regardless of causality, in the APEX study in relapsed / refractory multiple myeloma patients (see **CLINICAL TRIALS**) are presented in Tables 13. All adverse events occurring at $\geq 10\%$ are included.

Table 13: Most Commonly Reported ($\geq 10\%$ in VELCADE arm) Adverse Events in the APEX study using the 1.3 mg/m² dose (N=663)						
	VELCADE (N=331)			Dexamethasone (N=332)		
	All Events %	Grade 3 %	Grade 4 %	All Events %	Grade 3 %	Grade 4 %
Adverse Event	100	61	14	98	44	16
Body as a Whole-General Disorders						
Asthenic conditions (fatigue, malaise, weakness)	61	12	<1	45	6	0
Pyrexia	35	2	0	16	1	<1
Rigors	11	0	0	2	0	0
Oedema lower limb	11	0	0	13	<1	0
Gastro-Intestinal System Disorders						
Diarrhea	57	7	0	21	2	0
Nausea	57	2	0	14	0	0
Constipation	42	2	0	15	1	0
Vomiting	35	3	0	6	1	0
Abdominal pain	16	2	0	4	<1	0
Central & Peripheral Nervous System Disorders						
Peripheral Neuropathy*	36	7	<1	9	<1	<1
Paresthesia and dysesthesia	27	2	0	11	<1	0
Headache	26	<1	0	13	<1	0
Dizziness (excluding vertigo)	14	<1	0	10	0	0
Blood and lymphatic system disorders						
Thrombocytopenia	35	26	4	11	5	1
Anemia	26	9	<1	22	10	<1
Neutropenia	19	12	2	2	1	0
Psychiatric disorders						
General	35	3	<1	49	5	1
Insomnia	18	<1	0	27	2	0
Metabolic and Nutritional Disorders						
Appetite decreased and anorexia	34	3	0	9	<1	0
Respiratory System disorders						
Cough	21	<1	0	11	<1	0
Dyspnoea	20	5	<1	17	3	<1
Skin and subcutaneous tissue disorders						
Rash	18	1	0	6	0	0
Infections and infestations						
Lower respiratory/lung infections	15	4	<1	21	5	<1
Nasopharyngitis	14	<1	0	7	0	0
Herpes zoster	13	2	0	5	1	<1
Musculoskeletal and connective tissue disorders						
Bone pain	16	4	0	15	3	0
Pain in limb	15	2	0	7	<1	0
Back pain	14	3	0	10	1	0
Arthralgia	14	<1	0	11	2	0
Muscle cramps	12	0	0	15	<1	0
Myalgia	12	<1	0	5	<1	0

*Peripheral neuropathy includes all terms under peripheral neuropathy not elsewhere classified (NEC), (Peripheral neuropathy not otherwise specified (NOS), peripheral neuropathy aggravated, peripheral sensory neuropathy and peripheral motor neuropathy and neuropathy NOS).

Serious Adverse Events (SAEs)

In the APEX study, 44% of patients from the VELCADE treatment arm experienced a SAE during the study, as did 43% of dexamethasone-treated patients. The most commonly reported SAEs in the VELCADE treatment arm were pyrexia (6%), diarrhoea (5%), dyspnoea and pneumonia (4%) and vomiting (3%). In the dexamethasone group, the most common SAEs were pneumonia (7%), pyrexia (4%) and hyperglycaemia (3%). Twenty five percent (25%) and 18% of VELCADE and dexamethasone patients respectively were discontinued from treatment due to adverse events assessed as drug related by the investigators. The most common for VELCADE discontinuation was peripheral neuropathy (8%) and for dexamethasone was psychotic disorder and hyperglycaemia (2% each).

In the APEX study, 4 deaths were considered to be VELCADE-related: 1 case each of cardiogenic shock, respiratory insufficiency, congestive heart failure and cardiac arrest. Four (4) deaths were considered dexamethasone-related: 2 cases of sepsis, 1 case of bacterial meningitis and 1 case of sudden death at home. In the phase II studies 2 deaths were reported and considered by the investigator to be possibly related to VELCADE: 1 case of cardiopulmonary arrest and 1 case of respiratory failure.

Adverse reactions

The following adverse reactions were considered to have at least a possible or probable causal relationship to VELCADE by the investigators during 5 non-comparative phase II studies and 1 comparative phase III trial (APEX) in 663 patients with relapsed or refractory multiple myeloma, of whom 331 received VELCADE as single agent. The safety database comprises data from patients with multiple myeloma or B-cell lymphocytic leukaemia. Patients were treated with VELCADE as a single agent, or in combination with dexamethasone.

Adverse drug reactions are listed below by system organ class and frequency. Frequencies are defined as: Very common (>1/10); common (>1/100, <1/10); uncommon (>1/1,000, <1/100); rare (>1/10,000, <1/1,000); very rare (<1/10,000), including isolated reports.

Infections and infestations

Common: herpes zoster, pneumonia, bronchitis, sinusitis, nasopharyngitis, herpes simplex.

Uncommon: candidal infection, gastroenteritis, upper and lower respiratory tract infection, infection, influenza, fungal infection, sepsis, urinary tract infection, catheter related infection, haemophilus infection, pneumonia pneumococcal, post herpetic neuralgia, bacteraemia, blepharitis, bronchopneumonia, cytomegalovirus infection, infectious mononucleosis, varicella, oral candidiasis, pleural infection.

Neoplasms benign, malignant and unspecified (including cysts and polyps)

Uncommon: tumour lysis syndrome (see **PRECAUTIONS**)

Blood and lymphatic system disorders

Very Common: thrombocytopenia (see **PRECAUTIONS**), anaemia, neutropenia.

Common: leukopenia, lymphopenia.

Uncommon: lymphadenopathy, febrile neutropenia, pancytopenia, haemolytic anaemia, thrombocytopenic purpura.

Immune system disorders

Uncommon: hypersensitivity, immunocomplex mediated hypersensitivity.

Metabolism and nutrition disorders

Very Common: appetite decreased.

Common: dehydration, hyperglycaemia, hypokalaemia.

Uncommon: hypercalcaemia, hyperkalaemia, hyperuricaemia, hyponatraemia, hypernatraemia, hypocalcaemia, hypomagnesaemia, hypophosphataemia, hypoglycaemia, appetite increased, cachexia, vitamin B12 deficiency.

Endocrine disorders

Uncommon: Inappropriate antidiuretic hormone (ADH) secretion.

Psychiatric disorders

Common: insomnia, anxiety, confusion, depression.

Uncommon: agitation, delirium, restlessness, mood swings, mental status changes, sleep disorder, irritability, hallucinations, abnormal dreams.

Nervous system disorders

Very Common: peripheral neuropathy, peripheral sensory neuropathy (see **PRECAUTIONS**), headache, paraesthesia.

Common: dizziness (excluding vertigo), dysgeusia, peripheral neuropathy aggravated, polyneuropathy, dysaesthesia, hypoaesthesia, tremor.

Uncommon: convulsions, syncope, disturbance in attention, increased activity, ageusia, somnolence, migraine, peripheral motor neuropathy, jerky movements, dizziness postural, sciatica, cognitive disorder, mononeuropathy, paresis, restless leg syndrome, speech disorder, intracranial haemorrhage, paraplegia, subarachnoid haemorrhage.

Eye disorders

Common: vision blurred (see **PRECAUTIONS**), eye pain.

Uncommon: dry eye, conjunctivitis, eye discharge, vision abnormal, eye haemorrhage, photophobia, eye irritation, lacrimation increased, conjunctival hyperaemia, eye swelling.

Ear and labyrinth disorders

Common: vertigo.

Uncommon: tinnitus, deafness, hypoacusis, hearing impaired.

Cardiac disorders

Uncommon: Development or exacerbation of congestive heart failure (see **PRECAUTIONS**), cardiac failure, ventricular hypokinesia, pulmonary oedema and acute pulmonary oedema, cardiac arrest, cardiogenic shock, tachycardia, sinus tachycardia, supraventricular tachycardia, arrhythmia, atrial fibrillation, palpitations, sinus arrest, atrioventricular block complete, angina pectoris, angina unstable, myocardial infarction.

Rare: New onset of decreased left ventricular ejection fraction.

Vascular disorders

Common: hypotension, orthostatic and postural hypotension (see **PRECAUTIONS**), phlebitis, haematoma, hypertension.

Uncommon: flushing, petechiae, hot flushes, ecchymosis, purpura, cerebral hemorrhage, vasculitis, vein discolouration, vein distended, wound hemorrhage, pulmonary hypertension, cerebrovascular accident.

Respiratory, thoracic and mediastinal disorders

Very Common: dyspnoea.

Common: epistaxis, dyspnoea exertional, cough, rhinorrhoea.

Uncommon: nasal congestion, wheezing, pleural effusion, hoarseness, chest wall pain, hypoxia, pulmonary congestion, rhinitis, asthma, hyperventilation, orthopnoea, sinus pain, throat tightness, productive cough, respiratory alkalosis, respiratory arrest, tachypnoea.

Gastrointestinal disorders (see PRECAUTIONS)

Very Common: nausea, diarrhoea, vomiting, constipation.

Common: abdominal pain, dyspepsia, loose stools, abdominal pain upper, flatulence, abdominal distension, hiccups, mouth ulceration, pharyngolaryngeal pain, stomatitis, dry mouth.

Uncommon: ileus paralytic, abdominal discomfort, eructation, gastrointestinal motility disorder, oral pain, retching, antibiotic associated colitis, change in bowel habit, diarrhoea haemorrhagic, gastrointestinal haemorrhage, spleen pain, colitis, dysphagia, oesophagitis, gastritis, gastro-oesophageal reflux disease, gastrointestinal pain, gingival bleeding, gingival pain, haematemesis, hiatus hernia, irritable bowel syndrome, oral mucosal petechiae, rectal haemorrhage, salivary hypersecretion, tongue coated, tongue discolouration, enteritis, faecal impaction, acute pancreatitis.

Hepatobiliary disorders (see PRECAUTIONS)

Uncommon: hyperbilirubinaemia, hepatitis, hepatic haemorrhage, hypoproteinaemia

Skin and subcutaneous tissue disorders

Very Common: rash.

Common: pruritus, erythema, periorbital oedema, urticaria, rash pruritic, sweating increased, dry skin, eczema.

Uncommon: night sweats, rash erythematous, alopecia, contusion, pruritus generalised, rash macular, rash papular, skin nodule, rash generalized, dermatitis, eyelid oedema, nail disorder, photosensitivity reaction, skin discolouration, dermatitis atopic, hair texture abnormal, heat rash, psoriasis, vasculitic rash, face oedema, pressure sore, ichthyosis.

Musculoskeletal and connective tissue disorders

Very Common: myalgia.

Common: pain in limb, muscle cramps, arthralgia, bone pain, peripheral swelling, muscle weakness, back pain, musculoskeletal pain.

Uncommon: joint stiffness, buttock pain, joint swelling, muscle spasms, muscle twitching or sensation of heaviness, muscle stiffness, swelling, pain in jaw.

Renal and urinary disorders

Common: renal impairment, dysuria.

Uncommon: renal failure acute, renal colic, haematuria, proteinuria, urinary frequency, difficulty in micturition, renal failure, oliguria, urinary retention, loin pain, urinary incontinence, micturition urgency.

General disorders and administration site conditions

Very Common: fatigue (see **PRECAUTIONS**), pyrexia.

Common: weakness, rigors, malaise, influenza like illness, oedema peripheral, pain, lethargy, oedema, chest pain, asthenia.

Uncommon: fall, mucosal inflammation, feeling cold, chest pressure sensation, injection site phlebitis, mucosal haemorrhage, tenderness, injection site erythema, neuralgia, chest discomfort, groin pain, chest tightness, extravasation inflammation.

Investigations

Common: weight decreased, blood lactate dehydrogenase increased.

Uncommon: alanine aminotransferase increased, aspartate aminotransferase increased, blood alkaline phosphatase increased, blood creatinine increased, blood urea increased, gamma-glutamyltransferase increased, blood amylase increased, blood bilirubin increased, blood phosphate decreased, liver function tests abnormal, red blood cell count decreased, weight increased, white blood cell count decreased, blood bicarbonate decreased, heart rate irregular, C-reactive protein increased.

Injury, poisoning and procedural complications

Uncommon: catheter related complications, post procedural pain, post procedural haemorrhage, burns.

Reproductive system and breast disorders

Uncommon: testicular pain, erectile dysfunction.

Potentially immunocomplex-mediated reactions (see **PRECAUTIONS**)

Uncommon: potentially immunocomplex-mediated reactions, such as serum-sickness – type reaction, polyarthritis with rash and proliferative glomerulonephritis.

Post Marketing Experience

Clinically significant adverse reactions are listed if they have been reported during post approval use of VELCADE and have not been reported in clinical trials:

Blood and lymphatic system disorders

Rare: disseminated intravascular coagulation.

Cardiac Disorders

Rare: atrioventricular block complete, cardiac tamponade.

Ear and labyrinth disorders

Rare: deafness bilateral.

Eye Disorders

Rare: ophthalmic herpes, optic neuropathy and blindness.

Gastrointestinal disorders

Rare: ischemic colitis, acute pancreatitis.

Hepatobiliary disorders

Rare: liver failure

Infections and infestations

Rare: herpes meningoencephalitis

Immune System Disorders

Rare: angioedema

Nervous system disorders

Rare: encephalopathy, autonomic neuropathy, septic shock.

Respiratory, thoracic and mediastinal disorders

Rare: acute diffuse infiltrative pulmonary disease (see **PRECAUTIONS**)
pulmonary hypertension

Skin and subcutaneous tissue disorders

Rare: acute febrile neutrophilic dermatosis (Sweet's syndrome).

Very Rare: Stevens-Johnson Syndrome and toxic epidermal necrolysis

DOSAGE AND ADMINISTRATION

Recommended Dosage

VELCADE IS FOR INTRAVENOUS USE ONLY. Intrathecal administration has resulted in death.

Previously Untreated Multiple Myeloma

Transplant Eligible

1. VELCADE plus thalidomide-dexamethasone

During the induction stage, VELCADE (bortezomib) is administered twice weekly in combination with thalidomide-dexamethasone for three 3-week treatment cycles. Following stem cell transplantation, patients receive two 5-week cycles of VELCADE-thalidomide-dexamethasone. The treatment regimen is shown in Table 14.

Table 14: Recommended dosage regimen for VELCADE when used in combination with thalidomide and dexamethasone

Induction Therapy: Twice weekly VELCADE (3 cycles)											
Week	1				2				3		
Vc (1.3 mg/m ²)	Day 1	--	--	Day 4	Day 8	--	--	--	Day 11	--	--
t (100 mg)-Cycle 1	Day 1-7				Day 8-14				--		
t (200 mg)-Cycle 2-3	Day 1-7				Day 8-14				Day 15-21		
d (40 mg)	Day 1	Day 2	--	Day 4	Day 5	Day 8	Day 9	--	Day 11	Day 12	--

Consolidation Therapy: Once Weekly VELCADE (2 cycles)										
Week	1		2		3		4		5	
Vc (1.3 mg/m ²)	Day 1	--	Day 8	--	Day 15	--	Day 22	--	--	--
t (100 mg)	Day 1-7		Day 8-14		Day 15-21		Day 22-28		Day 29-35	
d (40 mg)	Day 1	Day 2	Day 8	Day 9	Day 15	Day 16	Day 22	Day 23	--	--

Vc = VELCADE; t = thalidomide; d = dexamethasone

2. VELCADE plus dexamethasone

VELCADE (bortezomib) is administered as an IV injection in combination with oral dexamethasone for four 3-week treatment cycles as shown in Table 15.

Table 15: Recommended dosage regimen for VELCADE when used in combination with dexamethasone

Week	1		2				3	
Vc (1.3 mg/m ²)	Day 1	Day 4	Day 8	--	Day 11	--	--	--
d (40 mg)-All Cycles	Day 1-4--		--				--	
d (40 mg)-Cycle 1-2			Day 9-12					

Vc = VELCADE; d = dexamethasone

Non-Transplant Eligible

VELCADE (bortezomib) is administered as a 3-5 second bolus IV injection in combination with oral melphalan and oral prednisone for nine 6-week treatment cycles as shown in Table 16. In Cycles 1-4, VELCADE is administered twice weekly (days 1, 4, 8, 11, 22, 25, 29 and 32). In Cycles 5-9, VELCADE is administered once weekly (days 1, 8, 22 and 29).

Table 16: Recommended Dosage Regimen for VELCADE when used in combination with melphalan and prednisone for Patients with Previously Untreated Multiple Myeloma

Twice Weekly VELCADE (Cycles 1-4)												
Week	1				2		3	4		5		6
Vc (1.3 mg/m ²)	Day 1	--	--	Day 4	Day 8	Day 11	rest period	Day 22	Day 25	Day 29	Day 32	rest period
m(9 mg/m ²) p(60 mg/m ²)	Day 1	Day 2	Day 3	Day 4	--	--	rest period	--	--	--	--	rest period

Once Weekly VELCADE (Cycles 5-9)										
Week	1				2	3	4	5	6	
Vc (1.3 mg/m ²)	Day 1	--	--	--	Day 8	rest period	Day 22	Day 29	rest period	
m (9 mg/m ²) p (60 mg/m ²)	Day 1	Day 2	Day 3	Day 4	--	rest period	--	--	rest period	

Vc = VELCADE; m = melphalan, p=prednisone

Dose Management Guidelines

Dose modification and re-initiation of therapy when VELCADE is administered in combination with melphalan and prednisone

Prior to initiating a new cycle of therapy:

- Platelet count should be $\geq 70 \times 10^9/L$ and the ANC should be $\geq 1.0 \times 10^9/L$
- Non-hematological toxicities should have resolved to Grade 1 or baseline

Table 17: Dose Modifications During Subsequent Cycles

Toxicity	Dose modification or delay
Haematological toxicity during a cycle:	
<ul style="list-style-type: none"> • If prolonged Grade 4 neutropenia or thrombocytopenia, or thrombocytopenia with bleeding is observed in the previous cycle 	Consider reduction of the melphalan dose by 25% in the next cycle.
<ul style="list-style-type: none"> • If platelet count $\leq 30 \times 10^9/L$ or ANC $\leq 0.75 \times 10^9/L$ on a VELCADE dosing day (other than day 1) 	Velcade dose should be withheld
<ul style="list-style-type: none"> • If several VELCADE doses in a cycle are withheld (≥ 3 doses during twice weekly administration or ≥ 2 doses during weekly administration) 	VELCADE dose should be reduced by 1 dose level (from 1.3 mg/m ² to 1 mg/m ² , or from 1 mg/m ² to 0.7 mg/m ²)
GRADE ≥ 3 NON-HAEMATOLOGICAL TOXICITIES	VELCADE therapy should be withheld until symptoms of the toxicity have resolved to Grade 1 or baseline. Then, VELCADE may be reinitiated with one dose level reduction (from 1.3 mg/m ² to 1 mg/m ² , or from 1 mg/m ² to 0.7 mg/m ²). For VELCADE-related neuropathic pain and/or peripheral neuropathy, hold and/or modify VELCADE as outlined in Table 18.

For additional information concerning melphalan and prednisone, see manufacturer's prescribing information.

Table 18: Recommended Dose Modification for VELCADE-related Neuropathic Pain and/or Peripheral Sensory or Motor Neuropathy.

Severity of Peripheral Neuropathy Signs and Symptoms	Modification of Dose and Regimen
Grade 1 (paraesthesias, weakness and/or loss of reflexes) without pain or loss of function	No action
Grade 1 with pain or Grade 2 (interfering with function but not with activities of daily living)	Reduce VELCADE to 1.0 mg/m ²
Grade 2 with pain or Grade 3 (interfering with activities of daily living)	Withhold VELCADE therapy until toxicity resolves. When toxicity resolves reinstate with a reduced dose of VELCADE at 0.7 mg/m ² and change treatment schedule to once per week.
Grade 4 (sensory neuropathy which is disabling or motor neuropathy that is life threatening or leads to paralysis)	Discontinue VELCADE

NCI Common Toxicity Criteria

Relapsed / Refractory Multiple Myeloma

The recommended dose of VELCADE is 1.3 mg/m²/dose administered as a 3-5 second bolus intravenous injection twice weekly for two weeks (days 1, 4, 8, and 11) followed by a 10-day rest period (days 12-21). This 3-week period is considered a treatment cycle. At least 72 hours should elapse between consecutive doses of VELCADE.

It is recommended that patients with a confirmed complete response receive 2 additional cycles of VELCADE beyond a confirmation. It is also recommended that responding patients who do not achieve a complete remission receive a total of 8 cycles of VELCADE therapy.

For extended therapy of more than 8 cycles, VELCADE may be administered on the standard schedule or on a maintenance schedule of once weekly for 4 weeks (days 1, 8, 15, and 22) followed by a 13-day rest period (days 23 to 35) (see **CLINICAL TRIALS** for a summary of dose administration during clinical trials).

Dose Modification and Reinitiation of Therapy

VELCADE therapy should be withheld at the onset of any Grade 3 non-haematological or Grade 4 haematological toxicities excluding neuropathy as discussed below (see **PRECAUTIONS**). Once the symptoms of the toxicity have resolved, VELCADE therapy may be reinitiated at a 25% reduced dose (1.3 mg/m²/dose reduced to 1.0 mg/m²/dose; 1.0 mg/m²/dose reduced to 0.7 mg/m²/dose). Table 18 above contains the recommended dose modification for the management of patients who experience VELCADE-related neuropathic pain and/or peripheral sensory neuropathy (Table 12 above). Patients with pre-existing severe neuropathy should be treated with VELCADE only after careful risk/benefit assessment.

Patients with Renal Impairment

The pharmacokinetics of VELCADE are not influenced by the degree of renal impairment. Therefore, dosing adjustments of VELCADE are not necessary for patients with renal insufficiency. Since dialysis may reduce VELCADE concentrations, the drug should be administered after the dialysis procedure (see **PHARMACOKINETICS**).

Patients with Hepatic Impairment

Patients with mild hepatic impairment do not require a starting dose adjustment and should be treated per the recommended VELCADE dose. Patients with moderate or severe hepatic impairment should be started on VELCADE at a reduced dose of 0.7 mg/m² per injection during the first cycle, and a subsequent dose escalation to 1.0 mg/m² or further dose reduction to 0.5 mg/m² may be considered based on patient tolerance (see Table 19).

Table 19: Recommended Starting Dose Modification for VELCADE in Patients with Hepatic Impairments

	Bilirubin Level	SGOT (AST) Levels	Modification of Starting Dose
Mild	≤1.0x ULN	>ULN	None
	>1.0x – 1.5x ULN	Any	None
Moderate	>1.5x – 3x ULN	Any	Reduce VELCADE to 0.7 mg/m ² in the first cycle. Consider dose escalation to 1.0 mg/m ² or further reduction to 0.5 mg/m ² in subsequent cycles based on patient tolerability
Severe	>3x ULN	Any	

SGOT = serum glutamic oxaloacetic transaminase;
AST = aspartate aminotransferase; ULN = upper limit of normal range

Administration

VELCADE is administered as a 3-5 second bolus intravenous injection through a peripheral or central intravenous catheter followed by a flush with 0.9% sodium chloride solution for injection.

Instructions for Use and Handling and Disposal

Administration Precautions: VELCADE is an antineoplastic. Caution should be used during handling and preparation. Proper aseptic technique should be used. Use of gloves and other protective clothing to prevent skin contact is recommended. In clinical trials, local skin irritation was reported in 5% of patients, but extravasation of VELCADE was not associated with tissue damage.

Reconstitution/Preparation for Intravenous Injection: Prior to use, the contents of each vial must be reconstituted to yield a solution with a concentration of 1 mg/mL, the contents of each 5 mL vial must be reconstituted with 1 mL of normal (0.9%) saline, Sodium Chloride for injection. The contents of each 10 mL vial must be reconstituted with 3.5 mL of normal (0.9%) saline, Sodium Chloride for Injection. The reconstituted product should be a clear and colourless solution.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit. If any discoloration or particulate matter is observed, the reconstituted product should not be used.

Procedure for proper disposal: Any unused product or waste material should be disposed of in accordance with local requirements.

OVERDOSAGE

Cardiovascular safety pharmacology studies in monkeys and dogs showed that IV doses approximately two to three times the recommended clinical dose on a mg/m² basis are associated with increases in heart rate, decreases in contractility, hypotension and death. The decreased cardiac contractility and hypotension responded to acute intervention with positive inotropic or pressor agents. In dog studies, a slight increase in the corrected QT interval was observed at a lethal dose.

In patients, overdosage more than twice the recommended dose has been associated with the acute onset of symptomatic hypotension and thrombocytopenia with fatal outcomes.

There is no known specific antidote for VELCADE overdosage. In the event of overdosage, patient's vital signs should be monitored and appropriate supportive care given to maintain

blood pressure (such as fluids, pressors, and/or inotropic agents) and body temperature (see **DOSAGE AND ADMINISTRATION** and **PRECAUTIONS**).

PRESENTATION

VELCADE is supplied in a 5 mL or 10 mL, type I, glass vial with a grey bromobutyl stopper and aluminum seal. The cap colour of the 5 mL vial is green, and the cap colour for the 10 mL vial is royal blue. The vial is contained in a transparent blister pack consisting of a tray with a lid. The 5 mL vial contains 11 mg powder (1.0 mg bortezomib) for injection and the 10 mL vial contains 38.5 mg powder (3.5 mg bortezomib) for injection.

VELCADE is available in cartons containing 1 vial. Product is for single use in one patient only.

Storage

Unopened vials: Store below 25°C. Keep the container in the outer carton in order to protect from light.

Reconstituted solution: VELCADE contains no antimicrobial preservative. The chemical and physical in-use stability of the reconstituted solution has been demonstrated for 8 hours at 25°C when it is stored under normal lighting conditions in the original vial and/or syringe prior to administration. However, to reduce microbiological hazard, use as soon as possible after dilution and if storage is necessary hold at 2-8°C for up to 8 hours.

POISON SCHEDULE OF THE DRUG

Prescription Only Medicine

SHELF LIFE

Unopened vial: 3 years

SPONSOR

JANSSEN-CILAG Pty Ltd
PO Box 9222 Newmarket
Auckland New Zealand
Telephone: (09) 523 8700

Date of Preparation: 07 March 2012

© VELCADE is a registered trademark of Millennium Pharmaceuticals, Inc.