



New Zealand Data Sheet

APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Presentation

APO-RISPERIDONE is available as oral, film-coated tablets of the following strengths:

0.5 mg APO-RISPERIDONE: Brownish-red, capsule shaped, film coated tablets engraved "APO" on one side, "RI" score ".5" on the other side.

1 mg APO-RISPERIDONE: White, capsule shaped, film coated tablets engraved "APO" on one side, "RI" score "1" on the other side.

2 mg APO-RISPERIDONE: Light orange, capsule shaped, film coated tablets engraved "APO" on one side, "RI" score "2" on the other side.

3 mg APO-RISPERIDONE: Beige, capsule shaped, film coated tablets engraved "APO" on one side, "RI" score "3" on the other side.

4 mg APO-RISPERIDONE: Light green, capsule shaped, film coated tablets engraved "APO" on one side, "RI" score "4" on the other side.

APO-RISPERIDONE is also available as a 1 mg/ml oral solution.

This product is not able to deliver all approved dose regimens.

Uses

Actions

Risperidone is a compound, which belongs to a new class of antipsychotic agents, the benzisoxazole derivatives.

Risperidone is a selective monoaminergic antagonist having a high affinity for serotonergic 5-HT₂ and dopaminergic D₂ receptors. Risperidone binds also to alpha₁-adrenergic receptors and, with lower affinity, to H₁-histamine and alpha₂-adrenergic receptors. Risperidone has no affinity for cholinergic receptors. The antipsychotic activity of risperidone is considered to be attributable to both risperidone and its active metabolite 9-hydroxy risperidone. Risperidone, as a potent D₂ antagonist, improves the positive symptoms of schizophrenia but causes less depression of motor activity and induction of catalepsy than classical neuroleptics. Balanced central serotonin and dopamine antagonism may reduce extrapyramidal side effect liability and extend the therapeutic activity to the negative and affective symptoms of schizophrenia.

Pharmacokinetics

Risperidone is completely absorbed after oral administration, reaching peak plasma concentrations within 1 to 2 hours. The absorption is not affected by food and thus APO-RISPERIDONE may be given with or without meals.

Risperidone is partly metabolised by CYP2D6 to 9-hydroxy risperidone, which has similar pharmacological activity to risperidone. Another metabolic pathway is N-dealkylation.

After oral administration to psychotic patients, risperidone is eliminated with a half-life of about 3 hours. The elimination half-life of 9-hydroxy risperidone and of risperidone plus 9-hydroxy risperidone is 24 hours.



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and Risperidone 1mg/ml oral solution

Steady-state of risperidone is reached within 1 day in most patients. Steady-state of 9-hydroxy risperidone is reached within 4-5 days of dosing. Risperidone plasma concentrations are dose-proportional within the therapeutic dose range.

Risperidone is rapidly distributed. The volume of distribution is 1-2 L/kg. In plasma, risperidone is bound to albumin and alpha₁-acid glycoprotein. The plasma protein binding of risperidone is 88%, while that of 9-hydroxy risperidone is 77%.

One week after administration, 70% of the dose is excreted in the urine and 14% in the faeces. In urine, risperidone plus 9-hydroxy risperidone represents 35-45% of the dose.

A single-dose study showed higher active plasma concentrations and a reduced clearance of risperidone plus 9-hydroxy risperidone by 30% in the elderly and 60% in patients with renal insufficiency. Risperidone plasma concentrations were normal in patients with liver insufficiency, but the unbound risperidone in plasma was increased by about 35%.

Indications

APO-RISPERIDONE is indicated for the treatment of schizophrenia and other psychotic disorders. These include first episode psychoses, acute schizophrenic exacerbations, chronic schizophrenia and other psychotic conditions, in which positive symptoms (such as hallucinations, delusions, thought disturbances, hostility, suspiciousness), and/or negative symptoms (such as blunted affect, emotional and social withdrawal, poverty of speech) are prominent.

APO-RISPERIDONE is also indicated for the treatment and long term control of mania in bipolar disorder. These episodes are characterised by symptoms such as elevated, expansive or irritable mood, inflated self esteem, decreased need for sleep, pressured speech, racing thoughts, distractibility, or poor judgement, including disruptive or aggressive behaviors.

APO-RISPERIDONE also alleviates affective symptoms (such as depression, guilt-feelings, anxiety) associated with schizophrenia. In addition, APO-RISPERIDONE also appears effective in maintaining the clinical improvement during continuation therapy in patients who have shown an initial response to treatment with this agent.

APO-RISPERIDONE is also indicated for the treatment of behavioural and psychological symptoms of dementia such as aggressiveness (verbal outburst, physical violence), activity disturbance (agitation, wandering) or psychotic symptoms.

APO-RISPERIDONE is also indicated for the treatment of conduct and other disruptive behaviour disorders in children (over 5 years), adolescents and adults with subaverage intellectual functioning or mental retardation, or average IQ, in whom destructive behaviours (e.g. aggression, impulsivity and self-injurious behaviours) are prominent. APO-RISPERIDONE is also effective in maintaining the clinical improvement during continuation therapy in children and adolescents who have shown an initial treatment response. Pharmacological treatment should be an integral part of a more comprehensive treatment program, including psychosocial and educational intervention. Treatment with APO-RISPERIDONE for patients with disruptive behaviour disorders should be initiated only in consultation with a specialist, including child and adolescent psychiatrists, paediatric neurologists, developmental paediatricians, or other physicians conversant in the diagnosis and treatment of conduct and other disruptive behaviour disorders.

APO-RISPERIDONE is indicated for the treatment of autism in children and adolescents.

Dosage and Administration

This product is not able to deliver all approved dose regimens.



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and Risperidone 1mg/ml oral solution

APO-RISPERIDONE may be given as tablets or as oral solution.

Schizophrenia

Switching from other antipsychotics

When medically appropriate, gradual discontinuation of the previous treatment while APO-RISPERIDONE therapy is initiated is recommended. Also if medically appropriate, when switching patients from depot antipsychotics, initiate APO-RISPERIDONE therapy in place of the next scheduled injection. The need for continuing existing anti-Parkinson medications should be re-evaluated periodically.

Adults

APO-RISPERIDONE may be given once daily or twice daily. Patients should start with 2 mg/day of APO-RISPERIDONE. The dose may be increased on the second day to 4 mg. From then on the dosage can be maintained unchanged, or further individualised, if needed. Most patients will benefit from daily doses between 4 and 6 mg. In some patients a slower titration phase and a lower starting and maintenance dose may be appropriate.

Doses above 10 mg/day have not been shown to be superior in efficacy to lower doses and may cause extrapyramidal symptoms. Since the safety of doses above 16 mg/day has not been evaluated, doses above this level should not be used.

A benzodiazepine may be added to APO-RISPERIDONE when additional sedation is required.

Elderly

A starting dose of 0.5 mg twice daily is recommended. This dosage can be individually adjusted with 0.5 mg twice daily increments to 1 - 2 mg twice daily.

APO-RISPERIDONE is well tolerated in the elderly.

Children

Experience is lacking in children aged less than 15 years.

Bipolar Mania

APO-RISPERIDONE should be administered on a once daily schedule, starting with 2mg. Dosage adjustments, if indicated, should occur at intervals of not less than 24 hours and in dosage increments of 1mg per day. A dosing range between 2 and 6mg per day is recommended. As with all symptomatic treatments, the continued use of APO-RISPERIDONE must be evaluated and justified on an ongoing basis.

Behavioral Disturbances in Patients with Dementia



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and Risperidone 1mg/ml oral solution

A starting dose of 0.25 mg twice daily is recommended. This dosage can be individually adjusted by increments of 0.25 mg twice daily, not more frequently than every other day, if needed. The optimum dose is 0.5 mg twice daily for most patients. Some patients, however, may benefit from doses up to 1 mg twice daily.

Once patients have reached their target dose, a once daily dosing regimen can be considered. As with all symptomatic treatments, the continued use of APO-RISPERIDONE must be evaluated and justified on an on-going basis.

Conduct and other disruptive behavior disorders

For Subjects \geq 50kg

A starting dose of 0.5mg once daily is recommended. This dosage can be individually adjusted by increments of 0.5mg once daily not more frequently than every other day, if needed. The optimum dose is 1mg once daily for most patients. Some patients, however, may benefit from 0.5mg once daily while others may require 1.5mg once daily.

For Subjects <50kg

A starting dose of 0.25mg once daily is recommended, which can be individually adjusted by increments of 0.25mg once daily not more frequently than every other day, if needed. The optimum dose is 0.5mg once daily for most patients, although some patients may benefit from 0.25mg once daily while others may require 0.75mg once daily.

As with all symptomatic treatments, the continued use of APO-RISPERIDONE must be evaluated and justified on an on-going basis.

Autism

APO-RISPERIDONE can be administered once or twice daily. Patients experiencing somnolence may benefit from a switch in dosing from once daily to either once daily at bedtime, or twice daily. APO-RISPERIDONE should be administered based on body weight. Dosing should begin at 0.25 mg or 0.5 mg/day based upon weight (see Table 2 below for relative weight categories). On Day 4 of treatment, the dose may be increased up to 0.5 mg or 1.0mg/day. This dose should be maintained and response assessed at approximately day 14. Only in patients not achieving sufficient clinical response should additional dose increases be considered. Dose increases may proceed at \geq 2-week intervals in increments of 0.25 mg for patients <20 kg or 0.5 mg for patients \geq 20 kg. Based upon current studies, the maximum dose studied did not exceed a total daily dose of 1.5 mg in patients <20 kg, 2.5 mg in patients \geq 20 kg and 3.5 mg in patients >45 kg.

The table of the maximum daily doses provides a reference for titration and dosing by weight based upon current studies, and may serve as a guide according to clinical need:



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Doses of APO-RISPERIDONE in Paediatric Patients with Autistic Disorder

Weight Categories	Days 1 - 3	Days 4 - 14+	Increments if dose increases are needed	Dose Range
Dose by Weight in mg/day				
< 20 kg	0.25 mg	0.5 mg	+0.25 mg ≥2 week intervals	at 0.5 mg-1.5 mg
≥ 20 kg	0.5 mg	1.0 mg	+0.5 mg ≥2 week intervals	at 1.0 mg-2.5 mg*
Dose Range in mg/kg/day				
All	0.01 mg/kg/d	0.02 mg/kg/d	+0.01 mg/kg/day at ≥2 week intervals	0.02 mg/kg/d-0.06 mg/kg/d

* Subjects weighing >45 kg may require higher doses: maximum dose studied was 3.5 mg/day.

Once sufficient response has been achieved and maintained consideration may be given to gradually lowering the dose to achieve optimum balance of effectiveness and tolerance. Clinical experience was limited in autistic adolescents and in autistic children with an IQ>84 as not many of these patients were included in the trials. As with all symptomatic treatments, the continued use of APO-RISPERIDONE in children and adolescents with autism must be evaluated and justified on an ongoing basis.

Renal and Hepatic Impairment

Patients with renal impairment have less ability to eliminate risperidone plus 9-hydroxy risperidone than normal adults. Patients with impaired hepatic function have increases in plasma concentration of the unbound risperidone.

Irrespective of the indication, starting and consecutive dosing should be halved, and dose titration should be slower for patients with renal or hepatic impairment.

APO-RISPERIDONE should be used with caution in these groups of patients.

Contraindications

APO-RISPERIDONE is contraindicated in patients with a known hypersensitivity to the product.

Warnings and Precautions

Warnings

Elderly Patients with Dementia

Overall Mortality



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and Risperidone 1mg/ml oral solution

Elderly patients with dementia treated with atypical antipsychotic drugs have an increased mortality compared to placebo in a meta-analysis of 17 controlled trials of atypical antipsychotic drugs, including APO-RISPERIDONE. In placebo-controlled trials with APO-RISPERIDONE in this population, the incidence of mortality was 4.0% (40/1009) for APO-RISPERIDONE treated patients compared to 3.1% (22/712) for placebo-treated patients. The mean age (range) of patients who died was 86 years (range 67-100).

Concomitant use with Frusemide

In the APO-RISPERIDONE placebo-controlled trials in elderly patients with dementia, a higher incidence of mortality was observed in patients treated with frusemide plus Risperidone (7.3% [15/206]; mean age 89 years, range 75-97) when compared to patients treated with risperidone alone (3.1% [25/803]; mean age 84 years, range 70-96) or frusemide alone (4.1% [5/121]; mean age 80 years, range 67-90). The increase in mortality in patients treated with frusemide plus risperidone was observed in two of the four clinical trials.

No pathophysiological mechanism has been clearly identified to explain this finding, and no consistent pattern for cause of death was observed. Nevertheless, caution should be exercised and the risks and benefits of this combination should be considered prior to the decision to use. There was no increased mortality among patients taking other diuretics concomitantly with risperidone. Irrespective of treatment, dehydration was an overall risk factor for mortality and should therefore be carefully avoided in elderly patients with dementia.

Cerebrovascular Adverse Events

In placebo-controlled trials in elderly patients with dementia, there was a significantly higher incidence of cerebrovascular adverse events, such as stroke (including fatalities) and transient ischaemic attacks in patients (mean age 85 years, range 73-97) treated with APO-RISPERIDONE compared with patients treated with placebo. The pooled data from six placebo-controlled studies in mainly elderly patients (>65 years of age) with dementia showed that cerebrovascular adverse events (serious and non-serious combined) occurred in 3.3% (33/989) of patients treated with risperidone and 1.2% (8/693) of patients treated with placebo. The Odds Ratio (95% exact confidence interval) was 2.96 (1.33, 7.45).

Risperidone may interfere with activities requiring mental alertness. Therefore, patients should be advised not to drive or operate machinery until their individual susceptibility is known.

Precautions

Due to the alpha-blocking activity of Risperidone, orthostatic hypotension can occur, especially during the initial dose-titration period. APO-RISPERIDONE should be used with caution in patients with known cardiovascular disease (eg. heart failure, myocardial infarction, conduction abnormalities, dehydration, hypovolaemia, or cerebrovascular disease), and the dosage should be gradually titrated as recommended (see **DOSAGE AND ADMINISTRATION**). A dose reduction should be considered if hypotension occurs. Special care should be taken in patients taking medications to lower blood pressure.

Tardive Dyskinesia

Medicines with dopamine receptor antagonistic properties have been associated with the induction of tardive dyskinesia characterised by rhythmical involuntary movements, predominantly of the tongue and/or face. It has been reported that the occurrence of extrapyramidal symptoms



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and Risperidone 1mg/ml oral solution

is a risk factor for the development of tardive dyskinesia. Because APO-RISPERIDONE has a lower potential to induce extrapyramidal symptoms than classic neuroleptics, it should have a reduced risk of inducing tardive dyskinesia. If signs and symptoms of tardive dyskinesia appear, the discontinuation of all antipsychotic medicines should be considered.

Neuroleptic Malignant Syndrome

The Neuroleptic Malignant Syndrome, characterised by hyperthermia, muscle rigidity, autonomic instability, altered consciousness and elevated creatine phosphokinase (CPK) levels has been reported to occur with classical neuroleptics. In this event, all antipsychotic medicines, including APO-RISPERIDONE, should be discontinued.

Physicians should weigh the risks versus benefits when prescribing antipsychotics including APO-RISPERIDONE to patients with Parkinson's disease or Dementia with Lewy Bodies (DLB) since both groups may be at increased risk of Neuroleptic Malignant Syndrome as well as having an increased sensitivity to antipsychotic medications. Manifestation of this increased sensitivity can include confusion, obtundation, postural instability with frequent falls, in addition to extrapyramidal symptoms.

Hyperglycaemia and Diabetes Mellitus

Hyperglycaemia, in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, has been reported in patients treated with atypical antipsychotics including APO-RISPERIDONE. Assessment of the relationship between atypical antipsychotic use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Given these confounders, the relationship between atypical antipsychotic use and hyperglycaemia related adverse events is not completely understood. However, epidemiological studies suggest an increased risk of treatment-emergent hyperglycaemia-related adverse events in patients treated with atypical antipsychotics. Precise risk estimates for hyperglycaemia related adverse events in patients treated with atypical antipsychotics are not available.

Patients with an established diagnosis of diabetes mellitus who are started on atypical antipsychotics should be monitored regularly for worsening of glucose control. Patients with risk factors for diabetes mellitus (e.g. obesity, family history of diabetes) who are starting treatment with atypical antipsychotics should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics should be monitored for symptoms of hyperglycaemia including polydipsia, polyuria, polyphagia and weakness. Patients who develop symptoms of hyperglycaemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some cases, hyperglycaemia has resolved when the atypical antipsychotic was discontinued; however, some patients required continuation of antidiabetic treatment despite discontinuation of the suspect drug.

QT Interval

As with other antipsychotics, caution should be exercised when APO-RISPERIDONE is prescribed in patients with a history of cardiac arrhythmias, in patients with congenital long QT syndrome, and in concomitant use with drugs known to prolong the QT interval.

Dysphagia



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and Risperidone 1mg/ml oral solution

Oesophageal dysmotility and aspiration have been associated with antipsychotic drug use. Aspiration pneumonia is a common cause of morbidity and mortality in patients with advanced Alzheimer's dementia. APO-RISPERIDONE and other antipsychotic drugs should be used cautiously in patients at risk for aspiration pneumonia.

Other Precautions

Classical neuroleptics are known to lower the seizure threshold. Caution is recommended when treating patients with epilepsy.
Patients may be advised to refrain from excessive eating in view of the possibility of weight gain.

Use in the elderly

It is recommended to halve both the starting dose and the subsequent dose increments in geriatric patients.

Use in renal impairment

It is recommended to halve both the starting dose and the subsequent dose increments in patients with renal insufficiency.

Use in hepatic impairment

It is recommended to halve both the starting dose and the subsequent dose increments in patients with hepatic insufficiency.

Pregnancy and Lactation

The safety of APO-RISPERIDONE for use during human pregnancy has not been established. Reversible extrapyramidal symptoms in the neonate were observed following postmarketing use of risperidone during the last trimester of pregnancy. Although, in experimental animals, Risperidone did not show direct reproductive toxicity, some indirect, prolactin- and CNS-mediated effects were observed. No teratogenic effect of risperidone was noted in any study. Therefore, APO-RISPERIDONE should only be used during pregnancy if the benefits outweigh the risks. In animal studies, risperidone and 9-hydroxy risperidone are excreted in the milk. It has been demonstrated that this excretion also occurs in human breast milk. Therefore, women receiving APO-RISPERIDONE should not breast feed.

Paediatric Use

APO-RISPERIDONE had no adverse effects on cognitive function in paediatric patients. In combined, long-term, open-label trials, mean changes in cognitive function tests were small and did not increase or decrease over time.

A mean increase of 7.5 kg after 12 months of APO-RISPERIDONE treatment was observed, somewhat higher than the expected weight gain (approximately 3 to 3.5 kg per year) for children predominantly between 5 and 12 years of age.

APO-RISPERIDONE treatment for up to 3 years showed no adverse effects on growth and sexual maturation. No differences were observed between risperidone and placebo groups in measurements of sexual maturation, using the Tanner scale, and no adverse events suggestive



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and Risperidone 1mg/ml oral solution

of delayed pubertal maturation were reported. The mean change in height after 1 year of treatment with risperidone was within the expected growth range in this population.

Experience of risperidone treatment in children with schizophrenia aged less than 15 years is lacking. Experience is lacking in children with conduct and other disruptive behavior disorders aged less than 5 years. Experience is lacking in children with autism aged less than 5 years.

Adverse Effects

Clinical Trial Data

The safety of risperidone was evaluated from a clinical trial database consisting of 9712 patients exposed to one or more doses of risperidone for the treatment of various psychiatric disorders in adults, elderly patients with dementia, and pediatrics. Of these 9712 patients, 2626 were patients who received risperidone while participating in double-blind, placebo-controlled trials. The conditions and duration of treatment with risperidone varied greatly and included (in overlapping categories) double-blind, fixed- and flexible-dose, placebo- or active-controlled studies and open-label phases of studies, inpatients and outpatients, and short-term (up to 12 weeks) and longer-term (up to 3 years) exposures.

The majority of all adverse reactions were mild to moderate in severity.

Double-Blind, Placebo-Controlled Data - Adult Patients

Adverse drug reactions (ADRs) reported by $\geq 1\%$ of risperidone-treated adult patients in nine 3- to 8-week double-blind, placebo-controlled trials are shown in Table 1.

Table 1. Adverse Drug Reactions Reported by $\geq 1\%$ of risperidone-Treated Adult Patients in Double-Blind Placebo-Controlled Studies			
System/Organ Class Adverse Reaction	RISPERIDONE ≤ 8 mg/day (N=853) %	RISPERIDONE >8-16 mg/day (N=198) %	PLACEBO (N=687) %
Infections and Infestations			
Nasopharyngitis	2.1	4.0	1.7
Upper respiratory tract infection	1.5	2.5	1.5
Sinusitis	0.7	1.5	0.6
Urinary tract infection	0.5	2.5	0.1



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Blood and Lymphatic System Disorders			
Anaemia	0.1	1.0	0.1
Immune System Disorders			
Hypersensitivity	0.1	1.0	0.1
Psychiatric Disorders			
Insomnia	16.2	25.3	13.2
Anxiety	7.7	11.1	4.4
Nervousness	0.5	1.0	0.1
Nervous System Disorders			
Parkinsonism*	19.3	17.2	7.9
Akathisia*	9.8	10.1	2.7
Somnolence	6.8	1.5	2.0
Dizziness	6.3	3.5	3.9
Sedation	4.6	3.0	1.3
Tremor*	4.2	2.5	2.5
Dystonia*	3.8	3.5	1.0
Lethargy	2.6	0	1.3
Dizziness postural	1.2	0	0.1
Dyskinesia*	1.2	2.0	0.9
Syncope	0.4	1.0	0
Eye Disorders			
Vision blurred	2.1	1.0	0.7
Ear and Labyrinth Disorders			
Ear pain	0.1	1.0	0.3
Cardiac Disorders			
Tachycardia	1.1	2.5	0.1



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Vascular Disorders			
Orthostatic hypotension	1.3	0.5	0.1
Hypotension	0.2	1.0	0.3
Respiratory, Thoracic and Mediastinal Disorders			
Nasal congestion	2.0	6.1	1.3
Dyspnoea	0.8	2.0	0
Epistaxis	0.5	1.5	0.1
Sinus congestion	0.5	1.0	0.6
Gastrointestinal Disorders			
Nausea	6.4	4.0	2.6
Constipation	4.6	9.1	3.6
Dyspepsia	4.3	6.1	2.6
Vomiting	3.9	4.5	3.8
Diarrhoea	2.3	0.5	1.9
Salivary hypersecretion	2.3	1.0	0.4
Dry mouth	2.1	0	1.0
Abdominal discomfort	1.5	1.0	0.9
Abdominal pain	1.1	0.5	0.7
Stomach discomfort	1.1	1.0	0.6
Abdominal pain upper	0.7	1.0	0.1
Skin and Subcutaneous Tissue Disorders			
Rash	0.8	3.5	0.9
Dry skin	0.5	2.5	0.3



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Dandruff	0.2	1.0	0
Seborrhoeic dermatitis	0.2	1.0	0
Hyperkeratosis	0	1.0	0.3
Musculoskeletal and Connective Tissue Disorders			
Back pain	2.5	1.0	1.6
Arthralgia	1.5	2.5	0.6
Pain in extremity	1.2	1.0	2.2
Renal and Urinary Disorders			
Urinary incontinence	0.2	1.0	0.3
Reproductive System and Breast Disorders			
Ejaculation failure	0.4	1.0	0
General Disorders			
Fatigue	2.3	1.0	1.0
Asthenia	1.3	0.5	0.6
Pyrexia	1.3	1.0	0.7
Chest pain	0.8	1.5	0.4
Investigations			
Blood creatine phosphokinase increased	0.4	1.5	0.1
Heart rate increased	0.2	1.5	0.1

* Parkinsonism includes extrapyramidal disorder, musculoskeletal stiffness, Parkinsonism, cogwheel rigidity, akinesia, bradykinesia, hypokinesia, masked facies, muscle rigidity, and Parkinson's disease. Akathisia includes akathisia and restlessness. Dystonia includes dystonia, muscle spasms, muscle contractions involuntary, muscle contracture, oculogyration, tongue paralysis. Tremor includes tremor and Parkinsonian rest tremor. Dyskinesia includes dyskinesia, muscle twitching, chorea, and choreoathetosis.



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Double-Blind, Placebo-Controlled Data - Elderly Patients with Dementia

Adverse drug reactions (ADRs) reported by $\geq 1\%$ of risperidone-treated elderly patients with dementia in six 4- to 12-week double-blind, placebo-controlled trials are shown in Table 2. Table 2 includes only those ADRs that are either not listed in Table 1 or those ADRs that occurred at ≥ 2 times the frequency of the ADRs listed in Table 1.

Table 2. Adverse Drug Reactions (ADRs) Reported by $\geq 1\%$ of risperidone-Treated Elderly Patients with Dementia in Double-Blind Placebo-Controlled Studies: ADRs Not Listed in Table 1 or Reported at ≥ 2 Times the Frequency of ADRs Listed in Table 1.

System/Organ Class Adverse Reaction	RISPERIDONE (N=1009) %	PLACEBO (N=712) %
Infections and Infestations		
Urinary tract infection	12.9	10.3
Pneumonia	3.1	2.4
Cellulitis	1.1	1.3
Metabolism and Nutrition Disorders		
Decreased appetite	2.3	1.4
Psychiatric Disorders		
Confusional state	2.7	0.1
Nervous System Disorders		
Lethargy	7.6	2.2
Transient ischaemic attack	1.6	0.6
Depressed level of consciousness	1.3	0.3
Drooling	1.3	0
Cerebrovascular accident	1.1	0.4
Eye Disorders		
Conjunctivitis	2.7	1.1
Vascular Disorders		



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Hypotension	2.2	1.4
Respiratory, Thoracic and Mediastinal Disorders		
Cough	4.6	3.1
Rhinorrhoea	1.5	0.8
Gastrointestinal Disorders		
Dysphagia	1.5	1.3
Faecaloma	1.1	0.4
Skin and Subcutaneous Tissue Disorders		
Erythema	4.0	4.6
Musculoskeletal and Connective Tissue Disorders		
Posture abnormal	1.8	0.8
Joint swelling	1.5	0.3
General Disorders		
Oedema peripheral	7.7	3.9
Pyrexia	4.0	1.8
Gait disturbance	3.5	1.5
Pitting oedema	1.5	0.3
Investigations		
Body temperature increased	2.6	0.8



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Double-Blind, Placebo-Controlled Data - Pediatric Patients

Adverse drug reactions (ADRs) reported by $\geq 1\%$ of risperidone-treated pediatric patients in eight 3- to 8-week double-blind, placebo-controlled trials are shown in Table 3. Table 3 includes only those ADRs that are either not listed in Table 1 or those ADRs that occurred at ≥ 2 times the frequency of the ADRs listed in Table 1.

Table 3. Adverse Drug Reactions (ADRs) Reported by $\geq 1\%$ of risperidone-Treated Pediatric Patients in Double-Blind Placebo-Controlled Studies: ADRs Not Listed in Table 1 or Reported at ≥ 2 Times the Frequency of ADRs Listed in Table 1.

System/Organ Class Adverse Reaction	RISPERIDONE ≤ 3 mg/day (N=344) %	RISPERIDONE $>3-6$ mg/day (N=95) %	PLACEBO (N=349) %
Infections and Infestations			
Upper respiratory tract infection	5.2	2.1	3.4
Rhinitis	3.5	1.1	3.2
Influenza	1.7	0	1.7
Metabolism and Nutrition Disorders			
Increased appetite	17.2	3.2	7.2
Psychiatric Disorders			
Middle insomnia	1.7	0	0.9
Listless	0.9	1.1	0
Nervous System Disorders			
Somnolence	26.5	15.8	7.7
Headache	22.4	21.1	14.9
Sedation	20.1	14.7	4.0
Dizziness	8.1	13.7	2.3
Tremor	6.1	8.4	1.1
Drooling	4.9	2.1	1.1
Dysarthria	1.5	1.1	0



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Disturbance of attention	0.9	1.1	0.6
Balance disorder	0.9	1.1	0
Hypersomnia	0.6	1.1	0.9
Cardiac Disorders			
Palpitations	0.6	2.1	0
Respiratory, Thoracic and Mediastinal Disorders			
Cough	8.7	3.2	6.6
Rhinorrhoea	4.9	2.1	3.4
Epistaxis	3.8	4.2	1.7
Pharyngolaryngeal pain	3.8	2.1	1.7
Pulmonary congestion	0.3	1.1	0.3
Gastrointestinal Disorders			
Vomiting	13.7	8.4	9.2
Abdominal pain upper	8.4	6.3	4.6
Diarrhoea	6.7	2.1	6.0
Salivary hypersecretion	3.5	6.3	0.9
Stomach discomfort	2.9	0	1.4
Abdominal pain	2.3	2.1	0.6
Skin and Subcutaneous Tissue Disorders			
Pruritus	1.2	0	0
Acne	0.9	1.1	0
Musculoskeletal and Connective Tissue Disorders			



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Myalgia	1.2	1.1	0.9
Neck pain	0.3	1.1	0.3
Renal and Urinary Disorders			
Enuresis	6.4	1.1	5.2
Urinary incontinence	2.0	0	1.4
Pollakiuria	1.5	1.1	0.3
Reproductive System and Breast Disorders			
Galactorrhea	0.6	2.1	0
General Disorders			
Fatigue	19.2	18.9	4.9
Pyrexia	8.4	3.2	6.3
Feeling abnormal	1.2	0	0
Sluggishness	0.9	1.1	0
Chest discomfort	0.3	1.1	0
Investigations			
Weight increased	4.9	2.1	0.9
Blood prolactin increased	3.8	0	0.3

Other Clinical Trial Data

Adverse drug reactions (ADRs) reported in double-blind placebo-controlled clinical trials by < 1% of risperidone-treated adult or pediatric patients, or elderly patients with dementia, or at any rate by risperidone-treated patients in other studies, including double-blind, active-controlled and open-label studies are shown in Table 4.



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Table 4. Adverse Drug Reactions Reported in Double-Blind Placebo-Controlled Clinical Trials by <1% of risperidone-Treated Adult or Pediatric Patients, or Elderly Patients with Dementia, or at Any Rate by RISPERIDONE-Treated Patients in Other Studies, Including Double-Blind, Active-Controlled and Open-Label Studies

Infections and Infestations

Ear infection, Viral infection, Pharyngitis, Tonsillitis, Bronchitis, Eye infection, Localised infection, Cystitis, Otitis media, Onychomycosis, Acarodermatitis, Bronchopneumonia, Respiratory tract infection, Tracheobronchitis, Otitis media chronic

Blood and Lymphatic System Disorders

Granulocytopenia

Immune System Disorders

Drug hypersensitivity

Endocrine Disorders

Hyperprolactinemia

Metabolism and Nutrition Disorders

Polydipsia, Anorexia

Psychiatric Disorders

Agitation, Blunted affect, Sleep disorder, Libido decreased, Anorgasmia

Nervous System Disorders

Unresponsive to stimuli, Coordination abnormal, Loss of consciousness, Speech disorder, Hypoesthesia, Movement disorder, Tardive dyskinesia, Cerebral ischemia, Cerebrovascular disorder, Neuroleptic malignant syndrome, Diabetic coma

Eye Disorders

Ocular hyperemia, Eye discharge, Eye rolling, Eyelid edema, Eye swelling, Eyelid margin crusting, Dry eye, Lacrimation increased, Photophobia, Glaucoma, Visual acuity reduced

Ear and Labyrinth Disorders

Tinnitus

Cardiac Disorders

Sinus bradycardia, Sinus tachycardia, Palpitations, Atrioventricular block first degree, Bundle branch block left, Bundle branch block right, Atrioventricular block

Vascular Disorders



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Flushing

Respiratory, Thoracic, and Mediastinal Disorders

Wheezing, Pneumonia aspiration, Dysphonia, Productive cough, Respiratory tract congestion, Rales, Respiratory disorder, Nasal edema, Hyperventilation

Gastrointestinal Disorders

Fecal incontinence, Gastritis, Lip swelling, Cheilitis, Aptyalism

Skin and Subcutaneous Tissue Disorders

Skin discoloration, Skin lesion, Skin disorder, Rash erythematous, Rash papular, Rash generalised, Rash maculo-papular

Musculoskeletal and Connective Tissue Disorders

Musculoskeletal chest pain, Joint stiffness, Muscular weakness, Rhabdomyolysis

Renal and Urinary Disorders

Dysuria

Reproductive System and Breast Disorders

Menstruation irregular, Amenorrhea, Gynecomastia, Vaginal discharge, Erectile dysfunction, Ejaculation disorder, Menstrual disorder, Breast enlargement, Sexual dysfunction, Retrograde ejaculation

General Disorders

Thirst, Influenza-like illness, Edema, Malaise, Face edema, Discomfort, Generalised edema, Chills, Peripheral coldness, Drug withdrawal syndrome, Adverse drug reaction

Investigations

Alanine aminotransferase increased, Electrocardiogram abnormal, Eosinophil count increased, Aspartate aminotransferase increased, White blood cell count decreased, Blood glucose increased, Hemoglobin decreased, Hematocrit decreased, Body temperature decreased, Blood pressure decreased, Transaminases increased

Postmarketing Data

Adverse events first identified as ADRs during postmarketing experience with risperidone are included in Tables 5. The frequencies are provided according to the following convention:

Very common $\geq 1/10$

Common $\geq 1/100$ to $<1/10$

Uncommon $\geq 1/1,000$ to $<1/100$

Rare $\geq 1/10,000$ to $<1/1,000$

Very rare $<1/10,000$, including isolated reports



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and Risperidone 1mg/ml oral solution

In Table 5, ADRs are presented by frequency category based on spontaneous reporting rate.

Table 5. Adverse Drug Reactions Identified During Postmarketing Experience with risperidone by Frequency Category Estimated from Spontaneous Reporting Rates

Blood and Lymphatic Disorders	
<i>Very rare</i>	Agranulocytosis
<i>Very rare</i>	Thrombocytopenia ^a
Immune System Disorders	
<i>Very rare</i>	Anaphylactic reaction
Endocrine Disorders	
<i>Very rare</i>	Inappropriate antidiuretic hormone secretion
Metabolism and Nutrition Disorders	
<i>Very rare</i>	Diabetic ketoacidosis
<i>Very rare</i>	Water intoxication
Psychiatric Disorders	
<i>Very rare</i>	Mania
Cardiac Disorders	
<i>Very rare</i>	Atrial fibrillation
Respiratory, Thoracic, and Mediastinal Disorders	
<i>Very rare</i>	Sleep apnea syndrome
Gastrointestinal Disorders	
<i>Very rare</i>	Intestinal obstruction
<i>Very rare</i>	Pancreatitis
Hepatobiliary Disorders	
<i>Very rare</i>	Jaundice
Skin and Subcutaneous Tissue Disorders	
<i>Very rare</i>	Angioedema ^b
<i>Very rare</i>	Alopecia
Reproductive System and Breast Disorders	
<i>Very rare</i>	Priapism
General Disorders	
<i>Very rare</i>	Hypothermia
Investigations	
<i>Very rare</i>	Electrocardiogram QT prolonged ^c

^a Search terms included Thrombocytopenia, Platelet count decreased, Plateletcrit decreased, Platelet production decreased.



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and Risperidone 1mg/ml oral solution

^b Search terms included Angioneurotic oedema, C1 esterase deficiency acquired, Circumoral oedema, Eyelid edema, Face edema, Hereditary angioedema, Laryngeal oedema, Laryngotracheal oedema, Oculo-respiratory syndrome, Oedema mouth, Periorbital edema, Small bowel angioedema, Tongue oedema.

^c Search terms included Electrocardiogram QT corrected interval prolonged, Electrocardiogram QT interval abnormal, Electrocardiogram QT prolonged, Long QT syndrome, Long QT syndrome congenital.

Very rare cases of cardiac arrest, ventricular arrhythmias, torsades de pointes and sudden unexplained death have also been reported in postmarketing experience.

Interactions

Given the primary CNS effects of risperidone, it should be used with caution in combination with other centrally-acting medicines.

Caution is advised when prescribing risperidone with drugs known to prolong the QT interval, drugs causing electrolyte imbalance or metabolic inhibitors.

Carbamazepine has been shown to decrease the plasma levels of risperidone plus 9-hydroxy risperidone. Similar effects may be observed with other CYP 3A4 hepatic enzyme inducers. When carbamazepine or other CYP 3A4 hepatic enzyme inducers are initiated or discontinued, the physician should re-evaluate the dosing of risperidone.

Fluoxetine and paroxetine, CYP 2D6 inhibitors, increase the plasma concentration of risperidone, but less so of risperidone plus 9-hydroxy risperidone. When concomitant fluoxetine or paroxetine is initiated or discontinued, the physician should re-evaluate the dosing of risperidone.

Topiramate modestly reduced the bioavailability of risperidone, but not that of risperidone plus 9-hydroxy risperidone. Therefore, this interaction is unlikely to be of clinical significance.

Phenothiazines, tricyclic antidepressants and some beta-blockers may increase the plasma concentrations of risperidone but not those of risperidone plus 9-hydroxy risperidone. Amitriptyline does not affect the pharmacokinetics of risperidone or risperidone plus 9-hydroxy risperidone. Cimetidine and ranitidine increased the bioavailability of risperidone but only marginally that of risperidone plus 9-hydroxy risperidone.

Erythromycin, a CYP3A4 inhibitor, does not change the pharmacokinetics of risperidone and risperidone plus 9-hydroxy risperidone. The cholinesterase inhibitors galantamine and donepezil do not show a clinically relevant effect on the pharmacokinetics of risperidone and risperidone plus 9-hydroxy risperidone. When risperidone is taken together with other highly protein-bound medicines, there is no clinically relevant displacement of either medicine from the plasma proteins.

Risperidone does not show a clinically relevant effect on the pharmacokinetics of lithium, valproate, digoxin or topiramate.

See Precautions section (Elderly Patients with Dementia) regarding increased mortality in elderly dementia patients concomitantly receiving frusemide.

Food does not affect the absorption of risperidone.

Overdosage

Symptoms



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and Risperidone 1mg/ml oral solution

In general, reported signs and symptoms have been those resulting from an exaggeration of the known pharmacological effects of risperidone. These include drowsiness and sedation, tachycardia and hypotension, and extrapyramidal symptoms. Overdosages of up to 360 mg have been reported. The available evidence suggests a wide safety margin. In overdose, rare cases of QT-prolongation have been reported.

In case of acute overdosage, the possibility of multiple medicine involvement should be considered.

Treatment

Establish and maintain a clear airway and ensure adequate oxygenation and ventilation. Gastric lavage (after intubation, if the patient is unconscious) and administration of activated charcoal together with a laxative should be considered. Cardiovascular monitoring should commence immediately and should include continuous electrocardiographic monitoring to detect possible arrhythmias.

There is no specific antidote to risperidone. Therefore appropriate supportive measures should be instituted. Hypotension and circulatory collapse should be treated with appropriate measures such as intravenous fluids and/or sympathomimetic agents. In case of severe extrapyramidal symptoms, anticholinergic medication should be administered. Close medical supervision and monitoring should continue until the patient recovers.

Pharmaceutical Precautions

Shelf Life

APO-RISPERIDONE 0.5 mg, 1 mg, 2 mg, 3 mg and 4 mg Tablets 36 months when stored at or below 25°C in a dry place and protected from light.

APO-RISPERIDONE 1mg/ml Oral Solution 18 months when stored at or below 25°C. Do not refrigerate or freeze.

Medicine Classification

Prescription Only Medicine

Package Quantities

APO-RISPERIDONE 0.5 mg, 1 mg, 2 mg, 3 mg and 4mg tablets are available as blisters in a carton of 60.

APO-RISPERIDONE 1mg/ml Oral Solution:

Bottles of 30 & 100ml with a 3ml oral syringe enclosed.

Further Information

Tablets contain Lactose and Corn Starch



APO-RISPERIDONE

Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg tablets and
Risperidone 1mg/ml oral solution

Name and Address

Apotex NZ Ltd
32 Hillside Road
Glenfield
Private Bag 102-995
North Shore
North Shore City 0745
Auckland
Telephone: (09) 444 2073
Fax: (09) 444 2951

Date of Preparation

26 August 2010