

# DATA SHEET

## **VOLTFAST<sup>®</sup>** **Diclofenac Potassium** **50 mg Powder for Oral Solution**

---

### **Qualitative and quantitative composition**

---

The active ingredient is potassium-[o-[(2,6-dichlorophenyl)-amino]-phenyl]-acetate (= diclofenac potassium). One Voltfast<sup>®</sup> sachet contains 50 mg of diclofenac potassium.

For a full list of excipients, see List of excipients.

---

### **Pharmaceutical form**

---

Powder for oral solution.

Homogenous, white to light yellow powder.

---

### **Clinical particulars**

---

#### ***Therapeutic indications***

Short-term treatment in the following acute conditions:

- Post-traumatic pain, inflammation and swelling, e.g. due to sprains.
- Post-operative pain, inflammation and swelling, e.g. following dental or orthopaedic surgery.
- Painful and/or inflammatory conditions in gynaecology, e.g. primary dysmenorrhoea or adnexitis.
- Migraine attacks.
- Painful syndromes of the vertebral column.
- Non-articular rheumatism.
- As an adjuvant in severe painful inflammatory infections of the ear, nose, or throat, e.g. pharyngotonsillitis, otitis. In keeping with general therapeutic principles, the underlying disease should be treated with basic therapy, as appropriate. Fever alone is not an indication.

#### ***Dosage and method of administration***

After assessing the risk/benefit ratio in each individual patient, the lowest effective dose for the shortest possible duration should be used.

The contents of the sachet should be dissolved with stirring in a glass of natural (non-carbonated) water. The solution may remain slightly opalescent, but this should not influence the efficacy of the preparation. The solution should be swallowed preferably before meals.

## **Adults**

The recommended initial daily dose is 100 to 150 mg. In milder cases, 50 to 100 mg daily are usually sufficient. The daily dose should generally be divided in up to 3 doses.

In primary dysmenorrhoea, the daily dose should be individually adjusted and is generally 50 to 150 mg. A dose of 50 to 100 mg should be given initially and, if necessary increased over the course of several menstrual cycles up to a total maximum of 200 mg/day. Treatment should be started on appearance of the first symptoms and, depending on the symptomatology, continued for a few days.

In migraine, an initial dose of 50 mg should be taken at the first signs of an impending attack. In cases where pain relief within 2 hours after the first dose is not sufficient, a further dose of 50 mg may be taken. If needed, further doses of 50 mg may be taken at intervals of 4 to 6 hours, not exceeding a total dose of 200 mg per day.

## **Children and adolescents**

Voltfast 50 mg powder for oral solution is not recommended for use in children and adolescents below 14 years of age. For treatment in children and adolescents below 14 years of age, oral drops and suppositories of diclofenac 12.5 mg and 25 mg are available.

For adolescents aged 14 years and over, 50 to 100 mg daily are usually sufficient, given as 1 to 2 divided doses.

The maximum daily dose of 150 mg should not be exceeded.

The use of Voltfast 50 mg powder for oral solution in migraine attacks has not been established in children and adolescents.

## ***Contraindications***

Known hypersensitivity to the active substance or to any of the excipients.

Active gastric or intestinal ulcer, bleeding or perforation.

Last trimester of pregnancy (see Pregnancy and lactation).

Severe hepatic, renal or cardiac failure (see Special warnings and precautions for use).

Like other non-steroidal anti-inflammatory drugs (NSAIDs), Voltfast is also contraindicated in patients in whom attacks of asthma, urticaria, or acute rhinitis are precipitated by acetylsalicylic acid or other NSAIDs.

## ***Special warnings and precautions for use***

### **Warnings**

#### **Cardiovascular thrombotic events:**

Observational studies have indicated that non-selective NSAIDs may be associated with an increased risk of serious cardiovascular events including myocardial infarction and stroke, which may increase with dose or duration of use. Patients with cardiovascular disease or cardiovascular risk factors may also be at greater risk. To minimise the potential risk of an adverse cardiovascular event in patients taking an NSAID, especially in those with cardiovascular risk factors, the lowest effective dose should be used for the shortest possible duration (see 'Dosage and administration').

There is no consistent evidence that the concurrent use of aspirin mitigates the possible increased risk of serious cardiovascular thrombotic events associated with NSAID use.

**Hypertension:**

NSAIDs may lead to the onset of new hypertension or worsening of pre-existing hypertension and patients taking anti-hypertensives with NSAIDs may have an impaired anti-hypertensive response. Caution is advised when prescribing NSAIDs to patients with hypertension. Blood pressure should be monitored closely during initiation of NSAID treatment and at regular intervals thereafter.

**Heart failure**

Fluid retention and oedema have been observed in some patients taking NSAIDs, therefore caution is advised in patients with fluid retention or heart failure.

**Gastrointestinal effects:**

Gastrointestinal bleeding, ulceration or perforation, which may increase with dose or duration of use and which can be fatal, have been reported with all NSAIDs, including diclofenac, and may occur at any time during treatment, with or without warning symptoms or a previous history of serious gastrointestinal events. They generally have more serious consequences in the elderly. If gastrointestinal bleeding or ulceration occurs in patients receiving Voltfast, the medicinal product should be withdrawn.

Upper GI ulcers, gross bleeding or perforation caused by NSAIDs occur in approximately 1% of patients treated for 3-6 months and in about 2-4% of patients treated for one year. These trends continue with longer duration of use, increasing the likelihood of developing a serious GI event at some time during the course of therapy. However, even short-term therapy is not without risk.

Caution is advised in patients with risk factors for gastrointestinal events who may be at greater risk of developing serious gastrointestinal events, e.g. the elderly, those with a history of serious gastrointestinal events, smoking and alcoholism.

The concurrent use of aspirin and NSAIDs also increases the risk of serious gastrointestinal adverse events.

Doctors should warn patients about the signs and symptoms of serious gastrointestinal toxicity.

**Severe skin reactions**

Serious skin reactions, some of them fatal, including exfoliative dermatitis, Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TENS), have been reported very rarely in association with the use of NSAIDs, including Voltfast (see Adverse effects). These serious adverse events are idiosyncratic and are independent of dose or duration of use. Patients appear to be at highest risk of these reactions early in the course of therapy, the onset of the reaction occurring in the majority of cases within the first month of treatment. Patients should be advised of the signs and symptoms of serious skin reactions and to consult their doctor at the first appearance of skin rash, mucosal lesions or any other sign of hypersensitivity, and Voltfast should be discontinued.

As with other NSAIDs, allergic reactions, including anaphylactic/anaphylactoid reactions, can also occur in rare cases with diclofenac without earlier exposure to the drug.

Like other NSAIDs, Voltfast may mask the signs and symptoms of infection due to its pharmacodynamic properties.

**Precautions****General**

The concomitant use of Voltfast with systemic NSAIDs including cyclooxygenase-2 selective inhibitors, should be avoided due to the absence of any evidence demonstrating synergistic benefits and the potential for additive adverse effects.

Caution is indicated in the elderly on basic medical grounds. In particular it is recommended that the lowest effective dose be used in frail elderly patients or those with a low body weight.

Voltfast contains a source of phenylalanine and may be therefore harmful for patients with phenylketonuria.

### **Pre-existing asthma**

In patients with asthma, seasonal allergic rhinitis, swelling of the nasal mucosa (i.e. nasal polyps), chronic obstructive pulmonary diseases or chronic infections of the respiratory tract (especially if linked to allergic rhinitis-like symptoms), reactions on NSAIDs like asthma exacerbations (so-called intolerance to analgesics / analgesics-asthma), Quincke's oedema or urticaria are more frequent than in other patients. Therefore, special precaution is recommended in such patients (readiness for emergency). This is applicable as well for patients who are allergic to other substances, e.g. with skin reactions, pruritus or urticaria.

### **Gastrointestinal effects**

As with all NSAIDs, including diclofenac, close medical surveillance is imperative and particular caution should be exercised when prescribing Voltfast in patients with symptoms indicative of gastrointestinal (GI) disorders or with a history suggestive of gastric or intestinal ulceration, bleeding or perforation (see Adverse effects). The risk of GI bleeding is higher with increasing NSAID doses and in patients with a history of ulcer, particularly if complicated with haemorrhage or perforation and in the elderly.

To reduce the risk of GI toxicity in patients with a history of ulcer, particularly if complicated with haemorrhage or perforation, and in the elderly, the treatment should be initiated and maintained at the lowest effective dose.

Combination therapy with protective agents (e.g. proton pump inhibitors or misoprostol) should be considered for these patients, and also for patients requiring concomitant use of medicinal products containing low-dose acetylsalicylic acid (ASA)/aspirin or other medicinal products likely to increase gastrointestinal risk.

Patients with a history of GI toxicity, particularly the elderly, should report any unusual abdominal symptoms (especially GI bleeding). Caution is recommended in patients receiving concomitant medications which could increase the risk of ulceration or bleeding, such as systemic corticosteroids, anticoagulants, anti-platelet agents or selective serotonin-reuptake inhibitors (see Interaction with other medicinal products and other forms of interaction).

Close medical surveillance and caution should also be exercised in patients with ulcerative colitis or Crohn's disease, as their condition may be exacerbated (see Adverse effects).

### **Hepatic effects**

Close medical surveillance is required when prescribing Voltfast to patients with impaired hepatic function, as their condition may be exacerbated.

As with other NSAIDs, including diclofenac values of one or more liver enzymes may increase. During prolonged treatment with Voltfast, regular monitoring of hepatic function is indicated as a precautionary measure. If abnormal liver function tests persist or worsen, if clinical signs or symptoms consistent with liver disease develop, or if other manifestations occur (e.g. eosinophilia, rash), Voltfast should be discontinued. Hepatitis may occur with use of diclofenac without prodromal symptoms.

Caution is called for when using Voltfast in patients with hepatic porphyria, since it may trigger an attack.

### **Renal effects**

As fluid retention and oedema have been reported in association with NSAID therapy, including diclofenac, particular caution is called for in patients with impaired cardiac or renal function, history of hypertension, the elderly, patients receiving concomitant treatment with diuretics or medicinal products that can significantly impact renal function, and in those patients with substantial extracellular volume depletion from any cause, e.g. before and after major surgery (see Contraindications). Monitoring of renal function is recommended as a precautionary measure when using Voltfast in such cases. Discontinuation of therapy is normally followed by a recovery to the pre-treatment state.

### **Haematological effects**

Use of Voltfast is recommended only for short-term treatment. If, however, Voltfast is used for a prolonged period, monitoring of the blood count is recommended, as with other NSAIDs.

Like other NSAIDs, Voltfast may temporarily inhibit platelet aggregation. Patients with defects of haemostasis should be carefully monitored.

### ***Interaction with other medicinal products and other forms of interaction***

The following interactions include those observed with Voltfast powder for oral solution and/or other pharmaceutical forms of diclofenac.

***Lithium:*** If used concomitantly, diclofenac may raise plasma concentrations of lithium. Monitoring of the serum lithium level is recommended.

***Digoxin:*** If used concomitantly, diclofenac may raise plasma concentrations of digoxin. Monitoring of the serum digoxin level is recommended.

***Diuretics and antihypertensive agents:*** Like other NSAIDs, concomitant use of diclofenac with diuretics or antihypertensive agents (e.g. beta-blockers, angiotensin converting enzyme (ACE) inhibitors) may cause a decrease in their antihypertensive effect. Therefore, the combination should be administered with caution and patients, especially the elderly, should have their blood pressure periodically monitored. Patients should be adequately hydrated and consideration should be given to monitoring of renal function after initiation of concomitant therapy and periodically thereafter, particularly for diuretics and ACE inhibitors due to the increased risk of nephrotoxicity. Concomitant treatment with potassium-sparing drugs may be associated with increased serum potassium levels, which should therefore be monitored frequently (see Special warnings and precautions for use).

***Other NSAIDs and corticosteroids:*** Concomitant administration of diclofenac and other systemic NSAIDs or corticosteroids may increase the frequency of gastrointestinal adverse effects (see Special warnings and precautions for use).

***Anticoagulants and anti-platelet agents:*** Caution is recommended since concomitant administration could increase the risk of bleeding (see Special warnings and precautions for use). Although clinical investigations do not appear to indicate that diclofenac affects the action of anticoagulants, there are isolated reports of an increased risk of haemorrhage in patients receiving diclofenac and anticoagulants concomitantly. Close monitoring of such patients is therefore recommended.

***Selective serotonin reuptake inhibitors (SSRIs):*** Concomitant administration of systemic NSAIDs, including diclofenac, and SSRIs may increase the risk of gastrointestinal bleeding (see Special warnings and precautions for use).

***Antidiabetics:*** Clinical studies have shown that diclofenac can be given together with oral antidiabetic agents without influencing their clinical effect. However, there have been isolated reports of both hypoglycaemic and hyperglycaemic effects necessitating changes in the dosage of the antidiabetic agents during treatment with diclofenac. For this reason, monitoring of the blood glucose level is recommended as a precautionary measure during concomitant therapy.

***Methotrexate:*** Caution is recommended when NSAIDs, including diclofenac, are administered less than 24 hours before or after treatment with methotrexate, since blood concentrations of methotrexate may rise and the toxicity of this substance be increased.

***Cyclosporin:*** Diclofenac, like other NSAIDs, may increase the nephrotoxicity of cyclosporin due to the effect on renal prostaglandins. Therefore, it should be given at doses lower than those that would be used in patients not receiving cyclosporin.

***Quinolone antibacterials:*** There have been isolated reports of convulsions which may have been due to concomitant use of quinolones and NSAIDs.

**Potent CYP2C9 inhibitors:** Caution is recommended when co-prescribing diclofenac with potent CYP2C9 inhibitors (such as sulfinpyrazone and voriconazole), which could result in a significant increase in peak plasma concentrations and exposure to diclofenac due to inhibition of diclofenac metabolism.

**Phenytoin:** When using phenytoin concomitantly with diclofenac, monitoring of phenytoin plasma concentrations is recommended due to an expected increase in exposure to phenytoin.

## **Pregnancy and lactation**

### **Pregnancy**

The use of diclofenac in pregnant women has not been studied. Therefore, Voltfast should not be used during the first two trimesters of pregnancy unless the potential benefit to the mother outweighs the risk to the foetus. As with other NSAIDs, use of diclofenac during the third trimester of pregnancy is contraindicated owing to the possibility of uterine inertia and/or premature closure of the ductus arteriosus (see Contraindications). Animal studies have not shown any directly or indirectly harmful effects on pregnancy, embryonal/foetal development, parturition or postnatal development (see Preclinical safety data).

### **Lactation**

Like other NSAIDs, diclofenac passes into the breast milk in small amounts. Therefore, Voltfast should not be administered during breast feeding in order to avoid adverse effects in the infant.

### **Fertility**

As with other NSAIDs, the use of Voltfast may impair female fertility and is not recommended in women attempting to conceive. In women who have difficulties conceiving or who are undergoing investigation of infertility, withdrawal of Voltfast should be considered.

### **Effects on ability to drive and use machines**

Patients experiencing visual disturbances, dizziness, vertigo, somnolence or other central nervous system disturbances while taking Voltfast, should refrain from driving or using machines.

### **Adverse effects**

Adverse reactions (Table 1) are ranked under heading of frequency, the most frequent first, using the following convention: common ( $\geq 1/100$ ,  $< 1/10$ ); uncommon ( $\geq 1/1,000$ ,  $< 1/100$ ); rare ( $\geq 1/10,000$ ,  $< 1/1,000$ ); very rare ( $< 1/10,000$ ), including isolated reports.

The following adverse effects include those reported with Voltfast powder for oral solution and/or other pharmaceutical forms of diclofenac, with either short-term or long-term use.

#### **Table 1**

##### **Blood and lymphatic system disorders**

Very rare:	Thrombocytopenia, leukopenia, anaemia (including haemolytic and aplastic anaemia), agranulocytosis.
------------	---

##### **Immune system disorders**

Rare:	Hypersensitivity, anaphylactic and anaphylactoid reactions (including hypotension and shock).
-------	---

Very rare:	Angioneurotic oedema (including face oedema).
------------	---

**Psychiatric disorders**

Very rare: Disorientation, depression, insomnia, nightmare, irritability, psychotic disorder.

**Nervous system disorders**

Common: Headache, dizziness.

Rare: Somnolence.

Very rare: Paraesthesia, memory impairment, convulsion, anxiety, tremor, aseptic meningitis, taste disturbances, cerebrovascular accident.

**Eye disorders**

Very rare: Visual disturbance, vision blurred, diplopia.

**Ear and labyrinth disorders**

Common: Vertigo.

Very rare: Tinnitus, hearing impaired.

**Cardiac disorders**

Very rare: Palpitations, chest pain, cardiac failure, myocardial infarction.

**Vascular disorders**

Very rare: Hypertension, vasculitis.

**Respiratory, thoracic and mediastinal disorders**

Rare: Asthma (including dyspnoea).

Very rare: Pneumonitis.

**Gastrointestinal disorders**

Common: Nausea, vomiting, diarrhoea, dyspepsia, abdominal pain, flatulence, anorexia.

Rare: Gastritis, gastrointestinal haemorrhage, haematemesis, melaena, diarrhoea haemorrhagic, gastrointestinal ulcer (with or without bleeding or perforation).

Very rare: Colitis (including haemorrhagic colitis and exacerbation of ulcerative colitis or Crohn's disease), constipation, stomatitis, glossitis, oesophageal disorder, diaphragm-like intestinal strictures, pancreatitis.

**Hepatobiliary disorders**

Common: Transaminases increased.

Rare: Hepatitis, jaundice, liver disorder.

Very rare: Fulminant hepatitis, hepatic necrosis, hepatic failure.

**Skin and subcutaneous tissue disorders**

Common: Rash.

Rare: Urticaria.

Very rare: Bullous eruptions, eczema, erythema, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis (Lyell's syndrome), dermatitis exfoliative, loss of hair, photosensitivity reaction, purpura, allergic purpura, pruritus.

**Renal and urinary disorders**

Very rare: Acute renal failure, haematuria, proteinuria, nephrotic syndrome, interstitial nephritis, renal papillary necrosis.

**General disorders and administration site conditions**

Rare: Oedema.

## **Overdose**

### **Symptoms**

There is no typical clinical picture resulting from diclofenac overdosage. Overdosage can cause symptoms such as vomiting, gastrointestinal haemorrhage, diarrhoea, dizziness, tinnitus or convulsions. In the event of significant poisoning, acute renal failure and liver damage are possible.

### **Therapeutic measures**

Management of acute poisoning with NSAIDs, including diclofenac, essentially consists of supportive measures and symptomatic treatment. Supportive measures and symptomatic treatment should be given for complications such as hypotension, renal failure, convulsions, gastrointestinal disorder, and respiratory depression.

Special measures such as forced diuresis, dialysis or haemoperfusion are probably of no help in eliminating NSAIDs, including diclofenac, due to the high protein-binding and extensive metabolism.

Activated charcoal may be considered after ingestion of a potentially toxic overdose, and gastric decontamination (e.g. vomiting, gastric lavage) after ingestion of a potentially life-threatening overdose.

---

## **Pharmacological properties**

---

### ***Pharmacodynamic properties***

Pharmacotherapeutic group: Anti-inflammatory and antirheumatic products, non-steroids acetic acid derivatives and related substances (ATC code: M01A B05).

### **Mechanism of action**

Voltfast contains the potassium salt of diclofenac, a non-steroidal compound with pronounced analgesic, anti-inflammatory, and antipyretic properties.

Voltfast has a rapid onset of action, which makes it particularly suitable for the treatment of acute painful and inflammatory conditions. Inhibition of prostaglandin biosynthesis, which has been demonstrated in experiments, is considered to be fundamental to its mechanism of action. Prostaglandins play a major role in causing inflammation, pain, and fever.

Diclofenac potassium *in vitro* does not suppress proteoglycan biosynthesis in cartilage at concentrations equivalent to the concentrations reached in humans.

### **Pharmacodynamic effects**

Voltfast has been found to exert a pronounced analgesic effect in moderate and severe pain. In the presence of inflammation, e.g. due to trauma or following surgical interventions, it rapidly relieves both spontaneous pain and pain on movement and diminishes inflammatory swelling and wound oedema. Clinical studies have also revealed that in primary dysmenorrhoea the active substance is capable of relieving the pain and reducing the extent of bleeding. In migraine attacks Voltfast has been shown to be effective in relieving the headache and in improving the accompanying symptoms nausea and vomiting.

## ***Pharmacokinetic properties***

### **Absorption**

Diclofenac is rapidly and completely absorbed from diclofenac potassium. Mean peak plasma concentrations of 5.5 micromol/L are attained after 5 to 20 minutes after ingestion of one sachet of 50 mg.

Ingestion together with food is expected to have no influence on the amount of diclofenac absorbed although onset and rate of absorption may be slightly delayed.

The amount absorbed is in linear proportion to the size of the dose.

Since about half of diclofenac is metabolized during its first passage through the liver ("first pass" effect), the area under the concentration curve (AUC) is about half as large following oral or rectal administration as it is following a parenteral dose of equal size.

Pharmacokinetic behaviour does not change after repeated administration. No accumulation occurs provided the recommended dosage intervals are observed.

### **Distribution**

99.7 % of diclofenac binds to serum proteins, mainly to albumin (99.4 %). The apparent volume of distribution calculated is 0.12 to 0.17 L/kg.

Diclofenac enters the synovial fluid, where maximum concentrations are measured 2 to 4 hours after peak plasma values have been reached. The apparent half-life for elimination from the synovial fluid is 3 to 6 hours. Two hours after reaching peak plasma levels, concentrations of the active substance are already higher in the synovial fluid than in the plasma, and they remain higher for up to 12 hours.

### **Biotransformation**

Biotransformation of diclofenac takes place partly by glucuronidation of the intact molecule, but mainly by single and multiple hydroxylation and methoxylation, resulting in several phenolic metabolites (3'-hydroxy-, 4'-hydroxy-, 5-hydroxy-, 4',5-dihydroxy-, and 3'-hydroxy-4'-methoxy-diclofenac), most of which are converted to glucuronide conjugates. Two of these phenolic metabolites are biologically active, but to a much lesser extent than diclofenac.

### **Elimination**

Total systemic clearance of diclofenac from plasma is  $263 \pm 56$  mL/min (mean value  $\pm$  SD). The terminal half-life in plasma is 1 to 2 hours. Four of the metabolites, including the two active ones, also have short plasma half-lives of 1 to 3 hours. One metabolite, 3'-hydroxy-4'-methoxy-diclofenac, has a much longer plasma half-life. However, this metabolite is virtually inactive.

About 60 % of the administered dose is excreted in the urine as the glucuronide conjugate of the intact molecule and as metabolites, most of which are also converted to glucuronide conjugates. Less than 1 % is excreted as unchanged substance. The rest of the dose is eliminated as metabolites through the bile in the faeces.

### **Characteristics in patients**

No relevant age-dependent differences in the drug's absorption, metabolism or excretion have been observed.

In patients suffering from renal impairment, no accumulation of the unchanged active substance can be inferred from the single-dose kinetics when applying the usual dosage schedule. At a creatinine clearance of less than 10 mL/min, the calculated steady-state plasma levels of the hydroxy

metabolites are about 4 times higher than in normal subjects. However, the metabolites are ultimately cleared through the bile.

In patients with chronic hepatitis or non-decompensated cirrhosis, the kinetics and metabolism of diclofenac are the same as in patients without liver disease.

### ***Preclinical safety data***

Preclinical data from acute and repeated dose toxicity studies, as well as from genotoxicity, mutagenicity, and carcinogenicity studies with diclofenac revealed no specific hazard for humans at the intended therapeutic doses. There was no evidence that diclofenac had a teratogenic potential in mice, rats or rabbits.

Diclofenac had no influence on the fertility of parent animals in rats. The prenatal, perinatal and postnatal development of the offspring was not affected.

---

## **Pharmaceutical particulars**

---

### ***List of excipients***

Potassium hydrogen carbonate; mannitol; aspartame; saccharin sodium; glyceryl dibehenate; mint flavour; anise flavour.

### ***Incompatibilities***

Not applicable.

### ***Shelf life***

2 years.

### ***Special precautions for storage***

Do not store above 25°C.

Store in the original package in order to protect from moisture.

Voltfast powder for oral solution must be kept out of the reach and sight of children.

### ***Nature and contents of container***

Packs containing 9 sachets

### ***Instructions for use and handling***

No special requirements.

### ***Medicine classification***

Prescription Medicine

### ***Name and address***

Novartis New Zealand Limited  
Private Bag 65904

Mairangi Bay  
Auckland 0754  
Building G, 5 Orbit Drive  
Rosedale  
Auckland 0632

Telephone: 09 361 8100

***Date of preparation***

3 June 2010

(Ref: BPI 26 May2010)