

Sandimmun®

50mg / mL concentrate for solution for intravenous infusion
cyclosporin

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Sandimmun.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

The information in this leaflet was last updated on the date listed on the final page. More recent information on the medicine may be available.

You should ensure that you speak to your pharmacist or doctor to obtain the most up-to-date information on the medicine. You can also download the most up-to-date leaflet from

www.medsafe.govt.nz.

Those updates may contain important information about the medicine and its use of which you should be aware.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Sandimmun against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine. You may need to read it again.

What Sandimmun is used for

Sandimmun contains the active ingredient, cyclosporin. It belongs to a group of medicines called immunosuppressive agents. These medicines help to control your body's immune system.

Sandimmun is used for people who have had a bone marrow, kidney, heart, combined heart-lung, liver or pancreas transplant.

Sandimmun control your body's immune system and prevents the body from rejecting the new organ. It does this by blocking the development of special cells which would normally attack the transplanted tissue.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

Sandimmun is only available with a doctor's prescription.

It is not addictive.

Before you are given Sandimmun

When you must not have it

Do not have Sandimmun if you have ever had an allergic reaction to any of the following:

- cyclosporin, the active ingredient in Sandimmun
- Cremophor® EL, a castor oil derivative in Sandimmun concentrate for intravenous infusion
- (which may cause severe allergic reactions)
- Alcohol
- Any of the other ingredients of Sandimmun, listed at the end of this leaflet.

Symptoms of an allergic reaction may include hives or an itchy skin rash, swelling of the face, lips or

tongue, faintness, wheezing or troubled breathing.

If you think that you may be allergic to Sandimmun, ask your doctor for advice.

Experience with Sandimmun injection in children and adolescents is still limited. Children from 1 year of age have received Sandimmun in standard dosage with no particular problems. In several studies, children required and tolerated higher doses of Sandimmun per kg body weight than those used in adult.

Do not have Sandimmun after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

In that case, return it to your pharmacist.

Experience with Sandimmun injection in children is still limited. Children from 1 year of age have received Sandimmun in standard dosage with no particular problems. In several studies, children required and tolerated higher doses of Sandimmun per kg body weight than those used in adult.

Before you start to have it

If you have been prescribed Sandimmun, tell your doctor if you also have any of these health problems/medical conditions:

- high blood pressure that is not controlled
- any uncontrolled infection
- a poorly functioning immune system
- problems with your kidneys or liver
- severe heart, lung or blood vessel disease
- any type of cancer, including skin cancer

Your doctor may not want you to have Sandimmun or may want to

take special precautions if you have any of these conditions.

Tell your doctor if you are pregnant or intend to become pregnant.

Experience with Sandimmun in pregnancy is very limited. The use of immunosuppressant medicines, including cyclosporin, during pregnancy has been shown to increase the potential risk of problems in the mother and the unborn child. If it is necessary for you to take this medicine, your doctor will discuss with you the benefits and potential risks of taking it during pregnancy.

Tell your doctor if you are breast-feeding.

Breast-feeding is not recommended since cyclosporin, the active ingredient in Sandimmun, passes into breast milk and may affect your baby.

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop. Many other medicines may be affected by Sandimmun or they may affect how well Sandimmun works.

These include:

- St John's wort, an ingredient in many medicines that you can buy without a prescription from a pharmacy, health food shop or supermarket.
- methotrexate, a medicine to treat severe rheumatoid arthritis, some types of cancers, and psoriasis
- antibacterial amino glycoside-type agents (e.g. gentamicin, tobramycin)
- antifungal agents containing amphotericin B
- antibacterial agents containing ciprofloxacin
- cytostatics containing melphalan
- agents used to treat urinary tract infection containing trimethoprim
- medicines used to treat pain (non-steroid anti-inflammatory drugs, e.g. diclofenac)
- acid secretion inhibitors of the H₂-receptor antagonist type (e.g. cimetidine, ranitidine), which are used to reduce the amount of acid in your stomach
- other drugs which may affect the kidneys
- antibacterial agents of the macrolide type (e.g. clarithromycin, azithromycin, erythromycin)
- antifungal agents of the azole type (e.g. fluconazole, itraconazole, ketoconazole, voriconazole) or terbinafine
- oral contraceptives (Levonorgestrel and Norethisterone)
- protease inhibitors, used to treat or prevent infections caused by viruses
- imatinib, a medicine used to treat some cancers
- anthracycline anticancer medicines, such as doxorubicine
- certain blood pressure reducing agents of the calcium antagonist type (e.g. nifedipine, amlodipine, verapamil, diltiazem, amiodarone) or of the endothelin receptor antagonist type (e.g. bosentan, ambrisentan)
- certain anticonvulsives, used to prevent fits or seizures (e.g. carbamazepine, phenytoin)
- digoxin, a medicine used to treat heart failure
- colchicine, used to treat gout disease with painful, swollen joints
- HMG-CoA reductase inhibitors (statins), or fibric acid derivatives, which are used to treat high cholesterol
- prednisolone, a corticosteroid used to treat inflammatory diseases such as arthritis, asthma, colitis
- etoposide, used to treat cancer
- repaglinide, used to treat Type II diabetes
- aliskiren, used to treat high blood pressure
- dabigatran, used to prevent stroke
- potassium sparing drugs or potassium containing drugs
- triclopidine (a medicine that is used after a stroke)
- octreotide, a medicine used to treat excess growth hormone, relieve the symptoms of certain types of cancer, or having surgery on the pancreas
- orlistat, used to help with weight loss
- danazol, a medicine used to treat menstrual disorders
- allopurinol, a medicine used to treat gout (a disease with painful, swollen joints caused by uric acid crystals)
- metoclopramide, a medicine used to stop prevent nausea and vomiting
- cholic acid and derivatives, which are used to treat gallstones
- tacrolimus, everolimus, or sirolimus, which are medicines that lower your immunity
- mycophenolate sodium or mofetil used in transplant patients

You may need to take different amounts of your medicines or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking Sandimmun.

If you have not told your doctor about any of these things, tell him/her before you have Sandimmun.

How Sandimmun is given

You will be given Sandimmun by slow infusion. Sandimmun will be diluted with normal saline or 5% glucose before use.

Your doctor will adjust your dose to one that is ideal for you. To do this you will need to have regular blood tests and visits to the hospital, especially after the transplant. This

will give you a chance to talk to your doctor about your treatment and mention any problems that you may have.

Sandimmun is different from Neoral®

There is another medicine called Neoral which also contains cyclosporin. However, there are important differences between these two medicines.

Because of these differences, you must not change from Sandimmun to Neoral or from Neoral to Sandimmun unless it is under the strict supervision of your doctor.

Do not take Sandimmun and Neoral at the same time.

How much is given

The dose of Sandimmun is worked out for each person. It will depend on how much you weigh, what condition is being treated, how well Sandimmun works for you, and whether you have any side effects from this medicine.

Your dose may be changed from time to time. Your doctor will work out the correct dose of Sandimmun for you.

How it is given

Concentrate for intravenous infusion:

Sandimmun can be given by intravenous infusion (drip) for a short time after transplant surgery or in other cases where the medicine cannot be taken by mouth. The dose is given in hospital, usually over 2-6 hours.

When to take it

Solid organ transplantation

Patients in whom the use of Sandimmun concentrate for infusion is indicated should receive one-third of the recommended oral cyclosporine dose (See Neoral data sheet) shortly after surgery.

Bone marrow transplantation

The recommended i.v. dose is 3 to 5mg/kg per day, starting on the day before transplantation.

How long it is given

Keep having this medicine for as long as your doctor recommends. The length of treatment will depend on what medical condition you have.

If you forget to take it

If it is almost time for your next dose, skip the dose you missed and take the next dose when you are meant to.

Otherwise, take it as soon as you remember, and then go back to taking it as you would normally.

Do not take a double dose to make up for the one that you missed. This may increase the chance of you getting an unwanted side effect.

If you miss more than one dose, contact your doctor as soon as possible.

If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.

If you think that you have been given too much (Overdose)

Immediately tell your doctor or telephone the National Poisons Information Centre, Dunedin (telephone number 0800 POISON or 0800 764 766), or go to the Accident and Emergency Department at your nearest hospital if you think that you or anyone else may have taken too much Sandimmun. Do this even if there are no signs of discomfort or poisoning.

Keep the telephone numbers for these places handy.

While you are taking Sandimmun

Things you must do

Make sure that you keep all of your doctor's appointments and have any tests done that are ordered by your doctor.

Your doctor may ask you to have tests from time to time to check how well your kidneys and liver are working. It may be necessary to measure the amount of cyclosporin, as well as the levels of other chemicals (eg. potassium) in your blood. Your blood pressure will also be checked regularly.

Avoid eating large amounts of foods that are high in potassium.

In some people having Sandimmun, the amount of potassium in the blood can increase (called hyperkalaemia). The amount of potassium in the blood can also be increased by eating certain foods. Your doctor can tell you which foods to avoid.

If you become pregnant while having Sandimmun, tell your doctor.

Your doctor can discuss with you the risks of having it while you are pregnant

If you develop lumps anywhere on/in your body, or develop any moles, or you notice changes in existing moles, tell your doctor.

This may be an early sign of a cancer. Immunosuppressant medicines, including Sandimmun, may increase the risk of developing certain cancers, including skin cancer and lymphoma (cancer of the lymphatic system).

Limit your exposure to sunlight and UV light. If you go out in the sun, wear a hat, appropriate protective clothing and a sunscreen with a high protection factor.

This will help to prevent the development of skin cancer.

Take special care of your teeth and gums.

Taking good care of your teeth and gums will help to prevent dental and mouth infections.

If you experience any symptoms of infection (e.g. fever, sore throat), inform your doctor immediately. People taking immunosuppressant medicines are at a greater risk of getting infections.

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are having Sandimmun.

Tell any other doctor, dentist or pharmacist who treats you that you are being given Sandimmun.

Things you must not do

Do not have any vaccinations without first checking with your doctor.

Some vaccines may be less effective or they may cause unwanted side effects while you are taking Sandimmun.

Do not give this medicine to anyone else even if their condition seems similar to yours.

Do not take it to treat any other complaints unless your doctor tells you to.

Things to be careful of

Be careful driving, operating machinery or doing jobs that require you to be alert while you are having Sandimmun until you know how it affects you.

This medicine can cause tiredness, lack of energy or confusion in some people.

If you have any of these symptoms, do not drive or do anything else that could be dangerous.

Sandimmun concentrate for solution for infusion contains 34.4 % v/v ethanol (27.8 % w/v). A 100 mg dose of Sandimmun concentrate contains about half a gram of alcohol, equivalent to about 14 mL of beer or 6 mL of wine.

Alcohol may be harmful for those suffering from alcoholism, epilepsy, brain injury or liver disease as well as for pregnant or breast-feeding women and children.

Like other medicines that dampen the immune system, cyclosporin may influence your body's ability to fight against infection and may cause tumours or other malignancies, particularly of the skin.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are having Sandimmun, even if you do not think it is connected with the medicine.

All medicines can have side effects. Sometimes they are serious, but most of the time they are not. You may need medical treatment if you get some of the side effects. Your doctor may be able to relieve some of the side effects of Sandimmun by lowering the dose.

If you are over 65 years old, you should be especially careful while having this medicine. Report any side effects promptly to your doctor.

As people grow older, they are more likely to get side effects from medicines.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor if you notice any of the following and they worry you:

- tiredness, lack of energy
- excessive growth of body and facial hair
- overgrown, thickened, swollen or bleeding gums
- stomach upset, including nausea (feeling sick), vomiting, loss of appetite, diarrhoea, stomach cramps, stomach ulcer
- tremor (shaking)
- headache, including migraine
- sensitivity to light
- weight loss or gain
- feeling depressed (sad)

- flushing of face, acne, darkening of skin
- painful menstrual periods or lack of periods
- increase in size of breasts
- muscle cramps, tenderness or weakness
- blocked or stuffy nose
- pain of lower extremities

The above side effects are not usually serious.

Tell your doctor immediately if you notice any of the following:

- fever (temperature above 37°C)
- constant "flu-like" symptoms such as chills, sore throat, aching joints, swollen glands, or any other signs of infection
- unusual bleeding or bruising
- signs of allergy such as rash, itching or hives on the skin; swelling of the face, lips, tongue or other part of the body; shortness of breath, wheezing or troubled breathing
- new lumps or moles, or changes to existing moles, anywhere on the body
- swelling of the eyelids, hands or feet due to excess fluid
- a change in the amount of urine passed or in the number of times you urinate, pain on urinating, dark, bloody or smelly urine
- yellowing of the skin and/or eyes (jaundice) often accompanied by generally feeling unwell (for example, tiredness, lack of energy, loss of appetite, nausea and vomiting, pain in the abdomen)
- severe pain or tenderness in the stomach or abdomen
- vomiting blood or material that looks like coffee grounds; black sticky bowel motions or bloody diarrhoea
- unusual tiredness or weakness, which may be accompanied by dizziness, spots before the eyes, shortness of breath and pale skin
- numbness or "pins and needles" in the hands and feet
- disturbance in brain function which may cause a variety of symptoms, including confusion, personality changes, disorientation, agitation, inability to sleep, decreased responsiveness, weakness and

loss of coordination in arms and legs with or without abnormal speech or eye movements, seizures (fits), clumsiness, memory loss, difficulty speaking or understanding what others say, visual hallucinations (seeing things that are not there) or other problems with vision, coma, paralysis of part or all of the body, stiff neck

- buzzing or ringing in the ears, difficulty hearing

The above are serious side effects that need medical attention.

Some side effects may not give you any symptoms and can only be found when tests are done. Some of these side effects include:

- changes in kidney or liver function
- raised blood pressure
- increased potassium or cholesterol in the blood
- decreased magnesium in the blood
- an increase in the amount of uric acid in the blood, which can lead to gout
- an increase in blood sugar
- low white blood cell count
- low levels of red blood cells
- low levels of platelets in the blood

Your doctor will make sure that tests are done regularly to watch for these side effects.

Tell your doctor if you notice anything else that is making you feel unwell.

If you notice any other side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Other side effects not listed here may happen in some people.

After using Sandimmun

Storage

Sandimmun is usually stored in the doctor's surgery or clinic, or at the pharmacy. However, if you need to store Sandimmun:

Store the concentrate for i.v. infusion below 30°C.

Keep the medicine where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Keep Sandimmun in the original pack until it is time for it to be given.

Once an ampoule is opened, the contents should be used immediately.

Diluted infusion solution must be discarded after 24 hours.

Disposal

If your doctor tells you to stop having Sandimmun or it has passed its expiry date, ask your pharmacist what to do with any medicine you have left over.

Product description

What it looks like

Sandimmun concentrate for intravenous infusion is a clear, brown-yellow oily liquid in 5 mL glass ampoules; packs of 10.

Ingredients

Sandimmun concentrate for intravenous infusion contains:

- 50 mg cyclosporin per mL of infusion solution.

It also contains:

- polyoxyethylated castor oil (Cremophor® EL)
- ethanol (alcohol)
- macroglycerol ricinoleate

Sponsor

Sandimmun is supplied in New Zealand by:

Novartis New Zealand Limited
PO Box 99102
Newmarket
Auckland 1149
New Zealand

Telephone 0800 354 335

® = Registered Trademark

This leaflet was prepared in August 2022 based on the currently approved data sheet for this product.

Internal use only
Sim220822cNZ based on DS
sim220822iNZ