

Oxycodone Sandoz[®]

Oxycodone hydrochloride modified release tablet

Consumer Medicine Information (CMI)

What is in this leaflet

This leaflet answers some common questions about OXYCODONE SANDOZ tablets.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking this medicine against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What OXYCODONE SANDOZ tablets are taken for

OXYCODONE SANDOZ tablets contain oxycodone hydrochloride. Oxycodone belongs to a group of medicines called opioid analgesics.

OXYCODONE SANDOZ tablets are used to relieve moderate to severe persistent pain when other forms of treatment have failed or are otherwise inappropriate to provide sufficient management of pain.

Your doctor, however, may prescribe it for another purpose.

Ask your doctor if you have any questions about why it has been prescribed for you.

As with all strong painkillers, your body may become used to you taking OXYCODONE SANDOZ tablets. Taking it may result in physical dependence. Physical dependence means that you may experience withdrawal symptoms if you stop taking oxycodone suddenly, so it is

important to take it exactly as directed by your doctor.

This medicine is only available with a doctor's prescription.

Before you take it

Long-term use of Oxycodone Sandoz tablets may result in a decrease in sex hormone levels which may affect sperm production in men and the menstrual cycles in females. Talk to your doctor if you have concerns.

When you must not take it

Do not take OXYCODONE SANDOZ tablets if you:

- have any breathing problems such as acute asthma, respiratory depression (breathing slows or weakens) or other obstructive airways disease
- are severely drowsy or have a reduced level of consciousness
- suffer from irregular heartbeats or changes in the way the heart beats
- have heart disease due to long term lung disease
- have just consumed a large amount of alcohol, regularly consume large amounts of alcohol or have confusion and shaking due to alcohol withdrawal
- suffer from convulsions, fits or seizures
- have a head injury, a brain tumour or have raised pressure within the head, brain or spinal cord
- have sudden, severe abdominal pain
- have a condition where your stomach empties more slowly than it should, or have any condition that obstructs the bowel/stomach or affects bowel transit (movement of food or ingested material along the bowel)

- have swallowing difficulties or narrowing of the oesophagus
- have severe liver or kidney disease
- are about to have an operation (including surgery on your spine for pain relief in the next 24 hours) or have had an operation within the last 24 hours
- take medicine for depression called a 'monoamine oxidase inhibitor' or have taken any in the last two weeks.

Do not take OXYCODONE SANDOZ tablets if you are allergic to oxycodone, opioid painkillers, or any of the ingredients listed at the end of this leaflet.

This medicine contains lactose so consider this if you are lactose intolerant.

Do not take this medicine after the expiry date (EXP) printed on the pack.

If you take it after the expiry date has passed, it may not work very well.

Do not take it if the packaging is torn or shows signs of tampering.

Do not give this medicine to a child younger than 12 years of age.

Safety and effectiveness in children younger than 12 years of age have not been established.

Before you start to take it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have had any medical conditions, especially the following:

- sleep apnoea (temporarily stopping breathing while you sleep)
- low blood pressure

- increased prostate size or difficulty passing urine
- chronic lung, liver or kidney disease
- disease of your gall bladder or bile duct
- inflammation of the pancreas
- underactive adrenal glands
- underactive thyroid gland
- lactose intolerance
- inflammatory bowel disease or recent abdominal surgery
- diverticulitis (inflammation of bowel wall)
- oesophageal, stomach or intestinal disorders (including cancer in these areas), resulting in narrowing of the oesophagus, stomach or intestines
- severe mental condition involving losing contact with reality, hearing voices or an inability to think clearly
- an addiction or history of abuse of alcohol, opioids or other drugs

Pregnancy and breastfeeding

This medicine is not recommended to be taken during labour.

Oxycodone given to the mother during labour can cause breathing problems and signs of withdrawal in the newborn.

Tell your doctor if you are breast feeding or planning to breast-feed.

Oxycodone can pass into the breast milk and can affect the baby. Your doctor can discuss with you the risks involved.

If you have not told your doctor about any of the above, tell them before you take any OXYCODONE SANDOZ tablets.

Do not take this medicine if you are pregnant or intend to become pregnant whilst taking this medicine.

Like most medicines of this kind, OXYCODONE SANDOZ tablets are not recommended to be taken during pregnancy. Your doctor will discuss the risks of taking it if you are pregnant.

Addiction

You can become addicted to OXYCODONE SANDOZ even if you take it exactly as prescribed. OXYCODONE SANDOZ may become habit forming causing mental and physical dependence. If abused it may become less able to reduce pain.

Dependence

As with all other opioid containing products, your body may become used to you taking OXYCODONE SANDOZ. Taking it may result in physical dependence. Physical dependence means that you may experience withdrawal symptoms if you stop taking OXYCODONE SANDOZ suddenly, so it is important to take it exactly as directed by your doctor.

Tolerance

Tolerance to OXYCODONE SANDOZ may develop, which means that the effect of the medicine may decrease. If this happens, more may be needed to maintain the same effect.

Withdrawal

Continue taking your medicine for as long as your doctor tells you. If you stop having this medicine suddenly, your pain may worsen and you may experience some or all of the following withdrawal symptoms:

- nervousness, restlessness,
- agitations, trouble sleeping or anxiety
- body aches, weakness or stomach cramps
- loss of appetite, nausea, vomiting or diarrhoea
- increased heart rate, breathing rate or pupil size
- watery eyes, runny nose, chills or yawning
- increased sweating.

Taking other medicines

Tell your doctor if you are taking any other medicines or dietary supplements, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and OXYCODONE SANDOZ tablets may interfere with each other. These include:

- medicines to treat depression, psychiatric or mental disorders. Medicines to treat depression belonging to a group called monoamine oxidase inhibitors must be stopped 14 days before OXYCODONE SANDOZ tablets are taken.
- antidepressants eg; fluoxetine, paroxetine
- medicines to help you sleep
- medicines to put you to sleep during an operation or procedure
- medicines to relax your muscles
- medicines to lower blood pressure
- quinidine and other medicines to treat the heart
- medicines to treat convulsions e.g. phenytoin, carbamazepine
- medicines to thin the blood e.g. coumarin derivatives such as warfarin
- cimetidine, a medicine to treat stomach ulcers or heartburn
- medicines to relieve stomach cramps or spasms, to prevent travel sickness or to treat Parkinson's disease
- medicines to stop nausea or vomiting e.g. metoclopramide
- medicines to treat Parkinson's disease
- medicines to treat urinary incontinence
- other pain relievers including other opioids
- antifungals e.g. ketoconazole
- antibiotics e.g. clarithromycin, rifampicin
- medicines to treat HIV infection and AIDS e.g. ritonavir
- St John's wort (a herbal preparation)
- grapefruit and grapefruit juice
- medicines to treat epilepsy, pain and anxiety eg; gabapentin and pregablin

- alcohol

These medicines, dietary supplements or alcohol may be affected by OXYCODONE SANDOZ tablets, may affect how well OXYCODONE SANDOZ tablets work or may increase side effects. You may need to use different amounts of your medicines, or take different medicines

Your doctor or pharmacist has more information on medicines to be careful with or avoid while taking this medicine

How to take OXYCODONE SANDOZ tablets

How much to take

Your doctor will tell you exactly how much to take.

Follow the instructions given to you by your doctor and pharmacist exactly.

How to take it

Swallow OXYCODONE SANDOZ tablets whole with a full glass of water or other fluid.

Do not break, chew, crush or dissolve the tablets.

OXYCODONE SANDOZ tablets are only designed to work properly if swallowed whole. The tablets may release all their contents at once if broken, chewed, crushed or dissolved which can be dangerous and cause serious problems, such as an overdose, which may be fatal.

If you have trouble swallowing your tablets whole, talk to your doctor.

You must only take OXYCODONE SANDOZ tablets by mouth.

Taking this medicine in a manner other than that prescribed by your doctor can be harmful to your health.

There are no data on rectal administration of OXYCODONE SANDOZ tablets, therefore rectal administration of OXYCODONE SANDOZ tablets is not recommended.

When to take it

Take OXYCODONE SANDOZ tablets every 12 hours.

Take OXYCODONE SANDOZ tablets regularly to control the pain.

Taking them at the same time each day will assist in ensuring the best effect in improving your pain. If however, you begin to experience worsening pain and you are taking your OXYCODONE SANDOZ tablets as prescribed, contact your doctor as your dosage may have to be reviewed.

How long to take it

Continue taking your medicine for as long as your doctor tells you.

If you stop taking this medicine suddenly, your pain may worsen and you may experience withdrawal symptoms such as:

- body aches
- loss of appetite, nausea, stomach cramps or diarrhoea
- fast heart rate
- sneezing or runny nose
- chills, tremors, shivering or fever
- trouble sleeping
- increased sweating and yawning
- weakness
- nervousness or restlessness.

If you forget to take it

If you forget to take your tablets, contact your doctor or pharmacist for advice.

Do not take a double dose to make up for the dose you have missed.

This will increase the chance of you getting unwanted side effects.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering when to take your medicine, ask your pharmacist for hints.

For example, take your medicine at the same time each morning and evening such as 8 a.m. and 8 p.m.

If you take too much (overdose)

Immediately telephone your doctor, or the Poisons Information Centre (telephone 0800 POISON or 0800 764 766) for advice, or go to Accident and Emergency at your nearest hospital, if you think you or

anyone else may have taken too many OXYCODONE SANDOZ tablets.

Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

- **If you or someone else receives too much (overdose), and experience one or more of the symptoms below, call 111 for an ambulance. Keep the person awake by talking to them or gently shaking them every now and then. You should follow the steps even if someone other than you have accidentally used OXYCODONE SANDOZ that was prescribed for you. If someone takes and overdose, they may experience or more the following symptoms:**

- slow, unusual or difficult breathing
- drowsiness, dizziness or unconsciousness
- slow or weak heartbeat
- nausea or vomiting
- convulsions or fits

While you are taking it

Things you must do

Take OXYCODONE SANDOZ tablets exactly as your doctor has prescribed.

Before you start on a new medicine, remind your doctor and pharmacist that you are taking OXYCODONE SANDOZ tablets.

Tell any other doctors, dentists and pharmacists who treat you that you are taking this medicine.

If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine.

It may affect other medicines used during surgery.

If you become pregnant while taking this medicine, tell your doctor immediately.

Keep all of your doctor's appointments so that your progress can be checked.

Tell your doctor if your pain is getting worse. Also tell your doctor if you are having any problems or difficulties while you are being treated with OXYCODONE SANDOZ tablets.

Tolerance to oxycodone may develop which means that the effect of the medicine may decrease. If this happens, your doctor may review your dose so that you get adequate pain relief.

Keep enough OXYCODONE SANDOZ tablets with you to last over weekends and holidays.

Things you must not do

Do not drink alcohol while you are taking OXYCODONE SANDOZ tablets.

Drinking alcohol whilst taking OXYCODONE SANDOZ tablets may make you feel more sleepy and increase risk of serious side effects, such as shallow breathing with the risk of stopping breathing and loss of consciousness.

Do not take OXYCODONE SANDOZ tablets to treat any other complaint unless your doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not stop taking your medicine, exceed the dose recommended or change the dosage without checking with your doctor.

Over time your body may become used to you taking oxycodone so if you stop taking it suddenly, your pain may worsen and you may have unwanted side effects such as withdrawal symptoms. This is called physical dependence.

If you need to stop taking this medicine, your doctor will gradually reduce the amount you take each day, if possible, before stopping the medicine completely.

Things to be careful of

Do not drive or operate machinery until you know how OXYCODONE SANDOZ tablets affect you.

OXYCODONE SANDOZ tablets may cause drowsiness, dizziness, hallucinations, disorientation, blurred vision or other vision problems or may affect alertness. If you are affected, you should not drive or operate machinery. Discuss these effects with your doctor.

Be careful if you are elderly, unwell or taking other medicines.

Some people may experience side effects such as drowsiness, confusion, dizziness and unsteadiness, which may increase the risk of a fall.

If you feel light-headed, dizzy or faint when getting out of bed or standing up, get up slowly.

Standing up slowly will help your body get used to the change in position and blood pressure. If this problem continues or gets worse, talk to your doctor.

Tell your doctor if you suffer from nausea or vomiting when taking OXYCODONE SANDOZ tablets.

If you vomit after your dose, your pain may come back, as you may not have absorbed your medicine. If this happens, speak to your doctor. Your doctor may prescribe some medicine to help you stop vomiting.

Tell your doctor if taking OXYCODONE SANDOZ tablets causes constipation.

Your doctor can advise you about your diet, the proper use of laxatives and suitable exercise you can do to help you manage this.

There is potential for abuse of oxycodone and the development of addiction to oxycodone. It is important that you discuss this issue with your doctor.

Side effects

All medicines may have some unwanted side effects. Sometimes they are serious but most of the time they are not. As for many other medicines of this type, that is opioid analgesics, many side effects tend to reduce over time, with the exception of constipation. This means that the

longer you take this medicine, the less it may cause problems for you. Your doctor has weighed the risks of this medicine against the benefits they expect it will have for you.

Do not be alarmed by this list of possible side effects.

Not everybody experiences them.

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking OXYCODONE SANDOZ tablets.

This medicine helps most people with moderate to severe pain, but it may have unwanted side effects in a few people. Other side effects not listed here may also occur in some people.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor if you notice any of the following and they worry you:

- mild abdominal problems such as diarrhoea, feeling sick (nausea), decreased appetite or constipation
- dry mouth, hiccups, sore throat, trouble swallowing or changes in voice
- excessive sweating
- feeling anxious or nervous, trouble sleeping or abnormal dreams
- trouble with your balance
- new problems with your eyesight
- skin rash, itching, chills or fever
- muscle problems such as spasms, tremors or twitching
- swelling of legs or ankles
- absence of menstrual periods
- impotence
- decreased sexual drive

Tell your doctor as soon as possible if you notice any of the following and they worry you:

- stomach discomfort, vomiting, indigestion or abdominal pain
- choking, gagging, regurgitation, tablets stuck in throat or trouble swallowing the tablets
- abnormal thinking, changes in mood or feeling deep sadness

- drowsiness, feeling faint or fainting or dizziness especially when standing up
- slow or noticeable heartbeats
- headache or confusion
- unusual weakness, loss of strength or trouble walking
- fatigue, feeling of tiredness, drowsiness or lack of energy
- changes in passing urine such as the volume passed, pain or feeling the need to urinate urgently.

The above list includes serious side effects that may require medical treatment.

If any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:

- your breathing slows or weakens
- you have an allergic reaction: shortness of breath, wheezing, shallow or difficult breathing; swelling of the face, lips, tongue, throat or other parts of the body; rash, itching or hives on the skin
- seizures, fits or convulsions
- fast or irregular heartbeats
- chest pain or chest tightness.

The above list includes very serious side effects. You may need urgent medical attention or hospitalization.

Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

Other side effects not listed above may also occur in some people.

When seeking medical attention, take this leaflet and any remaining tablets with you to show the doctor.

After taking it

Storage

Keep your tablets in the blister pack or bottle until it is time to take them.

If you take the tablets out of the blister pack they may not keep as well.

Keep your tablets in a cool, dry place, where the temperature stays below 25°C.

Do not store it or any other medicine in the bathroom, near a sink or on a window sill.

Do not leave it in the car on hot days.

Heat and damp can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking the tablets or the tablets have passed their expiry date, ask your pharmacist how to dispose of medicines no longer required.

Product description

What it looks like

OXYCODONE SANDOZ® tablets are available in five strengths which are as follows (not all strengths are currently marketed in New Zealand):

5 mg - round, blue, biconvex

10 mg - round, white, biconvex

20 mg - round, pink, biconvex

40 mg - round, yellow, biconvex

80 mg - round, green, biconvex

OXYCODONE SANDOZ® tablets come in bottles or boxes containing blister packs of 20 tablets.

Ingredients

Active ingredients:

5 mg tablets contain 5 mg oxycodone hydrochloride

10 mg tablets contain 10 mg oxycodone hydrochloride

20 mg tablets contain 20 mg oxycodone hydrochloride

40 mg tablets contain 40 mg oxycodone hydrochloride

80 mg tablets contain 80 mg oxycodone hydrochloride.

Inactive ingredients:

- maize starch
- behenoyl polyoxylglycerides
- lactose monohydrate
- medium chain triglycerides
- copovidone
- hydrogenated castor oil
- colloidal anhydrous silica
- magnesium stearate

In addition, the tablet coating also contains:

- hypromellose
- microcrystalline cellulose
- stearic acid
- titanium dioxide
- indigo carmine aluminium lake (5mg tablets only)
- iron oxide red (20mg tablets only)
- iron oxide yellow (40mg tablets only)
- iron oxide black (80mg tablets only).
- 815063 Spectracol Green Lake (80mg tablets only).

This medicine does not contain sucrose, gluten or tartrazine.

Manufacturer / Sponsor

Novartis New Zealand Limited,
PO Box 99102, Newmarket,
Auckland 1149

Tel: 0800 354 335

This leaflet was revised in September 2021.

®: OXYCODONE SANDOZ is a registered trademark