

Vimcosa tablets

Contains the active ingredient lacosamide (pronounced “la-KOE-sa-mide”)

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Vimcosa.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Vimcosa against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What Vimcosa is used for

Vimcosa film-coated tablets are used in patients over 4 years in combination with other medicines to control epilepsy.

Vimcosa film-coated tablets can only be used by itself in patients over 16 years. Epilepsy is a condition where you have repeated seizures. There are many different types of seizures, ranging from mild to severe.

This medicine belongs to a group of medicines called antiepileptics.

These medicines are thought to work by controlling brain chemicals which send signals to nerves so that seizures do not happen.

Your doctor may have

prescribed this medicine for another reason.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

There is no evidence that Vimcosa is addictive.

This medicine is available only with a doctor's prescription.

Vimcosa is not recommended for use in children under the age of 4 years as its safety and effectiveness has not been established in this age group.

Before treatment with Vimcosa

When you must not take it

Do not take Vimcosa if you have an allergy to:

- Lacosamide or any of the ingredients listed at the end of this leaflet

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin

Do not take Vimcosa if you have, or have had, a heart condition causing an uneven heart beat.

If you are not sure whether any of the above conditions apply to you, ask your doctor.

Do not take this medicine after the expiry date printed on the pack.

Do not take this medicine if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your doctor or pharmacist.

Before you start to take it

Tell your doctor or pharmacist if you:

- are taking any other medicines, especially barbiturates (such as phenobarbitone) or any other antiepileptic medicines (such as carbamazepine, lamotrigine or levetiracetam)
- have allergies to any other substances, such as foods, preservatives or dyes.

Tell your doctor if you have, or have had, any medical conditions, especially the following:

- heart problems
- kidney problems
- liver problems
- any mental health condition, such as depression.

Tell your doctor if you are pregnant or intend to become pregnant.

Vimcosa may affect your developing baby if you take it during pregnancy. However, it is very important to control your seizures while you are pregnant. Your doctor will outline and weigh up all the risks and benefits of taking Vimcosa

during pregnancy to help decide whether or not you should take it.

Tell your doctor if you are breastfeeding or plan to breastfeed.

Your doctor will discuss the risks and benefits of using Vimcosa if you are breastfeeding.

If you have not told your doctor or pharmacist about any of the above, tell them before you start taking Vimcosa.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Vimcosa may interfere with each other.

These include:

- medicines to treat heart problems
- medicine which may have an effect on your heart beat such as carbamazepine, lamotrigine or pregabalin.

Vimcosa does not interact with the oral contraceptive pill.

However, you may be given Vimcosa together with other antiepileptic medicines that do interact and may affect the effectiveness of your contraceptive. Your doctor may advise you to use an additional method of contraception if you take Vimcosa with other antiepileptic medicines.

How to take Vimcosa

How much to take

Follow all directions given to you by your doctor

carefully.

They may differ from the information contained in this leaflet

Your doctor will tell you how much Vimcosa you will need to take each day. This may depend on your condition, your body weight, and whether or not you are taking any other medicines.

Your doctor may start you on a low dose of Vimcosa first of 50 mg or 100 mg twice a day. Your doctor may slowly increase your dose up to a maximum of 300 mg twice a day, until you are taking enough to control your epilepsy and you are not having any seizures

If you do not understand the instructions on the box, ask your doctor or pharmacist for help.

How to take it

Swallow the tablets whole with a full glass of water.

Do not crush or chew the tablets.

When to take it

Take Vimcosa twice a day, once in the morning and once at night. Take it at about the same time each day.

Taking your medicine at the same time each day will have the best effect. It will also help you remember when to take it.

It does not matter if you take this medicine before or after food.

If you forget to take it

Contact your doctor if you have missed one or more doses.

Do not take a double dose

to make up for the dose that you missed.

How long to take it

Most antiepileptic medicines take time to work, so do not be discouraged if you do not feel better straight away.

Continue taking your medicine for as long as your doctor tells you to.

This medicine helps control your condition, but does not cure it. Therefore, you must take your medicine every day, even if you feel well.

Do not stop taking Vimcosa, or change the dosage, without checking with your doctor. Do not let yourself run out of medicine over the weekend or on holidays. Stopping Vimcosa suddenly may cause unwanted side effects or make your condition worse. Your doctor will slowly reduce your dose before you can stop taking it completely.

If you take too much (overdose)

Immediately telephone your doctor or the Poisons Information Centre (telephone in Australia 13 11 26, in New Zealand 0800 POISON or 0800 764 766), or go to Accident and Emergency at your nearest hospital if you think that you or anyone else may have taken too much Vimcosa. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Symptoms of overdose may include feeling dizzy, drowsy or having an upset stomach.

While being treated with Vimcosa

Things you must do

Tell your doctor immediately if you notice an

increase in seizures.

Tell your doctor immediately if you have symptoms of depression or thoughts of harming yourself.

Tell any other doctors, dentists, and pharmacists who are treating you that you are taking this medicine.

If you are about to be started on any new medicine, tell your doctor, dentist or pharmacist that you are taking Vimcosa.

Before you have any surgery or emergency treatment, tell your doctor or dentist that you are taking Vimcosa.

Tell your doctor if you feel Vimcosa is not helping your condition. Your doctor may need to change your medicine.

Tell your doctor if, for any reason, you have not taken your medicine exactly as prescribed. Otherwise, your doctor may think it is not working effectively and change your treatment unnecessarily.

If you become pregnant while taking this medicine, tell your doctor.

Be sure to keep all of your doctor's appointments so that your progress can be checked.

Your doctor will check your progress and may want to take some tests from time to time. This helps to prevent unwanted side effects.

Things you must not do

Do not give Vimcosa to anyone else, even if their symptoms seem similar to yours or they have the same condition as you.

Do not take Vimcosa to treat any other complaints unless

your doctor tells you to.

Do not stop taking Vimcosa or change the dosage unless your doctor tells you to. Stopping Vimcosa suddenly may cause unwanted side effects or make your condition worse.

Things to be careful of

Be careful before you drive or use any machines or tools until you know how Vimcosa affects you.

As with other antiepileptic medicines, Vimcosa may cause dizziness or drowsiness. This is more frequent at the beginning of treatment or after an increase in the dose.

If you are feeling dizzy or drowsy, do not drive a car, operate machinery, or do anything else that could be dangerous.

As a safety precaution, do not take Vimcosa with alcohol

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Vimcosa.

This medicine helps most people with epilepsy but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Do not be alarmed by this list of side effects.

You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

If you get any side effects, do not stop taking Vimcosa without first talking to your

doctor or pharmacist.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- dizziness or problems with balance or coordination
- headache
- nausea (feeling sick) or vomiting
- feeling tired, drowsy or sleepy
- forgetfulness
- tremors
- itching.

The above list includes the more common side effects of your medicine. They are mostly mild and short-lived.

Tell your doctor as soon as possible if you notice any of the following:

- feelings of depression
- feeling aggressive or agitated
- spinning sensations
- double vision or blurred vision
- having trouble sleeping.

The above list includes serious side effects that may require medical attention.

If any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:

- thoughts of harming yourself
- more frequent or more severe seizures
- fainting or feeling lightheaded
- heart palpitations or a rapid or irregular pulse
- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin.

The above list includes more serious side effects. You may need urgent medical attention or hospitalisation.

Tell your doctor if you notice anything else that is making you feel unwell.

Other side effects not listed above may happen in some people.

After using Vimcosa

Storage

Tablets

Keep your tablets in the pack until it is time to take them.

If you take the tablets out of the pack they will not keep well.

Keep your tablets in a cool dry place where the temperature stays below 30°C

Do not store Vimcosa or any other medicine in the bathroom or near a sink.

Do not leave your medicine on a window sill or in the car on hot days.

Heat and dampness can destroy some medicines.

Keep Vimcosa where children cannot reach it.

A locked cupboard at least one- and-a-half metres above the ground is a good place to store medicines.

Keep Vimcosa where young children cannot reach it.

A locked cupboard at least one and- a-half meters above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Product description

What it looks like

Tablets

Vimcosa film-coated tablets are available in four strengths:

- 50 mg – Pink, oval

shaped, film-coated tablets debossed with "LAC" on one side and "50" on other side.

(Blister pack: Aust R 353576)

(Bottle pack: Aust R 353580)

- 100 mg – Dark yellow, oval shaped, film-coated tablets debossed with "LAC" on one side and "100" on other side.
(Blister pack: Aust R 353583)
(Bottle pack: Aust R 353597)
- 150 mg – Peach, oval shaped, film-coated tablets debossed with "LAC" on one side and "150" on other side.
(Blister pack: Aust R 353572)
(Bottle pack: Aust R 353588)
- 200 mg – Blue, oval shaped, film-coated tablets debossed with "LAC" on one side and "200" on other side.
(Blister pack: Aust R 353578)
(Bottle pack: Aust R 353586)

What Vimcosa Tablets are packed in:

The film-coated tablets are packed in

- PVC-Aclar Blister Carton: 14's, 56's, 168's Tablets
- HDPE Bottle: 56's Tablets

Ingredients

Active ingredient:

Each Vimcosa film-coated tablet contains either 50 mg, 100 mg, 150 mg or 200 mg of lacosamide as the active ingredient.

Inactive ingredients:

- Microcrystalline cellulose
- Crospovidone
- Hyprolose
- Colloidal anhydrous silica

- Magnesium stearate
- Opadry II 85F20249 Purple (50 mg)
- Opadry 85F520062 Yellow (100 mg)
- Opadry II 85F27043 TAN (150 mg)
- Opadry II 85F505244 Blue (200 mg)

Do not take this medicine if you are allergic to any of these ingredients.

Australian Sponsor:

Pharmacor Pty Ltd.
Suite 803, Level 8,
Tower A, The Zenith,
821 Pacific Highway,
Chatswood, NSW 2067,
Australia
Tel: 1300 138 805
www.pharmacor.com.au

NZ Sponsor:

Pharmacor Limited.
c/- Wynn Williams
Level 25, Vero Centre 28
Shortland Street
Auckland Central
AUCKLAND, 1010 New
Zealand.
Tel: +64 800 172 553 or
0800 172 553

This leaflet was prepared in Sep 2026.