

# Flucloxacillin Eugia Injection

**Flucloxacillin sodium monohydrate**

500 mg & 1 g, powder for injection

---

## What is in this leaflet

---

Please read this leaflet carefully before you start using Flucloxacillin Eugia.

This leaflet answers some common questions about Flucloxacillin Eugia. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you using Flucloxacillin Eugia against the benefits they expect it will have for you.

**If you have any concerns about using this medicine, ask your doctor or pharmacist.**

Keep this leaflet with the medicine. You may need to read it again.

---

## What Flucloxacillin Eugia is used for

---

Flucloxacillin Eugia is an antibiotic that belongs to a group of medicines called penicillins. These antibiotics work by killing the bacteria that are causing your infection.

Flucloxacillin Eugia is an antibiotic used to treat some infections in different parts of the body caused by bacteria. It is used to:

- treat skin and soft tissue infections caused by bacteria which are resistant to penicillin.  
e.g. infections of the joints, respiratory tract, urinary tract, inflammation (pain, heat, swelling or redness) of the middle ear, inflammation of the endocardium (a membrane that covers the inside surface of the heart), infected blood and meningitis.
- prevent infections resulting from surgery, in particular chest, bone and joint procedures.

Flucloxacillin Eugia is also used in the prophylaxis of staphylococcal infections during major surgical procedures, particularly in cardiothoracic or orthopaedic surgery.

Flucloxacillin Eugia will not work against infections caused by viruses, such as colds.

Your doctor may have prescribed Flucloxacillin Eugia for another reason.

Ask your doctor if you have any questions about why Flucloxacillin Eugia has been prescribed for you.

This medicine is available only with a doctor's prescription.

There is no evidence that Flucloxacillin Eugia is addictive.

---

## **Before you are given Flucloxacillin Eugia**

---

### ***When you must not be given Flucloxacillin Eugia***

Do not use Flucloxacillin Eugia if:

- your skin and/or eyes have turned yellow (Jaundice) while, or soon after receiving flucloxacillin (Flucloxacillin Eugia).
- you are allergic to flucloxacillin, other penicillins or any of the ingredients listed at the end of this leaflet.  
Some of the symptoms of an allergic reaction may include skin rash, itching, hayfever, swelling and difficult breathing.
- you have had an allergic reaction to cephalosporins.
- you may have an increased chance of being allergic to Flucloxacillin Eugia if you are allergic to cephalosporins.

If you are not sure whether you should start using Flucloxacillin Eugia, talk to your doctor.

### ***Before you are given Flucloxacillin Eugia***

Flucloxacillin (Flucloxacillin Eugia) can cause jaundice (yellowing of skin and/or eyes) and severe liver disease which may last for a long time. This reaction is more frequent in older patients (particularly those over 55 years of age) and those who receive the drug for more than 14 days.

Tell your doctor if:

- you have an allergy to Flucloxacillin Eugia or other penicillins.
- you have had any type of allergic reaction to cephalosporin medicines.  
You may have an increased chance of being allergic to Flucloxacillin Eugia if you are allergic to cephalosporins.
- you have any allergies to any other medicines or any other substances, such as foods, preservatives or dyes. This may include medicines that you buy without a prescription from your pharmacy, supermarket or health food shop.
- your skin and/or eyes have turned yellow (Jaundice) while, or soon after, receiving any antibiotics.

- you have or have ever had any other health problems/medical conditions, including:
  - kidney or liver disease
  - asthma, hayfever or hives
  - glandular fever (infectious mononucleosis).
- you are pregnant or intend to become pregnant.  
Your doctor will discuss the risks and benefits of using Flucloxacillin Eugia during pregnancy.
- you are breast-feeding or plan to breast-feed.  
Your doctor will discuss the risks and benefits of using Flucloxacillin Eugia when breast-feeding.

If you have not told your doctor about any of the above, tell them before you start using Flucloxacillin Eugia.

**Special care should be taken in new-born or premature babies to avoid overdose or high levels of bilirubin, a chemical in the blood which can lead to brain damage.**

### ***Taking other medicines***

Tell your doctor if you are taking any other medicines, including medicines that you buy without a prescription from your pharmacy, supermarket or health food shop. You should also tell any health professional who is prescribing a new medication for you that you are taking Flucloxacillin Eugia.

Some medicines may interfere with Flucloxacillin Eugia. These include:

- probenecid - a medicine used to treat gout.
- antibiotics such as chloramphenicol, erythromycin, sulphonamides or tetracyclines.
- estrogen containing oral contraceptives - if you are using this type of medicine you should take additional precautions to prevent pregnancy.
- Aminoglycosides - If Flucloxacillin Eugia and an aminoglycoside are to be used simultaneously, the two antibiotics should not be mixed together.
- Methotrexate - medicine taken to treat cancer or autoimmune disorders.
- Warfarin - medicine to prevent blood clotting.
- Voriconazole and Posaconazole - medicine used to treat fungal infections.
- Paracetamol - caution is advised when Flucloxacillin Eugia is administered concomitantly with paracetamol due to the increased risk of high anion gap metabolic acidosis.

These medicines may be affected by Flucloxacillin Eugia, or may affect how well it works. You may need different amounts of your medicine, or you may need to take different medicines.

Your doctor or pharmacist has more information on medicines to be careful with or avoid while taking Flucloxacillin Eugia.

Flucloxacillin Eugia should not be mixed with blood products or other fluids containing protein.

---

## How Flucloxacillin Eugia is given

---

Flucloxacillin Eugia may be given in two ways:

- as a slow injection into a vein
- as a deep injection into a large muscle, a joint or the sac surrounding the lung.

Flucloxacillin Eugia must only be given by a doctor or nurse.

Your doctor will decide what the dose is and for how long you will receive Flucloxacillin Eugia. This depends on your infection and other factors, such as your weight. For most infections, Flucloxacillin Eugia is usually given in divided doses throughout the day. Sometimes only a single dose of Flucloxacillin Eugia is required for the treatment and prevention of certain infections.

---

## After you have been given Flucloxacillin Eugia

---

### *Things you must do*

If the symptoms of your infection do not improve within a few days, or if they become worse, tell your doctor.

**If you develop itching with swelling or skin rash or difficulty breathing after you have been given Flucloxacillin Eugia, contact your doctor immediately.**

**If you develop yellow eyes and/or skin (jaundice) while, or soon after receiving Flucloxacillin Eugia, contact your doctor immediately.**

**If you get severe diarrhoea, tell your doctor or pharmacist immediately. Do this even if it occurs up to several weeks after Flucloxacillin Eugia has been stopped.** Diarrhoea may mean that you have a serious condition affecting your bowel. You may need urgent medical care. **Do not take any diarrhoea medicine without first checking with your doctor.**

If you get a sore white mouth or tongue after you have been given Flucloxacillin Eugia, tell your doctor. Also tell your doctor if you get a vaginal itching or discharge. This may mean you have a fungal infection called thrush. Sometimes the use of Flucloxacillin Eugia allows fungi to grow and the above symptoms to occur. Flucloxacillin Eugia does not work against fungi.

If you become pregnant while you are receiving Flucloxacillin Eugia, tell your doctor.

If you have to have any tests tell your doctor you have been given Flucloxacillin Eugia. Flucloxacillin Eugia may affect the results of some tests.

If you are about to start taking any new medicine, tell your doctor and pharmacist that you are receiving Flucloxacillin Eugia.

Tell any doctor, dentist or pharmacist who is treating you that you have been given Flucloxacillin Eugia.

**During prolonged therapy, your doctor should carry out periodic tests to assess kidney and liver function and blood cell formation.**

**Be aware that superinfections may occur.**

This is a new infection resulting from the invasion of bacteria resistant to flucloxacillin. Flucloxacillin Eugia should be discontinued and another therapy commenced.

### ***Things to be careful of***

Be careful before you drive or use any machines or tools until you know how Flucloxacillin Eugia affects you.

Flucloxacillin Eugia generally does not cause any problems with your ability to drive a car or operate machinery. However, as with many other medicines, Flucloxacillin Eugia may cause dizziness, drowsiness or tiredness in some people.

If you experience dizziness, do not drive.

If you drink alcohol, dizziness may be worse.

---

## **In case of overdose**

---

### ***If you take too much (overdose)***

As Flucloxacillin Eugia injection is given to you under the supervision of your doctor, it is very unlikely that you will receive too much.

However, if you experience severe side effects after being given this medicine, tell your doctor immediately.

Immediately telephone your doctor or the National Poisons Centre (telephone 0800 POISON or 0800 764 766) (New Zealand) or Poisons Information Centre on 13 11 26 (Australia), or go to accident and emergency at your nearest hospital, if you think that you or anyone else may have taken too much Flucloxacillin Eugia.

Do this even if there are no signs of discomfort or poisoning.

A very large overdose of Flucloxacillin Eugia can cause brain upsets, including fits.

---

## Side Effects

---

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are using Flucloxacillin Eugia.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the adverse effects.

Ask your doctor or pharmacist to answer any questions you may have.

***Whilst being given Flucloxacillin Eugia***

**Tell your doctor immediately or go to casualty at your nearest hospital if you notice any of the following:**

- a severe rash
- wheezing
- irregular heart beat
- feeling faint
- yellow skin and/or eyes
- a red rash commonly seen on both sides of buttocks, upper inner thighs, armpits, neck.

**Also tell your doctor if you notice:**

- pain or redness at the site of injection
- oral thrush - white, furry, sore tongue and mouth
- vaginal thrush - sore and itchy vagina and/or discharge
- a mild rash

Other side effects not listed above may also occur in some patients. These include very rare conditions such as blood, liver and kidney problems or fits.

***After finishing Flucloxacillin Eugia***

**Tell your doctor if your skin and/or eyes have turned yellow (Jaundice) while or soon after receiving flucloxacillin (Flucloxacillin Eugia).**

**Tell your doctor immediately if you notice any of the following side effects, particularly if they occur several weeks after finishing treatment with Flucloxacillin Eugia:**

- severe abdominal cramps or stomach cramps
- watery and severe diarrhoea, which may also be bloody
- fever, in combination with one or both of the above.

These are rare but serious side effects. Flucloxacillin Eugia can change bacteria (which are normally present in the bowel and normally harmless) to multiply and therefore cause the above symptoms. You may need urgent medical attention.

**Do not take any diarrhoea medicine without first checking with your doctor.**

An illness consisting of a rash, swollen glands, joint pains and fever may occur about a week after treatment.

Other adverse effects not listed above may also occur in some patients. Tell your doctor if you notice any other effects.

Do not be alarmed by this list of possible adverse effects. You may not experience any of them.

---

## **After receiving Flucloxacillin Eugia**

---

### ***Storage***

Flucloxacillin Eugia must be kept out of reach and sight of children.

Flucloxacillin Eugia injection is usually stored in the pharmacy or on the ward. Your doctor, pharmacist or nurse is responsible for storing Flucloxacillin Eugia and disposing of any unused product correctly.

Store below 25°C.

---

## **Product description**

---

### ***What it looks like***

Flucloxacillin Eugia Injection is a white to almost white powder which has to be dissolved in water before it is injected.

Flucloxacillin Eugia is available in packs of 1, 5 and 10 vials.

## ***Ingredients***

### Active ingredient:

- 500 mg or 1 g flucloxacillin sodium monohydrate

### Inactive ingredients:

- None.

---

## **Sponsor Details**

---

Flucloxacillin Eugia is supplied in New Zealand by:

### **CARSL Consulting**

P O Box 766  
Hastings 4156, New Zealand  
Telephone: 06 875 0979  
[www.carsl.co.nz](http://www.carsl.co.nz)  
[carsl@xtra.co.nz](mailto:carsl@xtra.co.nz)

### **Distributed by**

Eugia Pharma (Australia) Pty Ltd  
25 Milton Parade  
MALVERN VIC 3144  
[www.eugiapharma.com.au](http://www.eugiapharma.com.au)  
[eugia.aus@eugiapharma.com](mailto:eugia.aus@eugiapharma.com)

---

## **Date of Preparation**

---

This leaflet was revised in September 2025.