

NEW ZEALAND DATASHEET – QINLOCK® (ripretinib) TABLETS

1 PRODUCT NAME

QINLOCK (ripretinib) tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each QINLOCK tablet contains 50 mg of ripretinib.

Ripretinib is a white to off-white crystalline solid. Ripretinib is a lipophilic, weak base compound, practically insoluble in aqueous media.

Excipients with known effect: sugars as lactose.

For the full list of excipients, see [Section 6.1](#) List of Excipients.

3 PHARMACEUTICAL FORM

QINLOCK tablets are white to off-white oval shaped tablets debossed with 'DC1' on one side.

4 CLINICAL PARTICULARS

4.1 THERAPEUTIC INDICATIONS

QINLOCK is a kinase inhibitor indicated for the treatment of adult patients with advanced gastrointestinal stromal tumours (GIST) who have received prior treatment with 3 or more kinase inhibitors, including imatinib.

4.2 DOSE AND METHOD OF ADMINISTRATION

Dosage

The recommended dosage of QINLOCK is 150 mg (three 50 mg tablets) orally once daily with or without food until disease progression or unacceptable toxicity.

Instruct patients to swallow tablets whole.

Advise patients to take QINLOCK at the same time each day.

Advise patients to take a missed dose if less than 8 hours have passed since the missed scheduled dose. If a patient misses a dose by more than 8 hours of the time it is usually taken, the patient should be instructed not to take the missed dose and simply resume the usual dosing schedule on the following day.

Advise patients not to take an additional dose if vomiting occurs after taking QINLOCK and to continue with their next scheduled dose.

Dose modification guidelines

The recommended dose reduction for adverse reactions is:

- QINLOCK 100 mg orally once daily.

Permanently discontinue QINLOCK in patients who are unable to tolerate 100 mg orally once daily.

The recommended dosage modifications of QINLOCK for adverse reactions are provided in [Table 1](#).

Table 1: Recommended Dose Modifications for QINLOCK

Adverse Reaction	Severity ^a	Dosage Modifications
Palmar-Plantar Erythrodysesthesia Syndrome [PPES] [see Section 4.4 Special Warnings and Precautions for Use]*	Grade 2	<ul style="list-style-type: none"> • Withhold QINLOCK until Grade ≤ 1 or baseline. If recovered within 7 days, resume QINLOCK at same dose; otherwise, resume at reduced dose. • Consider re-escalating QINLOCK if maintained at Grade ≤ 1 or baseline for at least 28 days. • If PPES recurs, withhold QINLOCK until Grade ≤ 1 or baseline and then resume QINLOCK at a reduced dose regardless of time to improvement.
	Grade 3	<ul style="list-style-type: none"> • Withhold QINLOCK for at least 7 days or until Grade ≤ 1 or baseline (maximum 28 days). Resume QINLOCK at a reduced dose. • Consider re-escalating QINLOCK if maintained at Grade ≤ 1 or baseline for at least 28 days.
Hypertension [see Section 4.4 Special Warnings and Precautions for Use]	Grade 3	<ul style="list-style-type: none"> • If symptomatic, withhold QINLOCK until symptoms have resolved and blood pressure is controlled. • If blood pressure is controlled to Grade ≤ 1 or baseline, resume QINLOCK at the same dose; otherwise, resume QINLOCK at reduced dose. • If Grade 3 hypertension recurs, withhold QINLOCK until symptoms have resolved and blood pressure is controlled. Resume QINLOCK at a reduced dose.
	Grade 4 Life-threatening consequences (e.g., malignant hypertension, transient or permanent neurologic deficit, hypertensive crisis)	Permanently discontinue QINLOCK.
Left Ventricular Systolic Dysfunction	Grade 3 or 4	Permanently discontinue QINLOCK.

Adverse Reaction	Severity ^a	Dosage Modifications
[see Section 4.4 Special Warnings and Precautions for Use]		
Arthralgia or Myalgia *	Grade 2	<ul style="list-style-type: none"> Withhold QINLOCK until Grade ≤ 1 or baseline. If recovered within 7 days, resume QINLOCK at same dose; otherwise, resume QINLOCK at reduced dose. Consider re-escalating QINLOCK if maintained at Grade ≤ 1 or baseline for at least 28 days. If arthralgia or myalgia recurs, withhold QINLOCK until Grade ≤ 1 or baseline and then resume QINLOCK at a reduced dose regardless of time to improvement.
	Grade 3	<ul style="list-style-type: none"> Withhold QINLOCK for at least 7 days or until Grade ≤ 1 or baseline (maximum of 28 days). Resume QINLOCK at a reduced dose. Consider re-escalating QINLOCK if maintained at Grade ≤ 1 or baseline for at least 28 days.
Other adverse reactions	Grade 3 or 4	<ul style="list-style-type: none"> Withhold QINLOCK until Grade ≤ 1 or baseline (maximum 28 days), and then resume QINLOCK at a reduced dose; otherwise, permanently discontinue. Consider re-escalating QINLOCK if no recurrence of the adverse reaction for at least 28 days. If Grade 3 or 4 recurs, permanently discontinue QINLOCK.

^a Graded per National Cancer Institute Common Terminology Criteria for Adverse Events version 4.03 (NCI CTCAE v4.03).

*Grade 4 events for Palmar-Plantar Erythrodysesthesia Syndrome (PPES) and Arthralgia or myalgia does not exist in CTCAE

Paediatrics

The safety and effectiveness of QINLOCK in paediatric patients have not been established.

Patients with renal impairment

No dose adjustment is recommended for patients with mild and moderate renal impairment [creatinine clearance (CrCl 30 to 89 mL/min estimated by Cockcroft-Gault)]. The pharmacokinetics and safety of QINLOCK in patients with severe renal impairment (CrCl 15 to 29 mL/min estimated by Cockcroft-Gault) have not been studied. No dosing recommendation can be made in patients with severe renal impairment.

Patients with hepatic impairment

No dose adjustment is recommended in patients with hepatic impairment (Child-Pugh A, B, or C) [see Section 5.2 Pharmacokinetic Properties].

Dose modifications for CYP3A inducers

Avoid concomitant strong or moderate CYP3A inducers during QINLOCK treatment.

If co-administration of a moderate CYP3A inducer can't be avoided, increase the QINLOCK dosing frequency to twice daily during the co-administration period (e.g. from the recommended dose of 150 mg once daily to 150 mg twice daily). Monitor for clinical response and tolerability. If the concomitant moderate CYP3A inducer is discontinued, reduce the dose back to once daily, 14 days after the discontinuation of the moderate CYP3A inducer [see [Section 4.5](#) Interactions with other medicines and other forms of interactions].

If co-administration of a strong CYP3A inducer absolutely can't be avoided, the above approach could be considered, but must be balanced against a risk of reduced efficacy due to reduced exposure [see [Section 4.5](#) Interactions with other medicines and other forms of interactions]. Warn patients of this risk, and monitor for clinical response and tolerability.

For patients who need to take QINLOCK twice daily: advise patients to take a missed dose if less than 4 hours have passed since they were meant to take it. Otherwise, leave it as a missed dose and just take the next dose as scheduled.

4.3 CONTRAINDICATIONS

Use of QINLOCK is contraindicated in patients with hypersensitivity to ripretinib or to any other component of QINLOCK tablets.

4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE

Palmar-Plantar Erythrodysesthesia Syndrome

In the double-blind period of a randomised, placebo-controlled phase 3 trial (INVICTUS), Grade 1-2 palmar-plantar erythrodysesthesia syndrome (PPES) occurred in 21% of the 85 patients who received QINLOCK [see [Section 4.8](#) Adverse Effects (Undesirable Effects)]. PPES led to dose discontinuation in 1.2% of patients, dose interruption in 2.4% of patients, and dose reduction in 1.2% of patients.

Based on severity, withhold QINLOCK and then resume at same or reduced dose [see [Section 4.2](#) Dose and Method of Administration].

New Primary Cutaneous Malignancies

In INVICTUS, cutaneous squamous cell carcinoma (cuSCC) occurred in 4.7% of the 85 patients who received QINLOCK, with a median time to event of 4.6 months (range: 3.8 to 6 months). In the pooled safety population, cuSCC and keratoacanthoma occurred in 7% and 1.9% of 351 patients, respectively.

In INVICTUS, melanoma occurred in 2.4% of the 85 patients who received QINLOCK. In the pooled safety population, melanoma occurred in 0.9% of 351 patients.

Perform dermatologic evaluations when initiating QINLOCK and routinely during treatment. Manage suspicious skin lesions with excision and dermatopathologic evaluation. Continue QINLOCK at the same dose.

Hypertension

In INVICTUS, Grade 1-3 hypertension occurred in 14% of the 85 patients who received QINLOCK, including Grade 3 hypertension in 7% [see [Section 4.8](#) Adverse Effects (Undesirable Effects)].

Do not initiate QINLOCK in patients with uncontrolled hypertension. Adequately control blood pressure prior to initiating QINLOCK. Monitor blood pressure as clinically indicated during treatment with QINLOCK and initiate or adjust antihypertensive therapy as appropriate.

Based on severity, withhold QINLOCK and then resume at same or reduced dose or permanently discontinue [see [Section 4.2](#) Dose and Method of Administration].

Cardiac failure

In INVICTUS, cardiac failure occurred in 1.2% of the 85 patients who received QINLOCK. In the pooled safety population, cardiac failure (cardiac failure, cardiac failure acute, acute left ventricular failure, and diastolic dysfunction) occurred in 1.7% of 351 patients, including Grade 3 adverse reactions in 1.1%.

In INVICTUS, Grade 3 decreased ejection fraction occurred in 1.3% of the 77 patients who received QINLOCK and who had a baseline and at least one post-baseline echocardiogram. In the pooled safety population, Grade 3 decreased ejection fraction occurred in 1.9% of the 263 patients in the pooled safety population who received QINLOCK and who had a baseline and at least one post-baseline echocardiogram.

In INVICTUS, cardiac dysfunction led to dose discontinuation in 1.2% of the 85 patients who received QINLOCK. The safety of QINLOCK has not been assessed in patients with a baseline ejection fraction below 50%.

Assess ejection fraction by echocardiogram or MUGA scan prior to initiating QINLOCK and during treatment, as clinically indicated. Permanently discontinue QINLOCK for Grade 3 or 4 left ventricular systolic dysfunction [see [Section 4.2](#) Dose and Method of Administration].

Refer to section 4.6 Fertility, Pregnancy and Lactation for potential Embryo-fetal toxicity.

Risk of impaired wound healing

Impaired wound healing complications can occur in patients who receive drugs that inhibit the vascular endothelial growth factor (VEGF) signaling pathway. Therefore, QINLOCK has the potential to adversely affect wound healing.

Withhold QINLOCK for at least 1 week prior to elective surgery. Do not administer for at least 2 weeks following major surgery and until adequate wound healing. The safety of resumption of QINLOCK after resolution of wound healing complications has not been established.

Phototoxicity

Ripretinib exhibits a potential for phototoxicity. It is recommended to advise patients to avoid or minimise exposure to direct sunlight, sunlamps, and other sources of ultraviolet radiation due to the risk of phototoxicity associated with ripretinib. Patients should be instructed to use measures such as protective clothing (long sleeves and hat) and sunscreen with high sun protection factor (SPF) during treatment with QINLOCK and for at least one week after discontinuation of treatment.

Use in the elderly

Of the 85 patients in INVICTUS who received QINLOCK 150 mg orally once daily, 24% were between 65 to 74 years of age and 9% were 75 years of age or older. Clinical studies of QINLOCK did not include sufficient numbers of patients aged 65 and older to determine whether they respond differently from younger patients.

Paediatric use

The safety and effectiveness of QINLOCK in paediatric patients have not been established.

Effects on laboratory tests

See [Section 4.8](#) Adverse effects (Undesirable Effects).

4.5 INTERACTIONS WITH OTHER MEDICINES AND OTHER FORMS OF INTERACTIONS

Effect of other medications on QINLOCK

Since both ripretinib and its active metabolite are mainly cleared by CYP3A, plasma concentrations of ripretinib and the active metabolite DP-5439 are expected to be increased by CYP3A inhibitors and decreased by CYP3A inducers.

Effect of Strong CYP3A Inhibitors: Coadministration of QINLOCK with itraconazole (a strong CYP3A inhibitor) increased ripretinib C_{max} by 36% and $AUC_{0-\infty}$ by 99%. Use strong CYP3A inhibitors with caution and monitor for increased toxicity.

Effect of strong CYP3A inducers: Coadministration of QINLOCK with rifampin (a strong CYP3A inducer) decreased ripretinib C_{max} by 18% and $AUC_{0-\infty}$ by 61% and also decreased DP-5439 $AUC_{0-\infty}$ by 57% with increased C_{max} by 37% [see [Section 4.2](#) Dose and method of administration].

In the presence of a strong CYP3A inducer, a doubled ripretinib dose (twice daily rather than once daily), is predicted to result in a 40% reduction in combined AUC of ripretinib and active metabolite DP-5439, compared to the usual recommended once daily dose with no inducer present.

Avoid concomitant strong CYP3A inducers during QINLOCK treatment [see [Section 4.2](#) Dose and method of administration].

Effect of moderate CYP3A inducers: Coadministration of QINLOCK with efavirenz (a moderate CYP3A inducer) was predicted to decrease ripretinib C_{max} by 24% and decrease $AUC_{0-\infty}$ by 56% [see [Section 4.2](#) Dose and method of administration].

In the presence of a moderate CYP3A inducer, a doubled ripretinib dose (twice daily rather than once daily), is predicted to result in a 17% reduction in combined AUC of ripretinib and active

metabolite DP-5439, compared to the usual recommended once daily dose with no inducer present.

Avoid concomitant moderate CYP3A inducers during QINLOCK treatment [see [Section 4.2](#) Dose and method of administration].

Effect of P-gp inhibitors: Ripretinib and its active metabolite DP-5439 are substrates of P-gp (MDR1) and BCRP, as indicated by *in vitro* studies. Ripretinib is an inhibitor of P-gp and BCRP. P-gp inhibitors may increase plasma concentrations of ripretinib and its active metabolite DP-5439. DP-5439 is an inhibitor of BCRP and MATE1 (Multidrug And Toxin Extrusion Protein 1).

Effect of Acid-Reducing Agents: Coadministration of QINLOCK with pantoprazole (a proton pump inhibitor) did not affect exposure to ripretinib.

Effect of QINLOCK on other medications

Effect of CYP2C8 inhibitors: Ripretinib is a weak inhibitor of CYP2C8. Co-administration of QINLOCK with repaglinide (a sensitive index substrate for CYP2C8) increased repaglinide AUC_{0-∞} by 26%. Repaglinide C_{max} was unchanged; therefore, dose adjustment is not required.

Ripretinib and DP-5439 are not inducers of CYP1A2, CYP2B6, or CYP3A4.

UGT1A substrates

In vitro studies suggested that ripretinib is an inhibitor of UGT1A1, UGT1A3, UGT1A4, UGT1A7, and UGT1A8. Clinical studies with UGT1A substrates have not been conducted.

4.6 FERTILITY, PREGNANCY AND LACTATION

Effects on fertility

Based on findings from animal studies, QINLOCK may impair fertility in males of reproductive potential. Decreased testis and epididymis weights, as well as atrophy of the testes and degeneration of the seminiferous epithelium were observed in male rats at exposure levels (AUC) similar to the human exposure at 150 mg once daily.

Use in pregnancy

Category D

There are no clinical data on the use of QINLOCK in pregnant women to inform a drug-associated risk of major birth defects and miscarriage.

Based on findings from animal studies, QINLOCK can cause embryo-fetal harm when administered to a pregnant woman. QINLOCK should not be used during pregnancy.

QINLOCK should not be used during pregnancy. In an embryo-fetal development study in which pregnant rats were administered daily doses of ripretinib during organogenesis, ripretinib was given from gestational days 6 through 18 at doses 1, 5, or 20 mg/kg/day. Dose-related malformations primarily associated with the cardiovascular and skeletal systems were observed at a dose of 20 mg/kg/day (approximately 0.4 times the human exposure at 150 mg once daily).

An increased incidence of anatomic variations, indicative of developmental toxicity, also occurred at 20 mg/kg/day. Variations included malpositioned carotid and subclavian artery origins, malpositioned subclavian artery, absent or elongated innominate artery, misshapen and nodulated ribs, bipartite, incompletely ossified, or unossified vertebral centra, small or misshapen vertebral arches, and reductions in ossified forelimb and hindlimb phalanges, hindlimb metatarsals, and caudal vertebrae were also observed at 20 mg/kg/day.

Verify the pregnancy status of females of reproductive potential prior to initiating QINLOCK. Advise female patients of reproductive potential to use effective contraception during treatment and for at least 1 week after the final dose. Advise male patients with female partners of reproductive potential to use effective contraception during treatment and for at least 1 week after the final dose. Effects of QINLOCK on contraceptive steroids have not been studied. A barrier method contraception should be added if systemic contraceptive steroids are used.

Use in lactation

There are no data regarding the presence of ripretinib or its metabolites in human milk or their effects on the breastfed infant or on milk production. Because of the potential for adverse reactions in breastfed infants, advise women not to breastfeed during treatment with ripretinib and for at least 1 week after the final dose.

4.7 EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

No data available.

4.8 ADVERSE EFFECTS (UNDESIRABLE EFFECTS)

Summary of the safety profile

In the Phase 3 double-blind, randomised (2:1), placebo-controlled trial (INVICTUS), 129 study participants with a diagnosis of advanced GIST were randomised to QINLOCK (N=85) or placebo (N=44) [see [Section 5.1 Pharmacodynamic Properties](#)]. The data described in this section reflect the safety population (N=128) who had received at least one dose of QINLOCK (N=85) or placebo (N=43). One study participant who was randomised to the placebo arm did not receive placebo. The safety results from the double-blind treatment period of INVICTUS are described below.

The most common adverse events ($\geq 20\%$) observed in patients treated with QINLOCK (all grades) were alopecia, fatigue, nausea, abdominal pain, constipation, myalgia, diarrhoea, decreased appetite, palmar-plantar erythrodysesthesia syndrome (PPES), and vomiting ([Table 2](#)). The most common Grade 3 or 4 laboratory abnormalities were increased lipase and decreased phosphate ([Table 3](#)).

For additional information on palmar-plantar erythrodysesthesia syndrome, new primary cutaneous malignancies such as squamous cell carcinoma of the skin, keratoacanthoma and melanoma, hypertension and cardiac failure (cardiac failure, cardiac failure acute, acute left ventricular failure, and diastolic dysfunction) see section 4.4 Special warnings and precautions for use.

Tabulated list of adverse events

Table 2 summarises the most frequently reported treatment-emergent adverse events (TEAEs) in $\geq 10\%$ of patients with advanced GIST who received QINLOCK in the double-blind treatment period of INVICTUS.

Table 2: Treatment-Emergent Adverse Events (TEAEs) Reported in $\geq 10\%$ of Patients Who Received QINLOCK with a Difference Between Arms of $>5\%$ Compared to Placebo in INVICTUS

Treatment-Emergent Adverse Events	QINLOCK (N=85)		Placebo (N=43)	
	Grades 1-4 %	Grades 3-4 %	Grades 1-4 %	Grades 3-4 %
Skin and subcutaneous tissue				
Alopecia	52	0	4.7	0
Palmar-plantar erythrodysesthesia syndrome	21	0	0	0
Dry skin	13	0	7	0
Pruritus	11	0	4.7	0
General				
Fatigue	42	3.5	23	2.3
Peripheral oedema	17	1.2	7	0
Asthenia	13	1.2	14	4.7
Gastrointestinal				
Nausea	39	3.5	12	0
Abdominal pain	36	7	30	4.7
Constipation	34	1.2	19	0
Diarrhoea	28	1.2	14	2.3
Vomiting	21	3.5	7	0
Stomatitis	11	0	0	0
Musculoskeletal and connective tissue				
Myalgia	32	1.2	12	0
Arthralgia	18	0	4.7	0
Muscle spasms	15	0	4.7	0
Metabolism and nutrition				
Decreased appetite	27	1.2	21	2.3
Investigations				
Decreased weight	19	0	12	0
Nervous system				
Headache	19	0	4.7	0
Vascular				
Hypertension	14	7	4.7	0
Respiratory, thoracic and mediastinal				
Dyspnoea	13	0	0	0

Table 3 summarises the most frequently reported treatment-emergent laboratory abnormalities in $\geq 10\%$ of patients with advanced GIST who received QINLOCK in the double-blind treatment period of INVICTUS.

Table 3: Laboratory Abnormalities Reported in ≥10% of Patients Who Received QINLOCK in the Double-Blind Treatment Period of INVICTUS^a

Laboratory Abnormality	QINLOCK ^a (N=85)		Placebo ^a (N=43)	
	Grades 1-4 %	Grades 3-4 ^b %	Grades 1-4 %	Grades 3-4 %
Haematology				
Increased activated partial thromboplastin time	35	0	9	0
Increased INR	21	3.8	15	0
Decreased neutrophil count	10	0	2.5	0
Chemistry				
Increased lipase	32	7	13	8
Decreased phosphate	26	4.9	2.5	0
Increased triglycerides	26	2.4	23	0
Decreased calcium	23	0	8	0
Increased blood bilirubin	22	0	5	2.5
Increased CPK	21	1.2	10	0
Decreased sodium	17	2.4	10	2.5
Increased creatinine	16	0	18	0
Increased serum amylase	13	1.2	5	0
Increased ALT	12	1.2	5	0
CPK=creatine phosphokinase; INR=international normalised ratio; ALT=alanine aminotransferase				
a. The denominator used to calculate the rate varied from 82 to 83 for QINLOCK and 34 to 40 for placebo based on the number of patients with a baseline value and at least one post-treatment value.				
b. Only includes Grade 3 laboratory abnormalities.				

Table 4: Dose Interruptions, Dose Reductions, and Treatment Discontinuations due to Adverse Reactions

Event	QINLOCK (N=85) %	Placebo (N=43) %
Dose interruption	23.5	20.9
Dose reduction	7.1	2.3
Treatment discontinuation	8.2	11.6

Other Adverse Reactions

The following additional adverse reactions have occurred in <10% of QINLOCK-treated patients in INVICTUS:

Peripheral sensory neuropathy (7.1% vs 2.3% placebo), dermatitis acneiform (5.9% vs 0% placebo), rash (4.7% vs 4.7% placebo), seborrheic keratosis (4.7% vs 0% placebo), skin papilloma (3.5% vs 0% placebo) and melanocytic naevus (2.4% vs 0% placebo).

Clinically relevant adverse reactions that occurred in <10% of patients in the pooled safety population included cardiac ischemic events (1.1%) (including acute coronary syndrome and fatal cardiac arrest or myocardial infarction). Photosensitivity occurred in 0.6% [see Section 4.4 Special Warnings and Precautions for Use].

Reporting suspected adverse effects

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at <https://pophealth.my.site.com/carmreportnz/s/> and drugsafety-STA@stbiopharma.com.

4.9 OVERDOSE

There is no known specific antidote for QINLOCK overdose. In the event of suspected overdose, interrupt QINLOCK, undertake general supportive measures, and observe until clinical stabilisation.

For risk assessment and advice on the management of overdose, contact the National Poisons Centre on 0800 POISON (0800 764 766).


5 PHARMACOLOGICAL PROPERTIES

5.1 PHARMACODYNAMIC PROPERTIES

Physicochemical properties

Chemical structure

The chemical structure of ripretinib is shown below:



Chemical name: 1-(4-bromo-5-[1-ethyl-7-(methylamino)-2-oxo-1,2-dihydro-1,6-naphthyridin-3-yl]-2-fluorophenyl)-3-phenylurea

Molecular Formula: C₂₄H₂₁BrFN₅O₂

Molecular Weight: 510.36 g/mol

CAS number

1442472-39-0

Mechanism of action

Ripretinib is a switch-control tyrosine kinase inhibitor with a dual mechanism of action. Ripretinib binds to both the switch pocket and the activation loop to lock the kinase in the inactive state, preventing downstream signalling and cell proliferation. This dual mechanism of action provides broad inhibition of KIT and PDGFRA kinase activity, including wild type and multiple primary and secondary mutations. Ripretinib also inhibits other kinases *in vitro*, such as PDGFRB, TIE2, VEGFR2, and BRAF.

Cardiac Electrophysiology

Following treatment with QINLOCK 150 mg once daily, no clinically meaningful QT interval prolongation was observed.

Clinical trials

The efficacy of QINLOCK was evaluated in INVICTUS, an international, multi-centre, randomised (2:1), double-blind, placebo-controlled trial. Eligible patients had unresectable, locally advanced or metastatic gastrointestinal stromal tumour (GIST) and had received prior treatment with imatinib, sunitinib, and regorafenib. Randomisation was stratified by prior lines of therapy (3 versus ≥4) and Eastern Cooperative Oncology Group (ECOG) performance status (0 versus 1 or 2). Patients received QINLOCK 150 mg or placebo orally once daily until disease progression or unacceptable toxicity. Tumour response assessments were performed every 28 days through for the first 4 months and then every 56 days thereafter.

The major efficacy outcome measure was progression-free survival (PFS) based on disease assessment by blinded independent central review (BICR) using modified RECIST 1.1 criteria, in which lymph nodes and bone lesions were not target lesions and a progressively growing new tumour nodule within a pre-existing tumour mass must meet specific criteria to be considered unequivocal evidence of progression. Additional efficacy outcome measures included objective

response rate (ORR) by BICR and overall survival (OS). Patients randomised to receive placebo could be treated with QINLOCK at the time of disease progression.

A total of 129 patients were randomised, 85 to QINLOCK and 44 to placebo.

Patient characteristics of the intent-to-treat (ITT) population in INVICTUS were median age of 60 years (range: 29 to 83 years), with 39% aged ≥ 65 years; 57% were male; 75% were White; and 92% had an ECOG performance status of 0 or 1. Sixty-three percent (63%) of patients received 3 prior therapies and 37% received 4 or more prior therapies. Sixty-six percent (66%) of patients randomised to placebo switched to QINLOCK after disease progression.

Efficacy results from INVICTUS are summarised in [Table 5](#).

Table 5. Efficacy Results of INVICTUS

	QINLOCK (N=85)	Placebo (N=44)
Progression-Free Survival (PFS)^a		
Number of events (%)	51 (60)	37 (84)
Progressive disease	46 (54)	32 (73)
Deaths	5 (6)	5 (11)
Median PFS (months) (95% CI)	6.3 (4.6, 6.9)	1.0 (0.9, 1.7)
Hazard ratio (95% CI) ^b	0.15 (0.09, 0.25)	
p-value ^c	< 0.0001	
Overall Response Rate (ORR)^a		
Overall Response Rate (%)	9	0
(95%, CI)	(4.2, 18)	(0, 8)
p-value ^d	0.0504	
Overall Survival (OS)^e		
Number of Deaths, N (%)	26 (31)	26 (59)
Median OS (months) (95% CI) ^e	15.1 (12.3, 15.1)	6.6 (4.1, 11.6)
Hazard Ratio (95% CI) ^c	0.36 (0.21, 0.62)	
BICR=Blinded Independent Central Review; CI=Confidence Interval		
a. Assessed per BICR		
b. Hazard ratio is based on Cox proportional hazards regression model. This model includes treatment and randomisation stratification factors as fixed factors.		
c. p-value is based on 2-sided stratified Log Rank test.		

	QINLOCK (N=85)	Placebo (N=44)
d.	Based on Fisher's exact test. The p-value is not statistically significant.	
e.	Not evaluated for statistical significance as a result of the sequential testing procedure for the secondary endpoints of ORR and OS.	

Figure 1: Kaplan-Meier Curve of Progression-Free Survival in INVICTUS





Figure 2: Kaplan-Meier Curve of Overall Survival in INVICTUS



5.2 PHARMACOKINETIC PROPERTIES

Absorption

Ripretinib reaches peak plasma concentrations at 4 hours after a single oral dose of 150 mg ripretinib (given as three tablets each containing 50 mg). The steady state AUC_{0-12h} observed in patients at 150 mg is 5678 ng•h/mL. Steady state is achieved by approximately Day 15. Administration with a high-fat meal increased ripretinib AUC_{0-24} and C_{max} by 30% and 22%, respectively. DP-5439 AUC_{0-24} and C_{max} were higher by 47% and 66%, respectively.

Distribution

Both ripretinib and its active metabolite DP-5439 bind to plasma proteins at $\geq 99\%$. The apparent volume of distribution (V_{ss}/F) is approximately 307 L.

Metabolism

Ripretinib was metabolised *in vitro*. CYP3A4/5 is the major metaboliser of ripretinib and its active metabolite, DP-5439, while CYP2C8 and CYP2D6 are only minor metabolisers.

Excretion

Following oral administration of ripretinib 150 mg once daily, the mean apparent oral clearance (CL/F) of ripretinib at steady-state is 15.3 L/hr and the mean plasma elimination half-life is 14.8 hours

In preclinical species, ^{14}C -labeled ripretinib dosed to Sprague-Dawley rats (oral) and beagle dogs (intravenous [iv]), resulted in greater than 87% of the radioactive dose being excreted in faeces and 1.8% or less in the urine.

PK analyses obtained from urine and faeces samples in 10 healthy volunteers showed that systemic elimination of ripretinib was not primarily attributed to the kidney. Through 1 week (168 hours) after a single oral administration of 50 mg ripretinib (given alone), 0.02% of the ripretinib dose was excreted as ripretinib in urine and 34.2% of the ripretinib dose was excreted as ripretinib in faeces.

Pharmacokinetics in special patient populations

Population pharmacokinetic analyses of demographic data indicate that no clinically meaningful differences in the pharmacokinetics of ripretinib were observed based on age (19 to 87 years), sex, race (White, Black, and Asian), body weight (39 to 138 kg), tumour type (GIST or other solid tumour), prior gastrectomy, mild to moderate renal impairment (CrCl 30 to < 90 mL/min estimated by Cockcroft-Gault) and mild hepatic impairment (total bilirubin \leq ULN and AST $>$ ULN or total bilirubin 1 to $1.5 \times$ ULN and any AST).

The effects of severe renal impairment (CrCl 15 to 29 mL/min) on the pharmacokinetics of ripretinib have not been studied.

Patients with Hepatic Impairment

In subjects with mild hepatic impairment, there were no clinically significant changes in AUC_{0-last} and C_{max} when compared to matched healthy subjects.

In subjects with moderate hepatic impairment, ripretinib AUC_{0-last} increased 2-fold while C_{max} was unchanged when compared to matched healthy subjects. The combined AUC_{0-last} of ripretinib and DP-5439 increased 1.5-fold.

In subjects with severe hepatic impairment, ripretinib AUC_{0-last} increased 2.6-fold and C_{max} decreased by 24% when compared to matched healthy subjects. The combined AUC_{0-last} of ripretinib and DP-5439 increased by 1.4-fold.

The observed magnitude of increase in ripretinib and the combination of ripretinib and DP-5439 exposures in subjects with hepatic impairment is unlikely to be clinically relevant based on the known safety profile of ripretinib.

5.3 PRECLINICAL SAFETY DATA

Genotoxicity

Ripretinib was not mutagenic in an *in vitro* bacterial reverse mutation (Ames) assay or clastogenic in an *in vivo* rat bone marrow micronucleus assay. Ripretinib was weakly positive in an *in vitro* clastogenicity assay in human lymphocytes without metabolic activation. Ripretinib's active metabolite (DP-5439) was not mutagenic in an *in vitro* bacterial reverse mutation test or clastogenic in an *in vitro* chromosomal aberration assay in isolated human lymphocytes. Ripretinib is not expected to pose a genotoxic risk.

Carcinogenicity

Carcinogenicity studies have not been conducted with ripretinib.

Phototoxicity

Ripretinib indicates a potential for photoirritation/phototoxicity based on absorption in the UV visible range (above 290 nm). *In vitro* phototoxicity assessment in 3T3 mouse fibroblast cells suggest that ripretinib exhibits a potential for phototoxicity at clinically relevant concentrations following exposure to UVA and UVB radiation.

6 PHARMACEUTICAL PARTICULARS

6.1 LIST OF EXCIPIENTS

Each QINLOCK tablet contains the following inactive ingredients:

- crospovidone;
- hypromellose acetate succinate;
- lactose monohydrate;
- magnesium stearate;
- microcrystalline cellulose;
- silicon dioxide.

6.2 INCOMPATIBILITIES

No incompatibilities have been identified.

6.3 SHELF LIFE

48 months stored below 25°C and protected from light.

6.4 SPECIAL PRECAUTIONS FOR STORAGE

Store below 25°C. Protect from light.

6.5 NATURE AND CONTENTS OF CONTAINER

QINLOCK 50 mg tablets are packaged with silica gel desiccant into white high-density polyethylene (HDPE) bottles. The bottles are closed with polypropylene child resistant closures with a polyethylene-faced induction heat seal liner. Each HDPE bottle contains 90 tablets.

6.6 SPECIAL PRECAUTIONS FOR DISPOSAL

In New Zealand, any unused medicine or waste material should be disposed of by taking to your local pharmacy.

7 MEDICINE SCHEDULE (POISONS STANDARD)

S4 – Prescription Only Medicine

8 SPONSOR

Specialised Therapeutics Limited
Level 1, The Lane, Botany Town Centre
588 Chapel Road, East Tamaki
Auckland, 2013
New Zealand
Phone: +64 9801 0299

Website: www.stbiopharma.com

9 DATE OF FIRST APPROVAL

19 January 2023

10 DATE OF REVISION

24 JUNE 2025

SUMMARY TABLE OF CHANGES

Section Changed	Summary of new information
4.4	Update to EF values under Cardiac failure
4.5	Update to information under Effect of QINLOCK on other medications