

# NEW ZEALAND DATA SHEET

## ABILIFY MAINTENA<sup>®</sup>

### 1 NAME OF THE MEDICINE

Abilify Maintena Injection with diluent 300 mg

Abilify Maintena Injection with diluent 400 mg

### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Abilify Maintena Injection with diluent 300mg: each vial contains 300 mg aripiprazole (as monohydrate).

Abilify Maintena Injection with diluent 400 mg: each vial contains 400 mg aripiprazole (as monohydrate).

For the full list of excipients, see section 6.1.

### 3 PHARMACEUTICAL FORM

Aripiprazole monohydrate is a white-to-off-white crystalline powder. Aripiprazole monohydrate is practically insoluble in water.

Abilify Maintena is presented as a sterile, single-dose, lyophilised powder for reconstitution with water for injections to give a prolonged-release suspension for injection to deliver 400 mg of aripiprazole in 400-mg/vial strength or 300 mg of aripiprazole in 300-mg/vial strength.

Abilify Maintena powder and solvent for prolonged-release suspension for injection is available as a one-month therapeutic kit. Each therapeutic kit contains one vial of lyophilised powder, one vial of water for injections, one 3 mL sterile syringe with a 21 gauge needle for reconstitution, one 3 mL sterile syringe without a needle, one 1-inch (25 mm) 23 gauge sterile safety needle for injection; one 1.5-inch (38 mm) 22 gauge sterile safety needle for injection and one 2-inch (51 mm) 21 gauge sterile safety needle for injection and one vial adapter.

### 4 CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

For the acute and maintenance treatment of schizophrenia in adults.

For maintenance treatment to prevent the recurrence of manic or mixed episodes of bipolar I disorder in adult patients as monotherapy.

#### 4.2 Dose and method of administration

##### *Recommended Dosage and Dosage Adjustment*

Abilify Maintena is intended for intramuscular injection only. For patients who have never taken aripiprazole, establish tolerability with oral aripiprazole prior to initiating treatment with Abilify Maintena. Due to the half-life of oral aripiprazole, it may take up to 2 weeks to fully assess tolerability.

The recommended starting and maintenance dose of Abilify Maintena is 400 mg. Titration of the dose of Abilify Maintena is not required. Abilify Maintena is to be administered by a healthcare professional only, once-monthly as a single injection into the deltoid or gluteal muscle (no sooner than 26 days after the previous injection). Sites of injections should be rotated between the two gluteal or deltoid muscles.

After the first Abilify Maintena injection, treatment with oral aripiprazole (10 mg to 20 mg), or other oral antipsychotic, should be continued for 14 consecutive days to maintain therapeutic antipsychotic concentrations during initiation of therapy.

If there are adverse reactions with the 400 mg dose, reduction of the dose to 300 mg once-monthly should be considered.

### ***Switching from oral antipsychotics***

For patients who have never taken oral or injectable aripiprazole, establish tolerability with oral aripiprazole prior to initiating treatment with Abilify Maintena. When switching from oral antipsychotics, patients may continue their current oral antipsychotic (oral aripiprazole or prescribed dose of other oral antipsychotic) for 14 days following the first dose of Abilify Maintena to maintain therapeutic plasma concentrations during the initiation of Abilify Maintena. Abilify Maintena should then be administered once monthly as described above.

### ***Switching from long-acting injectable antipsychotics***

For patients who have never taken oral or injectable aripiprazole, establish tolerability with oral aripiprazole prior to initiating treatment with Abilify Maintena. When switching patients from previous long-acting injectable antipsychotics, initiate Abilify Maintena therapy in place of the next scheduled injection, with 14 days of concurrent oral aripiprazole. Abilify Maintena should then be continued monthly.

### ***Discontinuation of Abilify Maintena***

If Abilify Maintena is discontinued, its prolonged-release characteristics must be considered.

### ***Missed doses***

**Table 1 Management of missed doses**

<b>If 2<sup>nd</sup> or 3<sup>rd</sup> dose is missed and time since last injection is:</b>	<b>Action</b>
>4 weeks and <5 weeks	The injection should be administered as soon as possible and then resume monthly injection schedule.
>5 weeks	Concomitant oral aripiprazole should be restarted for 14 days with next administered injection and then resume monthly injection schedule.
<b>If 4<sup>th</sup> or subsequent doses are missed and time since last injection is:</b>	<b>Action</b>
>4 weeks and <6 weeks	The injection should be administered as soon as possible and then resume monthly injection schedule.
>6 weeks	Concomitant oral aripiprazole should be restarted for 14 days with next administered injection and then resume monthly injection schedule.

### ***Dosage in Special Populations***

#### **Elderly Population**

The effectiveness and safety of Abilify Maintena in the treatment of schizophrenia in patients  $\geq 61$  years and in the treatment of bipolar I disorder in patients  $\geq 66$  years have not been evaluated.

## Renal Impairment

No dosage adjustment of Abilify Maintena is required for patients with renal impairment. See 5.2 Pharmacokinetic properties - Special Populations.

## Hepatic Impairment

Based on oral data no dosage adjustment of Abilify Maintena is required for patients with mild or moderate hepatic impairment. In patients with severe hepatic impairment, the data available are insufficient to establish dosage recommendations. In these patients dosing should be managed cautiously; use of oral aripiprazole should be considered. See 5.2 Pharmacokinetic properties - Special Populations.

## Other Special Populations

No dosage adjustment of Abilify Maintena is recommended based on gender, race or smoking status.

## Known CYP2D6 Poor Metabolisers

In patients who are known to be CYP2D6 poor metabolisers, the starting and maintenance dose of Abilify Maintena should be 300 mg. If Abilify Maintena is taken concomitantly with strong CYP3A4 inhibitors, the dose of Abilify Maintena should be reduced to 200 mg. See 4.5 Interaction with other medicines and other forms of interaction.

## Paediatric Population

The safety and efficacy of Abilify Maintena in children and adolescents aged 0-17 years have not been established. No data are available.

## Dose Adjustments due to Interactions

Dosage adjustments are recommended in patients taking concomitant strong CYP3A4 inhibitors or strong CYP2D6 inhibitors for more than 14 days. If the CYP3A4 inhibitor or CYP2D6 inhibitor is withdrawn, the Abilify Maintena dose may need to be increased. Refer to **Table 2**.

Avoid the concomitant use of CYP3A4 inducers with Abilify Maintena for more than 14 days, because the blood levels of aripiprazole will be decreased and may fall below the effective levels. Refer to **Table 2**.

**Table 2 Dose Adjustments of Abilify Maintena in patients taking concomitant strong CYP2D6 inhibitors, and/or strong CYP3A4 inhibitors, and/or CYP3A4 inducers for more than 14 days**

	Adjusted Dose
<b>Patients Taking 400 mg of Abilify Maintena</b>	
Strong CYP2D6 or Strong CYP3A4 inhibitors	300 mg
Strong CYP2D6 and Strong CYP3A4 inhibitors	200 mg
CYP3A4 inducers	Avoid use
<b>Patients Taking 300 mg of Abilify Maintena</b>	
Strong CYP2D6 or Strong CYP3A4 inhibitors	200 mg
Strong CYP2D6 and Strong CYP3A4 inhibitors	160 mg
CYP3A4 inducers	Avoid use

## Administration

Abilify Maintena is available in a therapeutic kit which contains one vial of lyophilised powder, one vial of diluent (water for injections), one 3 mL sterile syringe with a 21 gauge needle for reconstitution, one 3 mL sterile syringe without a needle, one 1-inch (25 mm) 23 gauge sterile safety needle for injection, one 1.5-inch (38 mm) 22 gauge sterile safety needle for injection, and one 2-inch (51 mm) 21 gauge sterile safety needle for injection and one vial adapter.

***For deep intramuscular deltoid or gluteal injection only: do not administer intravenously or subcutaneously.***

**The suspension should be injected immediately after reconstitution, but can be stored below 25°C for up to 4 hours in the vial. If the injection is not performed immediately after reconstitution, shake the vial vigorously for at least 60 seconds to re-suspend prior to injection. Abilify Maintena should be administered by a healthcare professional only, once-monthly. The injection should be injected slowly as a single injection (doses must not be divided) into the deltoid or gluteal muscle.**

Any unused product or waste material should be disposed of in accordance with local requirements.

## Reconstitution

### Step 1: Preparation Prior to Reconstitution of the Lyophilised Abilify Maintena Powder.

- a) Lay out the contents of the package and confirm that all components listed below are provided:
  - One vial containing aripiprazole 400 mg or 300 mg (as monohydrate) powder for reconstitution;
  - One vial containing water for injections 2.0 mL;
  - One 3 mL luer lock syringe with pre-attached 21 gauge, 1.5-inch (38 mm) hypodermic safety needle with needle protection device;
  - One 3 mL disposable syringe with luer lock tip;
  - One vial adapter;
  - One 23 gauge, 1-inch (25 mm) hypodermic safety needle with needle protection device;
  - One 22 gauge, 1.5-inch (38 mm) hypodermic safety needle with needle protection device; and
  - One 21 gauge, 2-inch (51 mm) hypodermic safety needle with needle protection device.
- b) The contents of the vial containing aripiprazole (as monohydrate) lyophilised powder should be suspended using the water for injections supplied in the carton.
- c) The vials containing aripiprazole (as monohydrate) lyophilised powder and water for injections are for single-use only.
- d) Use appropriate aseptic techniques throughout the reconstitution and reconstitute at room temperature.

- e) Select the amount of water for injections needed for reconstitution at room temperature.

400 mg Vial		300 mg Vial	
Dose	Water for Injections	Dose	Water for Injections
400 mg	1.9 mL	300 mg	1.5 mL

**It is important to note that there is more water for injections in the vial than is needed to reconstitute the contents of the vial containing aripiprazole (as monohydrate) lyophilised powder for prolonged-release suspension for injection.**

### **Step 2: Reconstitution of the Lyophilised Powder**

- Remove the cap of the vial of water for injections and remove the cap of the vial containing aripiprazole (as monohydrate) lyophilised powder and wipe the tops with a sterile alcohol swab.
- Using the syringe with pre-attached hypodermic safety needle, withdraw the pre-determined water for injections volume from the vial of water for injections into the syringe (see Diagram 1). A small amount of residual water for injections will remain in the vial following withdrawal.



Diagram 1

- Slowly inject the water for injections into the vial containing the aripiprazole (as monohydrate) lyophilised powder (see Diagram 2).




Diagram 2

- Withdraw air to equalise the pressure in the vial by pulling back slightly on the plunger. Subsequently, remove the needle from the vial. Engage the needle safety device by using the one-handed technique (see Diagram 3). Gently press the sheath against a flat surface until the needle is firmly engaged in the needle protection sheath. Visually confirm that the needle is fully engaged into the needle protection sheath, and discard.




Diagram 3

- e) Shake the vial vigorously for 30 seconds until the suspension appears uniform (see Diagram 4).



Diagram 4

- f) Visually inspect the reconstituted suspension for particulate matter and discolouration prior to administration. The reconstituted Abilify Maintena is a uniform, homogeneous suspension that is opaque and milky-white in colour. Do not use if reconstituted suspension contains particulate matter or any discolouration.
- g) If the injection is not performed immediately after reconstitution, keep the vial below 25°C for up to 4 hours and shake the vial vigorously for at least 60 seconds to re-suspend prior to injection.
- h) Do not store the reconstituted suspension in the syringe.

### Step 3: Preparation Prior to Injection

- a) Use appropriate aseptic techniques throughout injection of the reconstituted Abilify Maintena suspension.
- b) Remove the cover from the vial adapter package (see Diagram 5). Do not remove the vial adapter from the package.



Diagram 5

- c) Using the vial adapter package to handle the vial adapter, attach the pre-packaged disposable syringe to the vial adapter (see Diagram 6).



Diagram 6 Use the disposable syringe to remove the vial adapter from the package and discard the vial adapter package (see Diagram 7). Do not touch the spike tip of the adapter at any time.




Diagram 7

- d) Determine the recommended volume for injection.

**Abilify Maintena Reconstituted Suspension Volume to Inject**

400 mg Vial		300 mg Vial	
Dose	Volume to Inject	Dose	Volume to Inject
400 mg	2.0 mL	---	---
300 mg	1.5 mL	300 mg	1.5 mL
200 mg	1.0 mL	200 mg	1.0 mL
160 mg	0.8 mL	160 mg	0.8 mL

- e) Wipe the top of the vial of the reconstituted Abilify Maintena suspension with a sterile alcohol swab.
- f) Place and hold the vial of the reconstituted Abilify Maintena suspension on a hard surface. Attach the adapter-syringe assembly to the vial by holding the outside of the adapter and pushing the adapter's spike firmly through the rubber stopper, until the adapter snaps in place (see Diagram 8).




Diagram 8

- g) Slowly withdraw the recommended volume from the vial into the disposable syringe to allow for injection (see Diagram 9). A small amount of excess product will remain in the vial.



Diagram 9

#### Step 4: Injection Procedure

- Detach the disposable syringe containing the recommended volume of reconstituted Abilify Maintena suspension from the vial.
- Select one of the hypodermic safety needles listed in the table below according to the injection site and patient's weight.

Body weight	Injection site	Needle size
Non-obese	Deltoid	25 mm (1 inch) 23 gauge
	Gluteal	38 mm (1.5 inch) 22 gauge
Obese	Deltoid	38 mm (1.5 inch) 22 gauge
	Gluteal	51 mm (2 inch) 21 gauge

Attach the needle to the disposable syringe containing the suspension for injection. To avoid subcutaneous drug administration, examine the depth of subcutaneous fat at the injection site and select the appropriate needle size. Ensure the needle is firmly seated on the needle safety device with a push and clockwise twist and then pull the needle cap straight away from the needle (see Diagram 10).



Diagram 10

- Slowly inject the recommended volume as a single intramuscular injection into the deltoid or gluteal muscle (see Diagram 11). Do not massage the injection site. Do not administer intravenously or subcutaneously.






Diagram 11

### Step 5: Procedures After Injection

- Engage the needle safety device as described in Step 2 (d). Dispose of the vials, adapter, needles, and syringe appropriately after injection. **The vials containing aripiprazole (as monohydrate) lyophilised powder and water for injections are for single-use only.**
- Rotate sites of injections between the two deltoid and gluteal muscles. Look for signs or symptoms of inadvertent intravenous administration.

### 4.3 Contraindications

Hypersensitivity to aripiprazole or any of the excipients. See 6.1 List of excipients.

### 4.4 Special warnings and precautions for use

#### *Elderly patients with dementia-related psychosis*

##### **Increased mortality**

Elderly patients with dementia-related psychosis treated with atypical antipsychotic drugs, including aripiprazole, are at an increased risk of death compared to placebo. Analyses of 17 placebo-controlled trials (modal duration of 10 weeks) in these patients revealed a risk of death in the drug-treated patients of between 1.6 to 1.7 times that seen in placebo-treated patients. Over the course of a typical 10-week controlled trial, the rate of death in drug-treated patients was about 4.5%, compared to a rate of about 2.6% in the placebo group. Although the causes of death varied, most of the deaths appeared to be either cardiovascular (e.g., heart failure, sudden death) or infectious (e.g., pneumonia) in nature.

In three placebo-controlled trials with oral aripiprazole in elderly patients with psychosis associated with Alzheimer's disease (n= 938; mean age: 82.4 years; range: 56-99 years), patients treated with aripiprazole were at an increased risk of death compared to placebo. The rate of death in the aripiprazole-treated patients was 3.5% compared with 1.7% in the placebo group. Abilify Maintena is not indicated for the treatment of patients with dementia-related psychosis.

##### **Cerebrovascular adverse events**

In the same three 10-week placebo-controlled trials, cerebrovascular adverse reactions (e.g., stroke, transient ischaemic attack), including fatalities, were reported in patients (mean age: 84 years; range: 78-88 years). Overall, 1.3% of aripiprazole-treated patients reported cerebrovascular adverse reactions compared with 0.6% of placebo-treated patients in these trials. However, in one of these trials, a fixed-dose trial, there was a significant dose response relationship for cerebrovascular adverse reactions in patients treated with aripiprazole. The safety and efficacy of aripiprazole on the treatment of patients

with psychosis associated with dementia have not been established. Abilify Maintena is not indicated for the treatment of patients with dementia-related psychosis.

### ***General***

During antipsychotic treatment, improvement in the patient's clinical condition may take several days to some weeks. Patients should be closely monitored during this period.

### ***Suicide***

The possibility of a suicide attempt is inherent in psychotic illnesses and close supervision of high-risk patients should accompany drug therapy. As Abilify Maintena is to be administered by a healthcare professional, suicide due to an overdose is considered unlikely.

### ***Tardive dyskinesia***

In clinical trials of one year or less duration, there were uncommon reports of treatment-emergent tardive dyskinesia during treatment with aripiprazole. The risk of tardive dyskinesia increases with long-term exposure to antipsychotic treatment. If signs and symptoms of tardive dyskinesia appear in a patient on Abilify Maintena, dose reduction or discontinuation of treatment should be considered (see 4.2 Dose and method of administration). These symptoms can temporally deteriorate or can even arise after discontinuation of treatment.

### ***Neuroleptic malignant syndrome***

Neuroleptic Malignant Syndrome (NMS) is a potentially fatal symptom complex associated with antipsychotic medicinal products. In clinical trials, rare cases of NMS were reported during treatment with aripiprazole. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional signs may include elevated creatine phosphokinase, myoglobinuria (rhabdomyolysis), and acute renal failure. However, elevated creatine phosphokinase and rhabdomyolysis, not necessarily in association with NMS, have also been reported. If a patient develops signs and symptoms indicative of NMS, or presents with unexplained high fever without additional clinical manifestations of NMS, all antipsychotic medicinal products, including aripiprazole, must be discontinued.

### ***Seizure***

In clinical trials, uncommon cases of seizures were experienced during treatment with aripiprazole. As with other antipsychotic drugs, Abilify Maintena should be used cautiously in patients who have a history of seizure disorder or have conditions associated with seizures.

### ***Hyperglycaemia and diabetes mellitus***

Hyperglycaemia, in some cases extreme and associated with ketoacidosis or hyperosmolar coma or death, has been reported in patients treated with atypical antipsychotics including aripiprazole. In clinical trials, the observed differences in the incidence rates of hyperglycaemia-related adverse reactions (including diabetes) or in abnormal glycaemia laboratory values between Abilify Maintena (<1%) and placebo (0%) could be considered as of no major clinical concern.

Assessment of the relationship between atypical antipsychotic use and glucose abnormalities is complicated by the possibility of an increased background risk of diabetes mellitus in patients with schizophrenia and the increasing incidence of diabetes mellitus in the general population. Given these confounders, the relationship between atypical antipsychotic use and hyperglycaemia-related adverse events is not completely understood. However, epidemiological studies suggest an increased risk of treatment-emergent hyperglycaemia-related adverse events in patients treated with atypical

antipsychotics. Precise risk estimates for hyperglycaemia-related adverse events in patients treated with atypical antipsychotics are not available.

Patients with an established diagnosis of diabetes mellitus who are started on atypical antipsychotics, including aripiprazole, should be monitored regularly for worsening of glucose control. All patients who are starting treatment with atypical antipsychotics, including aripiprazole, should undergo fasting blood glucose testing at the beginning of treatment and periodically during treatment. Any patient treated with atypical antipsychotics, including aripiprazole, should be monitored for symptoms of hyperglycaemia including polydipsia, polyuria, polyphagia and weakness. Patients who develop symptoms of hyperglycaemia during treatment with atypical antipsychotics should undergo fasting blood glucose testing. In some cases, hyperglycaemia has resolved when the atypical antipsychotic was discontinued; however, some patients required continuation of anti-diabetic treatment despite discontinuation of the suspect drug.

In patients with significant treatment-emergent hyperglycaemia, discontinuation of Abilify Maintena should be considered.

### ***Cardiovascular disorders***

Aripiprazole should be used with caution in patients with known cardiovascular disease (e.g., history of myocardial infarction or ischemic heart disease, heart failure, or conduction abnormalities), cerebrovascular disease, or conditions which would predispose patients to hypotension (e.g. dehydration, hypovolaemia, and treatment with antihypertensive medications) or hypertension, including accelerated or malignant. Patients with a history of clinically significant cardiovascular disorders were excluded from clinical trials.

### ***QT interval***

In clinical trials of treatment with Abilify Maintena and oral aripiprazole, the incidence of QT prolongation was uncommon. As with other antipsychotics, aripiprazole should be used with caution in patients with a family history of QT prolongation. See also 4.8 Adverse effects (Undesirable effects) - Laboratory Tests - QT Interval.

### ***Orthostatic hypotension***

Aripiprazole may be associated with orthostatic hypotension, potentially due to its  $\alpha_1$ -adrenergic receptor antagonism. Aripiprazole may induce orthostatic hypotension, tachycardia, dizziness and sometimes syncope, especially at the initiation of treatment. In the double-blind controlled phase of the long-term clinical trials using Abilify Maintena, orthostasis-related events were reported in 2/534 (0.4%) patients. In the 12-week clinical trial in acutely relapsed patients, orthostasis-related events were reported in 1/167 (0.6%) patient treated with Abilify Maintena, while syncope and orthostatic hypotension each occurred in 2/172 (1.2%) patients treated with placebo.

### ***Venous thromboembolism***

Cases of venous thromboembolism (VTE) have been reported with antipsychotic drugs. Since patients treated with antipsychotics often present with acquired risk factors for VTE, all possible risk factors for VTE should be identified before and during treatment with Abilify Maintena and preventive measures undertaken.

### ***Falls***

Somnolence, postural hypotension, motor, and sensory instability have been reported with the use of antipsychotics, including aripiprazole, which may lead to falls. Caution should be taken when treating patients with diseases, conditions, or who are taking medications that could exacerbate these effects.

### ***Body temperature regulation***

Disruption of the body's ability to increase or reduce core body temperature has been attributed to antipsychotic agents, including aripiprazole. Appropriate care is advised when prescribing Abilify Maintena for patients who will be experiencing conditions which may contribute to an elevation in core body temperature, e.g., exercising strenuously, exposure to extreme heat, receiving concomitant medication with anticholinergic activity or being subject to dehydration.

Patients should be advised regarding appropriate care in avoiding overheating and dehydration.

### ***Dysphagia***

Oesophageal dysmotility and aspiration have been associated with antipsychotic drug use, including aripiprazole. Abilify Maintena and other antipsychotic drugs should be used cautiously in patients at risk of aspiration pneumonia (e.g., elderly patients).

### ***Akathisia***

*Class effect:* The presentation of akathisia may be variable and comprises subjective complaints of restlessness and an overwhelming urge to move and either distress or motor phenomena (such as pacing, swinging of the legs while seated, rocking from foot to foot), or both. Particular attention should be paid to the monitoring for such symptoms and signs as, left untreated, akathisia is associated with poor compliance and an increased risk of relapse.

### ***Leukopenia, neutropenia and agranulocytosis***

*Class Effect:* In clinical trial and/or post-marketing experience, events of leukopenia/neutropenia have been reported temporally related to antipsychotic agents, including aripiprazole. Agranulocytosis has also been reported.

Possible risk factors for leukopenia/neutropenia include pre-existing low white blood cell count (WBC) and history of drug-induced leukopenia/neutropenia. Patients with a history of a clinically significant low WBC or drug-induced leukopenia/neutropenia should have their complete blood cell (CBC) monitored frequently during the first few months of therapy, and discontinuation of Abilify Maintena should be considered at the first sign of a clinically significant decline in WBC in the absence of other causative factors.

Patients with clinically significant neutropenia should be carefully monitored for fever or other symptoms or signs of infection and treated promptly, if such symptoms or signs occur. Patients with severe neutropenia (absolute neutrophil count  $<1000/\text{mm}^3$ ) should discontinue Abilify Maintena and have their WBC followed until recovery. See 4.8 Adverse effects (Undesirable effects) - Laboratory Tests.

### ***Pathological gambling and impulse control disorders***

Patients can experience increased urges, particularly for gambling, and the inability to control these urges while taking aripiprazole. Other urges reported include: increased sexual urges, compulsive spending, binge or compulsive eating, and other impulsive and compulsive behaviours. It is important for prescribers to ask patients or their caregivers specifically about the development of new or increased gambling urges, sexual urges, compulsive spending, binge or compulsive eating, or other urges while being treated with aripiprazole. It should be noted that impulse-control symptoms can be associated with the underlying disorder; however, in some cases urges were reported to have stopped when the dose was reduced or the medication was discontinued. Impulse control disorders may result in harm to the patient and others if not recognized. Consider dose reduction or stopping the medication if a patient develops such urges while taking aripiprazole. See 4.8 Adverse Effects (Undesirable effects) - Post-Market Adverse Drug Reactions.

### ***Weight gain***

Antipsychotic drugs have been associated with metabolic changes, including weight gain. In long-term clinical trials there was no clinically relevant difference in the incidence of weight gain between Abilify Maintena and placebo. In a 12-week clinical trial in acutely relapsed patients, a potentially clinically relevant difference in the incidence of weight gain between Abilify Maintena and placebo was seen. See 4.8 Adverse Effects (Undesirable effects) - Weight.

All patients should have baseline and periodic monitoring of body weight, and other cardiometabolic parameters, including fasting glucose, full lipid profile and blood pressure, during treatment with any atypical antipsychotic including Abilify Maintena.

### ***Paediatric use***

The safety and efficacy of Abilify Maintena in children and adolescents aged 0-17 years have not been established. No data are available.

### ***Use in the elderly***

There are no data on the safety and efficacy of Abilify Maintena in patients  $\geq 61$  years of age.

## **4.5 Interaction with other medicines and other forms of interaction**

While no specific drug interaction studies have been performed with Abilify Maintena, the effects of co-administration of inhibitors of CYP2D6 and CYP3A4 were modelled as part of a population pharmacokinetic study but with no data accrued. The information below is therefore obtained from studies with oral aripiprazole.

### ***Potential for other medicinal products to affect aripiprazole***

Aripiprazole is metabolised by multiple pathways involving the CYP2D6 and CYP3A4 enzymes.

### **Inhibitors and Inducers of CYP2D6 and CYP3A4**

#### **Quinidine or other strong CYP2D6 inhibitors**

In a clinical trial of oral aripiprazole in healthy subjects, a strong inhibitor of CYP2D6 (quinidine) decreased oral clearance of aripiprazole by 52%, increased aripiprazole AUC by 107%, while  $C_{\max}$  was unchanged. The AUC and  $C_{\max}$  of dehydro-aripiprazole, the active metabolite, decreased by 32% and 47%, respectively. Other strong inhibitors of CYP2D6, such as fluoxetine, paroxetine, and bupropion may be expected to have similar effects and similar dose reduction should, therefore, be applied. See 4.2 Dose and method of administration, Dosage in Special Populations - Dose Adjustments due to Interactions.

#### **Ketoconazole or other strong CYP3A4 inhibitors**

In a clinical trial of oral aripiprazole in healthy subjects, a strong inhibitor of CYP3A4 (ketoconazole) decreased oral clearance of aripiprazole by 38%, increased aripiprazole AUC and  $C_{\max}$  by 63% and 37%, respectively. The AUC and  $C_{\max}$  of dehydro-aripiprazole increased by 77% and 43%, respectively. Other strong inhibitors of CYP3A4, such as itraconazole and HIV protease inhibitors may be expected to have similar effects and similar dose reductions should, therefore, be applied (see 4.2 Dose and method of administration, Dosage in Special Populations - Dose Adjustments due to Interactions). When considering concomitant administration of ketoconazole or other potent CYP3A4 inhibitors with Abilify Maintena, potential benefits should outweigh the potential risks to the patient.

Upon discontinuation of the CYP2D6 or CYP3A4 inhibitor, the dose of Abilify Maintena should be increased to the dose prior to the initiation of the concomitant therapy.

### **Carbamazepine or other CYP3A4 inducers**

In a clinical study in patients with schizophrenia or schizo-affective disorder, co-administration of carbamazepine (200 mg twice daily), a potent CYP3A4 inducer, with aripiprazole (30 mg daily) resulted in an approximate 70% decrease in AUC values of both aripiprazole and its active metabolite, dehydro-aripiprazole. Concomitant administration of Abilify Maintena and other potent inducers of CYP3A4 (such as rifampicin, rifabutin, phenytoin, phenobarbital, primidone, efavirenz, nevirapine and St. John's Wort) or potent inducers of CYP2D may be expected to have similar effects.

### **Valproate and lithium**

When either valproate or lithium was administered concomitantly with aripiprazole, there was no clinically significant change in aripiprazole concentrations.

### **Inhibitors and inducers of CYP1A1, CYP1A2, CYP2C9 and CYP2C19**

Aripiprazole is not metabolised by CYP1A1, CYP1A2, CYP2C9 and CYP2C19 *in vitro*, suggesting that interactions with medications or other factors (e.g., smoking), which are inhibitors or inducers of these enzymes, are unlikely.

### ***Potential for aripiprazole to affect other medicinal products***

#### **CNS Drugs (including Alcohol)**

Given the primary CNS effects of aripiprazole, caution should be used when Abilify Maintena is administered in combination with alcohol or other CNS drugs with overlapping adverse reactions, such as sedation. See 4.8 Adverse effects (Undesirable effects).

Patients should be advised to avoid alcohol while on Abilify Maintena.

When aripiprazole was administered concomitantly with valproate, lithium, lamotrigine, dextromethorphan, warfarin, omeprazole, escitalopram, venlafaxine or desvenlafaxine there was no clinically important change in concentrations of these drugs.

### **Effects of aripiprazole on substrates for CYP2D6, CYP2C9, CYP2C19, CYP3A4 and CYP1A2**

In clinical studies, oral doses of 10-30 mg/day of aripiprazole had no significant effect on the metabolism of substrates of CYP2D6 (dextromethorphan/3-methoxymorphinan ratio), 2C9 (warfarin), 2C19 (omeprazole), and 3A4 (dextromethorphan). Additionally, aripiprazole and dehydro-aripiprazole did not show potential for altering CYP1A2-mediated metabolism *in vitro*. Thus, Abilify Maintena is unlikely to cause clinically important medicinal product interactions mediated by these enzymes.

### **Antihypertensive agents**

Due to its  $\alpha_1$ -adrenergic receptor antagonistic activity, aripiprazole has the potential to enhance the effect of certain antihypertensive agents.

### **Medicines which cause QT prolongation or electrolyte imbalance**

If aripiprazole is administered concomitantly with medicinal products known to cause QT prolongation or electrolyte imbalance, caution should be used.

### **Effects on laboratory tests**

Drug interaction with laboratory tests has not been established.

## 4.6 Fertility, pregnancy and lactation

### *Effects on fertility*

Reproductive toxicity studies have not been performed on Abilify Maintena. The following information is taken from studies performed on oral aripiprazole, which showed that aripiprazole did not impair fertility in reproductive toxicity studies.

Aripiprazole had no effect on fertility in female rats treated orally with 2, 6, and 20 mg/kg/day (0.6, 2, and 6 times the oral maximum recommended human dose (MRHD) of 30 mg/day based on mg/m<sup>2</sup>) for 2 weeks prior to mating through gestation day 7. Drug-related effects (persistent dioestrus, and increased mating time, pre-implantation losses, and corpora lutea) observed at all doses were considered the result of perturbed oestrous cyclicity secondary to drug-mediated hyperprolactinaemia.

Aripiprazole had no effect on fertility in male rats treated with oral doses of 20, 40, and 60 mg/kg/day (6, 12, and 18 times the oral MRHD of 30 mg/day based on mg/m<sup>2</sup>) for 9 weeks prior to mating through mating. Disturbances of spermatogenesis were seen at 60 mg/kg/day and prostatic atrophy was seen at 40 and 60 mg/kg/day.

### *Use in pregnancy*

#### Pregnancy Category C

There are no adequate and well-controlled studies of aripiprazole in pregnant women. It is not known whether aripiprazole can cause fetal harm when administered to a pregnant woman or can affect reproductive capacity. Patients must be advised to notify their doctors if they become pregnant or intend to become pregnant during treatment with aripiprazole.

In animal studies, aripiprazole demonstrated developmental toxicity, including possible teratogenic effects in rats and rabbits. Pregnant rats were treated with oral doses of 3, 10, and 30 mg/kg/day (1, 3, and 9 times the oral Maximum Recommended Human Dose [MRHD] of 30 mg/day on a mg/m<sup>2</sup> basis) of aripiprazole during the period of organogenesis. At 30 mg/kg in the rat, treatment was associated with slightly prolonged gestation, and a slight delay in fetal development, as evidenced by decreased fetal weight, undescended testes, and delayed skeletal ossification. There were no adverse effects on embryofetal or pup survival. Delivered offspring had decreased bodyweights, and increased incidences of hepatodiaphragmatic nodules and diaphragmatic hernia at 30 mg/kg (the other dose groups were not examined for these findings). A low incidence of diaphragmatic hernia was also seen in the fetuses exposed to 30 mg/kg. Postnatally, delayed vaginal opening was seen at 10 and 30 mg/kg and impaired reproductive performance (decreased fertility rate, corpora lutea, implants, and live fetuses and increased post implantation loss, likely mediated through effects on female offspring) was seen at 30 mg/kg. Some maternal toxicity was seen at 30 mg/kg; however, there was no evidence to suggest that these developmental effects were secondary to maternal toxicity.

Pregnant rabbits were treated with oral doses of 10, 30, and 100 mg/kg/day (2, 3, and 11 times human exposure at the oral MRHD of 30 mg/day based on AUC) of aripiprazole during the period of organogenesis. Decreased maternal food consumption and increased abortions were seen at 100 mg/kg. Treatment caused increased fetal mortality (100 mg/kg), decreased fetal weight (30 and 100 mg/kg), increased incidence of a skeletal abnormality (fused sternebrae at 100 mg/kg), and minor skeletal variations (100 mg/kg).

Rats were treated with oral doses of 3, 10, and 30 mg/kg/day (1, 3, and 9 times the oral MRHD of 30 mg/day on a mg/m<sup>2</sup> basis) of aripiprazole from late gestation through weaning. At 30 mg/kg, maternal toxicity, slightly prolonged gestation, an increase in stillbirths, poor postnatal care/nursing, and decreases in pup weight (persisting into adulthood) and survival were seen.

*Non-teratogenic class effect:* Neonates exposed to antipsychotic drugs (including aripiprazole) during the third trimester of pregnancy are at risk of experiencing extrapyramidal neurological disturbances and/or withdrawal symptoms following delivery. There have been post-marketing reports of agitation, hypertonia, hypotonia, tremor, somnolence, respiratory distress, and feeding disorder in these neonates. These complications have varied in severity; while in some cases symptoms have been self-limited; in other cases neonates have required additional medical treatment or monitoring.

Abilify Maintena should be used during pregnancy only if the anticipated benefit to the mother outweighs the potential risk to the fetus and the administered dose and duration of treatment should be as low and as short as possible.

#### ***Use in labour and delivery***

The effect of aripiprazole on labour and delivery in humans is unknown.

#### ***Use in lactation***

Aripiprazole has been found in human breast milk. Patients should be advised not to breastfeed if they are taking Abilify Maintena.

In rats, there were adverse effects in dams and offspring following daily oral administration of aripiprazole from late gestation through weaning. See 4.6 Fertility, pregnancy and lactation – Use in Pregnancy.

### **4.7 Effects on ability to drive and use machines**

Abilify Maintena, like other antipsychotics, may have the potential to impair judgment, thinking, or motor skills. Patients should be cautioned about operating hazardous machinery, including motor vehicles, until they are reasonably certain that Abilify Maintena therapy does not affect them adversely.

The effects of this medicine on a person's ability to drive and use machines were not assessed as part of its registration.

### **4.8 Adverse effects (Undesirable effects)**

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare professionals are asked to report any suspected adverse reactions <https://pophealth.my.site.com/carmreportnz/s/>.

Abilify Maintena administered once monthly has been evaluated for safety in 3,453 adult patients in clinical trials in schizophrenia and bipolar I disorder. Of the 3,453 adult patients exposed to aripiprazole long-acting injectable, 2,567 patients have been treated with Abilify Maintena 400/300 mg. Of the 3,453 patients exposed to Abilify Maintena 1,226 patients have received at least 13 Abilify Maintena 400 mg/300 mg injections (i.e., have been treated for at least 12 months).

Abilify has been evaluated for safety in 13543 patients who participated in multiple-dose clinical trials across all approved indications including schizophrenia and bipolar I disorder, and who had approximately 7619 patient-years of exposure to oral aripiprazole and 749 patients with exposure to aripiprazole short acting injection. A total of 3390 patients were treated with oral aripiprazole for at least 180 days and 1933 patients treated with oral aripiprazole had at least 1 year of exposure.



## Adult patients with Schizophrenia

Abilify Maintena administered once monthly has been evaluated for safety in clinical trials in adult patients with schizophrenia. Of the 2,649 adult patients exposed to aripiprazole long-acting injectable, 2,567 patients have been treated with Abilify Maintena 400/300 mg.

The most frequently observed treatment emergent adverse events (TEAE) reported in  $\geq 5\%$  of patients treated with Abilify Maintena 300 mg – 400 mg in the two double-blind long term clinical trials were insomnia (10.9%), weight increased (9.4%), akathisia (8.1%), headache (7.9%), anxiety (6.6%), decreased weight (6.6%), nasopharyngitis (5.8%), and injection site pain (5.2%). Overall, treatment emergent adverse events (TEAEs) were similar to placebo and the majority were mild to moderate in severity. The TEAEs that occurred in the two double-blind long-term clinical trials with Abilify Maintena at a frequency of  $\geq 2\%$  are listed in **Table 3**.

**Table 3 Treatment–Emergent Adverse Events (TEAE) Reported for  $\geq 2\%$  of patients in both placebo and active controlled long-term clinical trials**

<b>System Organ Class MedDRA Preferred Term</b>	<b>ABILIFY MAINTENA™ 400 mg/300 mg (N = 534)</b>	<b>Oral Aripiprazole 10-30 mg (N = 266)</b>	<b>Aripiprazole IM Depot 50 mg/25 mg (N = 131)</b>	<b>Placebo (N = 134)</b>
	<b>n (%)</b>	<b>n (%)</b>	<b>n (%)</b>	<b>n (%)</b>
<b>Any TEAE</b>	<b>389 (72.8)</b>	<b>213 (80.1)</b>	<b>106 (80.9)</b>	<b>83 (61.9)</b>
<b>Gastrointestinal Disorders</b>				
Abdominal pain upper	4 (0.7)	4 (1.5)	3 (2.3)	0 (0.0)
Diarrhoea	15 (2.8)	9 (3.4)	6 (4.6)	3 (2.2)
Nausea	10 (1.9)	4 (1.5)	3 (2.3)	2 (1.5)
Toothache	14 (2.6)	13 (4.9)	3 (2.3)	3 (2.2)
Vomiting	12 (2.2)	4 (1.5)	1 (0.8)	3 (2.2)
<b>General Disorders and Administration Site Conditions</b>				
Fatigue	11 (2.1)	9 (3.4)	2 (1.5)	1 (0.7)
Injection site pain	28 (5.2)	6 (2.3)	1 (0.8)	5 (3.7)
Oedema peripheral	4 (0.7)	3 (1.1)	3 (2.3)	3 (2.2)
<b>Infections and Infestations</b>				
Bronchitis	7 (1.3)	5 (1.9)	5 (3.8)	2 (1.5)
Influenza	16 (3.0)	11 (4.1)	7 (5.3)	2 (1.5)
Nasopharyngitis	31 (5.8)	25 (9.4)	9 (6.9)	7 (5.2)
Pharyngitis	5 (0.9)	0 (0.0)	3 (2.3)	1 (0.7)
Upper respiratory tract infection	25 (4.7)	11 (4.1)	5 (3.8)	3 (2.2)
<b>Investigations</b>				

<b>System Organ Class MedDRA Preferred Term</b>	<b>ABILIFY MAINTENA™ 400 mg/300 mg (N = 534)</b>	<b>Oral Aripiprazole 10-30 mg (N = 266)</b>	<b>Aripiprazole IM Depot 50 mg/25 mg (N = 131)</b>	<b>Placebo (N = 134)</b>
	<b>n (%)</b>	<b>n (%)</b>	<b>n (%)</b>	<b>n (%)</b>
<b>Any TEAE</b>	<b>389 (72.8)</b>	<b>213 (80.1)</b>	<b>106 (80.9)</b>	<b>83 (61.9)</b>
Blood creatine phosphokinase increased	10 (1.9)	6 (2.3)	5 (3.8)	2 (1.5)
Blood pressure increased	6 (1.1)	1 (0.4)	0 (0.0)	3 (2.2)
Weight decreased	35 (6.6)	16 (6.0)	12 (9.2)	4 (3.0)
Weight increased	50 (9.4)	35 (13.2)	7 (5.3)	13 (9.7)
<b>Metabolism and Nutrition Disorders</b>				
Decreased appetite	6 (1.1)	1 (0.4)	3 (2.3)	0 (0.0)
<b>Musculoskeletal and Connective Tissue Disorders</b>				
Arthralgia	15 (2.8)	4 (1.5)	0 (0.0)	1 (0.7)
Back pain	16 (3.0)	14 (5.3)	15 (11.5)	3 (2.2)
Pain in extremity	11 (2.1)	7 (2.6)	2 (1.5)	6 (4.5)
<b>Nervous System Disorders</b>				
Akathisia	43 (8.1)	18 (6.8)	11 (8.4)	8 (6.0)
Dizziness	14 (2.6)	6 (2.3)	2 (1.5)	4 (3.0)
Headache	42 (7.9)	30 (11.3)	7 (5.3)	7 (5.2)
Sedation	13 (2.4)	3 (1.1)	1 (0.8)	1 (0.7)
Somnolence	14 (2.6)	12 (4.5)	2 (1.5)	1 (0.7)
Tremor	24 (4.5)	9 (3.4)	6 (4.6)	2 (1.5)
<b>Psychiatric Disorders</b>				
Agitation	9 (1.7)	2 (0.8)	0 (0.0)	3 (2.2)
Anxiety	35 (6.6)	13 (4.9)	10 (7.6)	10 (7.5)
Depression	7 (1.3)	3 (1.1)	0 (0.0)	3 (2.2)
Insomnia	58 (10.9)	37 (13.9)	18 (13.7)	12 (9.0)
Psychotic disorder	16 (3.0)	8 (3.0)	8 (6.1)	9 (6.7)
Restlessness	16 (3.0)	4 (1.5)	4 (3.1)	3 (2.2)
Schizophrenia	10 (1.9)	5 (1.9)	10 (7.6)	5 (3.7)
<b>Respiratory, Thoracic and Mediastinal Disorders</b>				
Cough	14 (2.6)	7 (2.6)	5 (3.8)	4 (3.0)

<b>System Organ Class MedDRA Preferred Term</b>	<b>ABILIFY MAINTENA™ 400 mg/300 mg (N = 534)</b>	<b>Oral Aripiprazole 10-30 mg (N = 266)</b>	<b>Aripiprazole IM Depot 50 mg/25 mg (N = 131)</b>	<b>Placebo (N = 134)</b>
	<b>n (%)</b>	<b>n (%)</b>	<b>n (%)</b>	<b>n (%)</b>
<b>Any TEAE</b>	<b>389 (72.8)</b>	<b>213 (80.1)</b>	<b>106 (80.9)</b>	<b>83 (61.9)</b>
Nasal congestion	3 (0.6)	1 (0.4)	1 (0.8)	3 (2.2)
<b>Vascular Disorders</b>				
Rash	2 (0.4)	4 (1.5)	4 (3.1)	1 (0.7)
Hypertension	7 (1.3)	4 (1.5)	4 (3.1)	3 (2.2)

The most frequently observed treatment emergent adverse event (TEAE) reported in ≥5% of patients treated with Abilify Maintena in the 12-week clinical trial in acutely relapsed patients were weight increased (16.8%), headache (14.4%), akathisia (11.4%), constipation (9.6%), cough (6.0%), dyspepsia (6.0%), agitation (5.4%), injection site pain (5.4%), sedation (5.4%) and toothache (5.4%). The TEAEs that occurred in at least 2% of patients and greater than placebo during the active treatment phase of the 12-week clinical trial in patients in the acute phase of schizophrenia are listed in **Table 4**.

**Table 4 - Incidence of Treatment-emergent Adverse Events Occurring in ≥ 2% of Aripiprazole IM Depot Patients with Schizophrenia and Greater than Placebo in the Acute Treatment Phase of the Acute Phase Placebo-controlled Trial**

<b>System organ class MedDRA preferred term</b>	<b>Aripiprazole IM Depot 400/300mg (N=167)</b>	<b>Placebo (N=172)</b>
	<b>n (%)</b>	<b>n (%)</b>
<b>Gastrointestinal Disorders</b>		
Abdominal discomfort	4 (2.4)	2 (1.2)
Constipation	16 (9.6)	12 (7.0)
Diarrhoea	5 (3.0)	4 (2.3)
Dry mouth	6 (3.6)	4 (2.3)
Toothache	9 (5.4)	8 (4.7)
Vomiting	5 (3.0)	2 (1.2)
<b>General Disorders and Administration Site Conditions</b>		
Fatigue	4 (2.4)	3 (1.7)
Injection site pain	9 (5.4)	1 (0.6)
<b>Infections and Infestations</b>		
Upper respiratory tract infection	6 (3.6)	3 (1.7)
<b>Investigations</b>		
Weight decreased	6 (3.6)	4 (2.3)

System organ class MedDRA preferred term	Aripiprazole IM Depot 400/300mg (N=167)	Placebo (N=172)
	n (%)	n (%)
Weight increased	28 (16.8)	12 (7.0)
<b>Musculoskeletal and Connective Tissue Disorders</b>		
Arthralgia	6 (3.6)	2 (1.2)
Back pain	7 (4.2)	4 (2.3)
Musculoskeletal pain	5 (3.0)	2 (1.2)
Myalgia	6 (3.6)	1 (0.6)
<b>Nervous System Disorders</b>		
Akathisia	19 (11.4)	6 (3.5)
Dizziness	6 (3.6)	3 (1.7)
Sedation	9 (5.4)	2 (1.2)
Tremor	5 (3.0)	1 (0.6)
<b>Psychiatric Disorders</b>		
Insomnia	8 (4.8)	8 (4.7)
<b>Respiratory, Thoracic and Mediastinal Disorders</b>		
Cough	10 (6.0)	10 (5.8)
Nasal congestion	4 (2.4)	2 (1.2)

### Adult patients with Bipolar I Disorder

Abilify Maintena administered once monthly has been evaluated for safety in clinical trials in 804 adult patients with bipolar I disorder.

In a double-blind, placebo-controlled, randomized withdrawal (maintenance) study, the most frequently observed TEAEs that were reported in  $\geq 5\%$  of patients receiving Abilify Maintena and with a greater frequency than in the placebo group were weight increased (23.5%), akathisia (21.2%), insomnia (7.6%) and anxiety (6.8%).

**Table 5 Incidence of Treatment-emergent Adverse Events Occurring in  $\geq 2\%$  of Aripiprazole IM Depot Patients and Greater than Placebo in the Placebo-controlled Phase of the Double-blind Trial in Adult Patients with Bipolar I Disorder**

System Organ Class MedDRA Preferred Term	Aripiprazole IM Depot (N=132) n (%)	Placebo (N=133) n (%)
<b>Subject With Any Treatment Emergent Adverse Events</b>	101 (76.5)	107 (80.5)
<b>Blood and Lymphatic System Disorders</b>		
Anaemia	3 (2.3)	0 (0.0)
<b>Eye Disorders</b>		

<b>System Organ Class MedDRA Preferred Term</b>	<b>Aripiprazole IM Depot (N=132) n (%)</b>	<b>Placebo (N=133) n (%)</b>
Vision Blurred	3 (2.3)	0 (0.0)
<b>Gastrointestinal Disorders</b>		
Constipation	4 (3.0)	4 (3.0)
Dry Mouth	4 (3.0)	3 (2.3)
Salivary Hypersecretion	3 (2.3)	3 (2.3)
<b>Infections and Infestations</b>		
Bronchitis	3 (2.3)	2 (1.5)
Influenza	3 (2.3)	2 (1.5)
Sinusitis	5 (3.8)	1 (0.8)
Urinary Tract Infection	4 (3.0)	2 (1.5)
<b>Injury, Poisoning and Procedural Complications</b>		
Procedural Pain	4 (3.0)	1 (0.8)
<b>Investigations</b>		
Blood Creatine Phosphokinase Increased	3 (2.3)	1 (0.8)
Weight Increased	31(23.5)	24 (18.0)
<b>Metabolism and Nutrition Disorders</b>		
Increased Appetite	4 (3.0)	1 (0.8)
<b>Nervous System Disorders</b>		
Akathisia	28 (21.2)	17 (12.8)
Somnolence	6 (4.5)	1 (0.8)
Tremor	3 (2.3)	2 (1.5)
<b>Psychiatric Disorders</b>		
Anxiety	9 (6.8)	6 (4.5)
Bipolar Disorder	5 (3.8)	5 (3.8)
Depression	4 (3.0)	3 (2.3)
Insomnia	10 (7.6)	10 (7.5)
Libido Decreased	3 (2.3)	2 (1.5)
Restlessness	6 (4.5)	5 (3.8)

### ***Injection Site Adverse Events***

Injection site assessments were completed after all injections during the double-blind, controlled phases of the two Abilify Maintena long-term trials and the acute schizophrenia trial. Analyses of injection site assessments (investigator-rated and subject reported Visual Analogue Scale (VAS)) were performed to evaluate the safety/tolerability of Abilify Maintena, see

. In both long-term trials, infrequent injection site reactions were observed; those seen were generally mild to moderate in severity, and resolved over time. Injection site pain (incidence 5.2%), has a median

onset on day 2 after the injection and a median duration of 4 days. In the acute phase trial, injection site pain (incidence 5.4%) was reported to be mild in severity, and resolved over time.

In the double-blind, placebo-controlled phase in patients with bipolar I disorder, the percentage of patients reporting injection site-related adverse reactions was similar between Abilify Maintena and placebo. The mean intensity of injection pain reported by patients on Abilify Maintena using a visual analog scale (0=no pain to 100=unbearably painful) was 5.2 (SD 9.0) for the first injection and 4.0 (SD 7.6) for the last injection in the double-blind, placebo-controlled phase.

**Table 6 Investigator Assessments of Pain, Swelling and Induration at the Injection Site and Patient VAS scores**

Treatment Group Dose (n)	Absence of Investigator Rated Pain, Redness, Swelling, and Induration at the Injection Site (% of patients) *		Mean VAS (patient rated pain from 0 mm (no pain) – 100 mm (unbearably painful)	
	First Injection	Last Injection	First Injection	Last Injection
<b>38-week Double-blind, Active-controlled trial, in Patients with Schizophrenia</b>				
<b>Double-blind, Active-controlled Phase</b>				
<b>Abilify Maintena</b> 400 mg/300 mg (n=265)	81.4 – 98.1	88.3 – 98.9	5.6	3.7
Oral aripiprazole 10-30 mg (n=266)	83.3 – 98.5	90.2 – 99.6	4.9	3.5
Aripiprazole Long-Acting Injectable 50 mg/25 mg (n=131)	90.7 – 99.2	90.0 – 99.2	3.3	2.4
<b>52-week Double-blind, Placebo-controlled Trial in Patients with Schizophrenia</b>				
<b>Long-Acting Injectable Stabilisation Phase (Open Label)**</b>				
<b>Abilify Maintena</b> 400 mg/300 mg (n=403)	75.3 – 96.2	77.3 – 97.0	6.0	4.5
<b>Double-blind, Placebo-controlled Phase</b>				
<b>Abilify Maintena</b> 400 mg/300 mg (n=269)	80.1 – 98.1	84.4 – 98.5	5.1	4.0
Placebo (n=134)	72.2 – 97.7	77.3 – 97.7	5.1	4.9
<b>12-week Double-blind, Placebo-controlled Acute Phase Trial in Patients with Schizophrenia</b>				
<b>Double-blind, Acute Treatment Phase</b>				
<b>Abilify Maintena</b> (n=167)	95.1-100	99.2-100	7.1	7.7
Placebo (n=172)	94.7-100	98.4-100	5.7	8.6
<b>52-week Double-blind, Placebo-controlled Trial in Patients with Bipolar I Disorder</b>				
<b>Double-blind, Placebo-controlled Phase</b>				
<b>Abilify Maintena</b> 400 mg/300 mg (n=132)	81.8-100	87.1-100	5.2	4.0
Placebo (n=133)	81.2-100	85-100	5.7	5.8

VAS = Visual Analogue Scale.

\*Range of per-cent is based on rating in the 4 domains (pain, redness, swelling, and induration)

\*\*The open-label analyses were done to understand the injection site reaction parameters after initiation of Abilify Maintena as well as during its continued use in the double-blind, placebo-controlled phase.

In an open label study comparing bioavailability of Abilify Maintena administered in the deltoid or gluteal muscle, injection site related reactions were slightly more frequent in the deltoid muscle. The majority were mild and improved on subsequent injections. Repeated occurrence of injection site pain was more frequent in the deltoid muscle.

### ***Leukopenia***

Neutropenia has been reported in the clinical program with Abilify Maintena and typically starts around day 16 after first injection, and lasts a median of 18 days.

### ***Extrapyramidal Symptoms (EPS)***

During the double-blind phases of the 38- and 52-week trials in schizophrenia, treatment-emergent EPS and EPS-related events were reported for Abilify Maintena (18.4%) and oral aripiprazole tablets 10-30 mg (11.7%). The most commonly reported EPS and EPS-related events in each group were akathisia events (Abilify Maintena: 8.2%; oral aripiprazole tablets 10-30 mg: 6.8%; and placebo: 6.0%), followed by parkinsonism events (6.9%; 4.1% and 3.0, respectively). Akathisia typically starts around day 10 after first injection, and lasts a median of 56 days.

There was minimal variation in EPS symptoms during the double-blind phases as assessed by mean changes from baseline in the Simpson-Angus Scale (SAS), Abnormal Involuntary Movement Scale (AIMS) and Barnes Akathisia Rating Scale (BARS) rating scales. The mean changes were not considered to be clinically relevant.

During the double-blind, placebo-controlled phase of the bipolar I disorder trial, 36/132 (27.3%) Abilify Maintena-treated subjects and 22/133 (16.5%) placebo-treated subjects had treatment-emergent EPS and EPS-related AEs. In the Abilify Maintena (vs. placebo) group, EPS and EPS-related events reported by  $\geq 2\%$  of subjects were akathisia and psychomotor hyperactivity events (22.0% vs 12.8%), parkinsonism events (5.3% vs 3.8%), dyskinetic (2.3% vs 1.5%), and dystonic events (2.3% vs 0.0%).

The most frequently reported treatment-emergent EPS and EPS-related AE was akathisia with 28/132 (21.2%) Abilify Maintena-treated subjects and 17/133 (12.8%) placebo-treated subjects experiencing an event. One Abilify Maintena-treated and no placebo-treated subjects experienced an SAE of akathisia and 2 Abilify Maintena-treated subjects were discontinued due to akathisia. There were no other EPS-related SAEs and no other TEAEs leading to discontinuation reported.

### ***Dystonia***

*Class Effect:* Symptoms of dystonia, prolonged abnormal contractions of muscle groups, may occur in susceptible individuals during the first few days of treatment. Dystonic symptoms include: spasm of the neck muscles, sometimes progressing to tightness of the throat, swallowing difficulty, difficulty breathing, and/or protrusion of the tongue. While these symptoms can occur at low doses, they occur more frequently and with greater severity with high potency and at higher doses of first generation antipsychotic drugs. An elevated risk of acute dystonia is observed in males and younger age groups.

### ***Weight***

During the double-blind, active-controlled phase of the 38-week trial in schizophrenia, the incidence of weight gain of  $\geq 7\%$  from baseline to last visit was 9.5% for the Abilify Maintena group and 11.7% for the oral aripiprazole tablets 10-30 mg group. The incidence of weight loss of  $\geq 7\%$  from baseline to last visit was 10.2% for Abilify Maintena and 4.5% for oral aripiprazole tablets 10-30 mg. During double-

blind treatment, mean change in body weight from baseline to last visit was -0.2kg for Abilify Maintena, +0.7kg for oral aripiprazole tablets.

During the double-blind, placebo-controlled phase of the 52-week Abilify Maintena schizophrenia trial, the incidence of weight gain of  $\geq 7\%$  from baseline to last visit was similar between Abilify Maintena and placebo: 6.4% for the Abilify Maintena group and 5.2% for the placebo group. The incidence of weight loss of  $\geq 7\%$  from baseline to last visit was 6.4% for the Abilify Maintena group and 6.7% for the placebo group. During double-blind treatment, mean change in body weight from baseline to last visit was -0.2kg for Abilify Maintena and -0.4kg for placebo ( $p=0.812$ ).

In the 12-week clinical trial in acutely relapsed schizophrenia patients, the incidence of potentially clinically relevant weight gain of  $\geq 7\%$  from baseline to last visit was 21.5% for the Abilify Maintena group and 8.5% for the placebo group. The mean change in body weight from baseline to last visit was +2.8 kg (N=144) for Abilify Maintena and +0.8 kg (N=141) for placebo with a median exposure of 85 days.

During the double-blind, placebo-controlled phase of the bipolar I disorder trial, the incidence of weight gain  $\geq 7\%$  at any time was 18.0% in the Abilify Maintena group and 12.9% for the placebo group; the incidence of weight loss  $\geq 7\%$  at any time was 9.4% in the Abilify Maintena group and 12.1% in the placebo group. At the last visit, the incidence of potentially clinically relevant weight gain was 13.3% in the Abilify Maintena group and 12.1% in the placebo group; the incidence of weight loss  $\geq 7\%$  at last visit was 5.5% in the Abilify Maintena group and 10.6% in the placebo group. The mean change (SD) from baseline at Week 52 was 1.3 (5.9) kg for the Abilify Maintena group and 1.5 (6.1) kg for the placebo group; at the last visit, the mean change from baseline was 0.9 (5.3) kg for the Abilify Maintena group and 0.0 (5.9) kg for the placebo group.

Patients should have baseline and periodic monitoring of body weight, and other cardiometabolic parameters, including fasting glucose, full lipid profile and blood pressure, during treatment with any atypical antipsychotic including Abilify Maintena.

### ***Laboratory Tests***

No clinically relevant mean changes from baseline in serum chemistry, haematology, urinalysis or other laboratory test (e.g., insulin, fasting insulin) results were observed during the acute or long-term clinical trials with Abilify Maintena.

Comparisons between oral aripiprazole and placebo in the proportions of patients experiencing potentially clinically significant changes in routine laboratory and lipid parameters revealed no medically important differences. Elevations of CPK (creatine phosphokinase), generally transient and asymptomatic, were observed in 3.5% of patients treated with oral aripiprazole as compared to 2.0% of patients treated with placebo.

### **QT Interval**

During double-blind treatment in the long-term trials, 1/534 (0.2%) Abilify Maintena subjects had a treatment-emergent adverse event related to QT interval change (prolonged ECG QT).

### **Prolactin**

Changes in prolactin levels were comparable across trials, irrespective of indication (schizophrenia or bipolar I disorder), and there were no clinically relevant mean changes from baseline to the last visit with regard to prolactin levels during the double-blind treatment phase of either trial.

In the double-blind active-controlled phase of the 38-week trial, from baseline to last visit there was a mean decrease in prolactin levels in the Abilify Maintena group ( $-0.33$  ng/mL) compared with a mean increase in the oral aripiprazole tablets 10-30 mg group ( $0.79$  ng/mL;  $p<0.01$ ). The incidence for Abilify



Maintena patients with prolactin levels >1 time the upper limit of normal (ULN) range at any assessment was 5.4% compared with 3.5% of oral aripiprazole tablets 10-30 mg, with a higher incidence in male patients than female patients in each treatment group.

In the double-blind placebo-controlled phase of the 52-week trial in schizophrenia, from baseline to last visit, there was a mean decrease in prolactin levels in the Abilify Maintena group (-0.38 ng/mL) compared with a mean increase in the placebo group (1.67 ng/mL). The incidence of Abilify Maintena patients with prolactin levels >1 time the ULN was 1.9% compared to 7.1% for placebo patients.

Of note, differences in the mean ( $\pm$  SD) change from the double-blind treatment phase baseline to the last visit of the double-blind treatment phase in prolactin were negligible between the Abilify Maintena and placebo groups and of little if any clinical relevance, indicating no implications for adverse effects on prolactin.

In the double-blind, placebo-controlled phase of the bipolar I disorder trial, the mean changes from baseline to last visit in prolactin were minimal in the Abilify Maintena 400 mg/300 mg group (0.15 ng/mL) compared to placebo (3.00 ng/mL) and none of the changes were considered to be clinically meaningful. There were no clinically meaningful differences in the incidence of prolactin levels above the ULN between treatment groups and no incidence of  $> 3 \times$  ULN reported during the double-blind, phase of this trial. No clinically meaningful differences in shifts in prolactin between treatment groups or gender were reported. During the double-blind phase, no Abilify Maintena-treated and 0.8% of placebo-treated subjects experienced the prolactin-related TEAE of hyperprolactinaemia.

### **Lipid Parameters**

In the two double-blind studies of 38- and 52-weeks' duration, the differences in the mean change from baseline (double-blind treatment phase) to the last visit in fasting lipid parameter values (total cholesterol, triglycerides, HDL, and LDL) were negligible between the Abilify Maintena 400 mg/300 mg group compared with oral aripiprazole tablets 10-30 mg group, aripiprazole IM depot 50 mg/25 mg group or placebo groups and could be considered as of no major clinical concern.

### ***Other Adverse Reactions Observed During the Clinical Trial Evaluation of Abilify Maintena***

All reported events in the Abilify Maintena group during the randomisation phase of the clinical trials, reported by less than 2% of subjects, and at least as frequently as in the placebo group are listed below.

### **Blood and Lymphatic System Disorders**

Anaemia, bicytopenia, leukopenia, lymphadenopathy, neutropenia, thrombocytopenia.

### **Cardiac Disorders**

Acute myocardial infarction, atrial fibrillation, bradycardia, cardio-respiratory arrest, first degree atrioventricular blocks, cardiac failure congestive, ventricular extrasystoles.

### **Ear and Labyrinth disorders**

Deafness, tinnitus, vertigo.

### **Eye Disorders**

Conjunctivitis allergic, eye irritation, eye pain, eyelid ptosis, oculogyric crisis, photophobia, vision blurred.

### **Gastrointestinal disorders**

Abdominal distension, abdominal pain, abdominal pain upper, anorectal discomfort, aphthous stomatitis, colitis, constipation, dental caries, diverticulum, dry mouth, dyspepsia, dysphagia, food poisoning, frequent bowel movements, gastritis, gastroesophageal reflux disease, gingival oedema,

gingival pain, gingivitis, haemorrhoidal haemorrhage, haemorrhoids, inguinal hernia, loose tooth, nausea, oral discomfort, periodontitis, poor dental condition, salivary hypersecretion, tongue disorder, tooth impacted, tooth loss, vomiting.

### **General disorders and administration site conditions**

Asthenia, chest discomfort, gait disturbance, influenza-like illness, infusion site haematoma, infusion site swelling, injection related reaction, injection site discomfort, injection site pruritus, injection site induration, injection site pain, injection site reaction, injection site swelling, lethargy, night sweats, oedema peripheral, pain, sluggishness, suprapubic pain, swollen tongue, thirst, vessel puncture site haematoma, vessel puncture site pain.

### **Hepatobiliary disorders**

Cholecystitis chronic, cholelithiasis, hepatic cirrhosis, hepatic steatosis, hepatosplenomegaly.

### **Immune System Disorders**

Drug hypersensitivity.

### **Infections and Infestations**

Acarodermatitis, anal abscess, appendicitis perforated, breast cellulitis, chlamydial infection, cellulitis, cystitis, dermatitis, ear infection, Escherichia UTI, folliculitis, fungal infection, fungal skin infection, furuncle, gastroenteritis, gastroenteritis viral, herpes virus infection, herpes zoster, hordeolum, impetigo, laryngitis, lice infestation, localised infection, mastitis, oral candidiasis, oropharyngitis fungal, otitis externa, otitis media, pharyngitis, pharyngitis streptococcal, pilonidal cyst, pneumonia, respiratory tract infection, sycosis barbae, trichomoniasis, viral rhinitis, subcutaneous abscess, tinea pedis, tooth abscess, tooth infection, urinary tract infections, vaginal infection, varicella, viral infection, viral upper respiratory tract infection, vulvovaginal mycotic infection.

### **Injury, poisoning and procedural complications**

Accident, ankle fracture, arthropod bite, carbon monoxide poisoning, chest injury, contusion, drug toxicity, excoriation, face injury, fall, foot fracture, gunshot wound, injury, joint dislocation, joint sprain, laceration, multiple injuries, muscle injury, muscle oedema, muscle strain, procedural pain, radius fracture, skeletal injury, skin laceration, spinal column injury, thermal burn, tooth fracture, wound.

### **Investigations**

Alkaline phosphatase increased, bilirubin increased, blood creatinine phosphokinase increased, blood insulin increased, cholesterol decreased, glucose decreased, glucose increased, lactate dehydrogenase increased, triglycerides decreased, triglycerides increased, electrocardiogram abnormal, electrocardiogram QT prolonged, electrocardiogram ST segment depression, electrocardiogram T wave amplitude decreased, electrocardiogram T wave inversion, gamma-glutamyltransferase increased, glucose urine present, glycosylated haemoglobin increased, heart rate decreased, hepatic enzyme increased, liver function test abnormal, liver function test increased, neutrophil count decreased, protein urine, waist circumference increased, white blood cell count decreased, white blood cells urine.

### **Metabolism and nutrition disorders**

Appetite disorder, decreased appetite, diabetes mellitus, gout, hypercholesterolaemia, hyperglycaemia, hyperinsulinaemia, hyperlipidaemia, hypertriglyceridaemia, hyperuricaemia, hypoglycaemia, increased appetite, overweight, type 2 diabetes mellitus.

### **Musculoskeletal and connective tissue disorders**

Arthritis, joint swelling, muscle rigidity, muscle haemorrhage, muscle spasm, muscle tightness, muscle twitching, musculoskeletal pain, myalgia, neck pain, nuchal rigidity, rotator cuff syndrome, sciatica, trismus.

### **Neoplasms benign malignant and unspecified**

Basal cell carcinoma, breast fibroma, pancreatic carcinoma.

### **Nervous system disorders**

Bradykinesia, brain injury, cogwheel rigidity, disturbance in attention, drooling, dyskinesia, dystonia, extrapyramidal disorder, hypersomnia, hypoaesthesia, migraine, movement disorder, paraesthesia, parkinsonism, parosmia, poor quality sleep, post-traumatic neck syndrome, psychomotor hyperactivity, restless leg syndrome, sedation, sinus headache, syncope, tardive dyskinesia, tension headache, dizziness, transient ischaemic attack.

### **Psychiatric Disorders**

Abnormal dreams, affect lability, apathy, bruxism, bulimia nervosa, delusion, dysphemia, dysphoria, hallucination auditory, hypersexuality, hypomania, hyposomnia, initial insomnia, irritability, libido decreased, middle insomnia, mood altered, nightmare, panic attack, panic reaction, psychomotor retardation, sleep disorder, social avoidant behaviour, somnambulism, suicidal ideation, suicide attempt, tension.

### **Renal and Urinary Disorders**

Asymptomatic bacteriuria, glycosuria, hypertonic bladder, micturition urgency, nephrolithiasis, pollakiuria.

### **Reproductive system and breast disorders**

Adnexa uteri pain, breast mass, breast tenderness, ejaculation delayed, erectile dysfunction, galactorrhoea, gynaecomastia, menorrhagia, ovarian cyst, sexual dysfunction, vulvovaginal dryness.

### **Respiratory Thoracic and Mediastinal disorders**

Allergic sinusitis, acute respiratory distress syndrome, asthma, dysphonia, dyspnoea, epistaxis, nasal septum deviation, non-cardiac chest pain, oropharyngeal pain, paranasal sinus hypersecretion, respiratory failure, respiratory tract congestion, rhinalgia, rhinitis allergic, sinus congestion, wheezing, hiccups.

### **Skin and Subcutaneous tissue disorders**

Acne, blister, dermatitis contact, dry skin, eczema, erythema, hyperkeratosis, pityriasis, pruritus, psoriasis, rash, rash macula, rosacea, skin induration, skin lesion, skin striae, urticaria.

### **Social circumstances**

Poor personal hygiene.

### **Vascular Disorders**

Orthostatic hypertension.

### ***Post-Market Adverse Drug Reactions***

The following adverse reactions have been reported during post-marketing surveillance with aripiprazole. The frequency of these reactions cannot be estimated from available post-marketing data and the causal relationship to the drug cannot be definitely established in the post-marketing scenario.

**Blood and lymphatic system disorders**

Leukopenia, neutropenia, thrombocytopenia, blood prolactin decrease.

**Endocrine disorders**

Hyperglycaemia, diabetes mellitus, diabetic ketoacidosis (DKA), diabetic hyperosmolar coma.

**Metabolism and nutrition disorders**

Anorexia, hyponatraemia.

**Psychiatric disorders**

Agitation, hypersexuality, pathological gambling, impulse-control disorders, obsessive-compulsive disorder, eating disorder.

**Nervous system disorders**

Speech disorder, grand mal convulsion.

**Eye disorders**

Diplopia.

**Vascular disorders**

Syncope, hypertension.

**Respiratory, thoracic and mediastinal disorders**

Aspiration pneumonia, hiccups.

**Gastrointestinal disorders**

Pancreatitis, dysphagia, diarrhoea.

**Hepato-biliary disorders**

Jaundice, hepatitis.

**Skin and subcutaneous tissue disorders**

Allergic reaction (e.g., anaphylactic reaction, angioedema, pruritus, or urticaria, rash, laryngospasm), hyperhidrosis, alopecia, drug reaction with eosinophilia and systemic symptoms (DRESS).

**Musculoskeletal and connective tissue disorders**

Rhabdomyolysis, myalgia, musculoskeletal stiffness.

**Renal and urinary disorders**

Urinary incontinence, urinary retention.

**Reproductive system and breast disorders**

Priapism.

**General disorders and administration site conditions**

Temperature regulation disorder (e.g., hypothermia, pyrexia), chest pain.

## **Investigations**

Blood creatine phosphokinase increased, blood glucose increased, blood glucose fluctuation, glycosylated haemoglobin increased, weight increased, weight decreased, Alanine Aminotransferase increased, Aspartate Aminotransferase increased, Gamma-glutamyltransferase increased.

Although a causal relationship has not been established, cases of suicide attempt, suicidal ideation, and completed suicide, have been reported post marketing.

Undesirable effects known to be associated with antipsychotic medication which have also been reported in association with aripiprazole are Neuroleptic Malignant Syndrome, tardive dyskinesia and seizure.

Uncommon occurrences of depression and tachycardia have also been reported in association with aripiprazole.

## **4.9 Overdose**

For risk assessment and advice on the management of overdose, contact the New Zealand Poisons Centre (Tel: 0800 POISON or 0800 764 766 for New Zealand) or the Poison Information Centre (Tel: 13 11 26 for Australia).

No cases of overdose associated with adverse reactions were reported in clinical studies with Abilify Maintena. Care must be taken to avoid inadvertent injection of this medicinal product into a blood vessel. Following any confirmed or suspected accidental overdose/inadvertent intravenous administration, close observation of the patient is needed and if any potentially medically serious sign or symptom develops, monitoring, which should include continuous electrocardiographic monitoring, is required. The medical supervision and monitoring should continue until the patient recovers.

A simulation of dose dumping showed that the predicted median aripiprazole concentration reaches a peak of 4,500 ng/mL which corresponds to approximately 9 times the upper therapeutic range. In case of dose dumping, aripiprazole concentrations are predicted to descend rapidly to the upper limit of the therapeutic window after approximately 3 days. By the 7th day, the median aripiprazole concentrations further decline to concentrations following an IM depot dose with no dose dumping. While overdose is less likely with parenteral than oral medicinal products, reference information for oral aripiprazole overdose is presented below.

### ***Signs and Symptoms***

In clinical trials and post-marketing experience, accidental or intentional acute overdose of aripiprazole alone was identified in adult patients with reported estimated doses up to 1,260 mg (42 times higher than the recommended daily aripiprazole dose, 30 mg) with no fatalities. The potentially medically important signs and symptoms observed included lethargy, increased blood pressure, somnolence, tachycardia, nausea, vomiting and diarrhoea. In addition, reports of accidental overdose with aripiprazole alone (up to 195 mg) in children have been received with no fatalities. The potentially medically serious signs and symptoms reported included somnolence, transient loss of consciousness and extrapyramidal symptoms.

### ***Management of overdose***

Management of overdose should concentrate on supportive therapy, maintaining an adequate airway, oxygenation and ventilation, and management of symptoms. The possibility of multiple medicinal product involvement should be considered. Therefore, cardiovascular monitoring should be started immediately and should include continuous electrocardiographic monitoring to detect possible arrhythmias. Following any confirmed or suspected overdose with aripiprazole, close medical supervision and monitoring should continue until the patient recovers.

### ***Haemodialysis***

Although there is no information on the effect of haemodialysis in treating an overdose with aripiprazole, haemodialysis is unlikely to be useful in overdose management, since aripiprazole is highly bound to plasma proteins.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

#### ***Mechanism of action***

The mechanism of action of aripiprazole, as well as other drugs having efficacy in schizophrenia and bipolar disorder, is unknown. It has been proposed that aripiprazole's efficacy is mediated through a combination of partial agonism at dopamine D<sub>2</sub> and serotonin 5-HT<sub>1A</sub> receptors and antagonism at serotonin 5-HT<sub>2A</sub> receptors.

Abilify Maintena activity is primarily due to the parent drug, aripiprazole. Aripiprazole exhibited antagonist properties in animal models of dopaminergic hyperactivity and agonist properties in animal models of dopaminergic hypoactivity. Aripiprazole exhibited high binding affinity *in vitro* for dopamine D<sub>2</sub> and D<sub>3</sub>, serotonin 5-HT<sub>1A</sub> and 5-HT<sub>2A</sub> receptors (K<sub>i</sub> values of 0.3, 0.8, 1.7, and 3.4 nM, respectively), and moderate affinity for dopamine D<sub>4</sub>, serotonin 5-HT<sub>2C</sub> and 5-HT<sub>7</sub>,  $\alpha$ 1-adrenergic, and histamine H<sub>1</sub> receptors (K<sub>i</sub> values of 44, 15, 39, 57, and 61 nM, respectively). Aripiprazole also exhibited moderate binding affinity for the serotonin reuptake site (K<sub>i</sub> value of 98 nM) but no appreciable affinity for muscarinic receptors (IC<sub>50</sub>>1000 nM).

The predominant metabolite in human plasma, dehydro-aripiprazole, has been shown to have a similar affinity for dopamine D<sub>2</sub> and D<sub>3</sub> receptors (K<sub>i</sub> values 0.4 and 0.5 nM, respectively) as the parent compound and a much lower affinity for the other receptor subtypes (serotonergic, noradrenergic and histaminergic receptors).

Interaction with receptors other than dopamine and serotonin subtypes may explain some of the other clinical effects of aripiprazole.

Aripiprazole oral doses ranging from 0.5 to 30 mg administered once a day to healthy subjects for 2 weeks produced a dose-dependent reduction in the binding of [<sup>11</sup>C]-raclopride, a D<sub>2</sub> receptor ligand, to the caudate and putamen detected by positron emission tomography.

#### ***Clinical trials***

##### ***Schizophrenia***

The efficacy and safety of Abilify Maintena in the treatment of adult patients with schizophrenia was established in one pivotal short-term, randomised, double-blind, placebo-controlled trial in acutely relapsed patients and one pivotal long-term, randomised, double-blind, placebo-controlled trial.

##### **Clinical Efficacy in the Acute Phase of Schizophrenia**

The efficacy of Abilify Maintena in adult patients in the acute phase of schizophrenia was established in one short-term (12 weeks), randomised, double-blind, placebo-controlled trial. Patients included in this trial met DSM-IV-TR criteria for schizophrenia and must have experienced an acute psychotic episode as defined by both a PANSS total score  $\geq 80$  and a PANSS score  $>4$  on each of four specific psychotic symptoms (conceptual disorganisation, hallucinatory behaviour, suspiciousness/persecution, unusual thought content) at baseline. Patients experiencing their first psychotic episode and those considered treatment resistant were excluded. Patients had a mean PANSS Total Score of 103 (range 82 to 144) and a CGIS score of 5.2 (markedly to severely ill) at entry.

In this study patients were administered Abilify Maintena (n=167) or IM placebo (n=172) on Days 0, 28, and 56. The dose could be adjusted down and up within the range of 400 mg to 300 mg on a one-time basis. Patients who had not taken aripiprazole previously had tolerability with oral aripiprazole (10 mg daily for 3 days) established prior to initiating treatment with Abilify Maintena or placebo. Patients randomised to Abilify Maintena also received concomitant oral aripiprazole, 10 to 20 mg/day, for the first two weeks of the study.

In the Abilify Maintena group, for 96.4% of patients, there was no difference between the starting dose and ending dose of Abilify Maintena (400 mg).

The primary endpoint was the change from baseline to Week 10 in PANSS Total Score. Abilify Maintena was superior to placebo in improving the PANSS total score, with Week 10 scores of -26.8 and -11.7, respectively (see Error! Reference source not found.7). A statistically significant difference ( $p \leq 0.0001$ ) was seen at each measured time point beginning at Week 1 and continuing through to study completion. The adjusted mean change in PANSS Total Score over time is shown in **Figure 1**


**Table 7 - Change from Baseline in PANSS Total Score at Week 10 in Acute Phase Schizophrenia**

Treatment Group	Primary Efficacy Measure: PANSS Total Score <sup>a</sup>		
	Mean Baseline Score (SD)	LS Mean Change from Baseline (SE)	Treatment Difference (95% CI)
Abilify Maintena 400 mg/300 mg	102.4 (11.4) N=162	-26.8 (1.6) N=99	-15.1 (-19.4, -10.8) $p < 0.0001$
Placebo	103.4 (11.1) N=167	-11.7 (1.6) N=81	--

SD: standard deviation; SE: standard error; LS Mean: least-squares mean; CI: confidence interval.

<sup>a</sup>Data were analysed using a mixed model for repeated measurements (MMRM) approach. The analysis included only patients who were randomly assigned to treatment, given at least one injection, had baseline and at least one post-baseline efficacy assessment.

<sup>b</sup>ifference (Abilify Maintena minus placebo) in LS mean change from baseline.




**Figure 1-Adjusted Mean Change from Baseline in PANSS Total Score (MMRM)**

For the key secondary endpoint, the change from baseline to Week 10 in CGI-S score, the treatment difference between the Abilify Maintena group (LS mean change -1.4) and the placebo group (LS mean change -0.6) was -0.8 (95% CI: -1.1, -0.6), which was statistically significant ( $p < 0.0001$ ).

Response was defined as a  $\geq 30\%$  reduction from baseline in PANSS total score. The responder rate was numerically higher in the Abilify Maintena group at all post-baseline time points; the treatment differences were statistically significant ( $p \leq 0.0013$ ) from Week 8 to Week 12 (see **Figure 2**). At Week 10, the responder rate was 37.0% in the Abilify Maintena group compared to 14.4% in the placebo group; the treatment difference was 22.7% (95% CI 12.9%, 32.4%).





**Figure 2 -Responder Rate in the Acute Phase Schizophrenia Study**

#### ***Clinical Efficacy in the Maintenance Phase of Schizophrenia***

The pivotal trial was a 52-week, randomised, double-blind, placebo-controlled trial conducted in adult patients with a current diagnosis of schizophrenia. This trial consisted of a screening phase and four treatment phases: Conversion, Oral Stabilisation, Abilify Maintena Stabilisation, and Double-blind Placebo-controlled. Patients eligible for the double-blind, placebo-controlled phase were randomly assigned in a 2:1 ratio to double-blind treatment with Abilify Maintena 400 mg (with an option to decrease to 300 mg for tolerability reasons) or placebo, respectively. The trial was completed early as a consequence of the positive pre-specified interim analysis and therefore only 26 patients completed 52 weeks of treatment. Eighty-seven per cent (87%) of subjects randomised to the 400 mg dose remained on this dose until either completing the trial duration or withdrawing from the trial.


The final efficacy analysis included 403 randomised patients and 80 exacerbations of psychotic symptoms/impending relapse events. The trial was terminated early because efficacy was demonstrated by the pre-specified interim analysis. The hazard ratio from the Cox proportional hazard model for the placebo to Abilify Maintena comparison was 5.029 (95% CI: 3.154, 8.018), thus patients in the placebo group had a 5-fold greater risk of experiencing impending relapse than patients in the Abilify Maintena group. The trial results support the efficacy for Abilify Maintena over 52 weeks of treatment.

The Kaplan-Meier curves of the time from randomisation to impending relapse during the 52-week, double-blind treatment phase for Abilify Maintena and placebo groups are shown in Figure 3.

The percentage of patients meeting the impending relapse criteria was significantly lower ( $p < 0.0001$ ) in the Abilify Maintena group (10.0%) than in the placebo group (39.6%).

The time to impending relapse was significantly shorter ( $p < 0.0001$ ) for subjects in the placebo group compared with subjects in the Abilify Maintena group.

**Figure 3 Kaplan-Meier Product Limit Plot of Time to Impending Relapse (Double-blind, Placebo-controlled Phase Efficacy Sample)**



IM = Intramuscular

Further, the superiority of Abilify Maintena compared to placebo is supported by the results of the analysis of PANSS total score, PANSS Positive and Negative Subscales, CGI-S, CGI-I. During the double-blind phase, significant differences in mean PANSS total and CGI-S scores were observed.

### ***Bipolar I Disorder***


#### ***Clinical Efficacy in Prevention of Recurrence of Manic or Mixed Episodes of Bipolar I Disorder***

The efficacy and safety of Abilify Maintena as treatment in adults aged 18 to 65 years was demonstrated in a 52-week, double-blind, placebo-controlled randomized withdrawal trial in patients who met DSM-IV-TR criteria for bipolar I disorder and who were experiencing a manic episode at trial entry. This trial consisted of a screening phase and 4 treatment phases: oral conversion, oral stabilization, Abilify Maintena stabilization, and a randomized, double-blind, placebo-controlled withdrawal phase.

Patients currently receiving oral treatment for their bipolar I disorder with medications other than aripiprazole monotherapy entered the oral conversion phase. In this phase, patients discontinued other treatments, such as mood stabilizers, antidepressants, or other antipsychotics over a period of 4 to 6 weeks and converted to aripiprazole monotherapy. Patients already treated with oral aripiprazole monotherapy at the time of trial entry and those subjects converted to oral aripiprazole monotherapy in the oral conversion phase proceeded to the oral stabilization phase. Patients fulfilling the stabilization requirement were assigned to receive, in a single-blind fashion, Abilify Maintena 400 mg and began an IM depot stabilization phase for a minimum of 12 weeks and a maximum of 28 weeks. Patients who demonstrated stability for 8 consecutive weeks were randomized into the 52-week double-blind,

placebo-controlled treatment phase. Of the 731 subjects who entered the trial, 466 subjects entered the conversion phase, 632 subjects entered the oral stabilization phase (including 265 subjects who entered the oral stabilization phase directly), and 425 subjects entered the IM depot stabilization phase. Stability was defined as: having achieved or maintained outpatient status; Young-Mania Rating Scale (YMRS) total score  $\leq 12$ ; Montgomery Asberg Depression Rating Scale (MADRS) total score  $\leq 12$  and no active suicidality; with active suicidality defined as a score of 4 or more on the MADRS Item 10 or an answer of “yes” on Question 4 or 5 on the Columbia Suicide Severity Rating Scale (C-SSRS).

The final efficacy analysis included 265 patients and the primary efficacy endpoint is shown in the Kaplan-Meier curves of the cumulative proportion of patients with recurrence of any mood episode (manic, mixed or depressive) during the 52-week, double-blind treatment phase for Abilify Maintena and placebo treatment groups (**Figure 4**). A total of 103 patients with recurrence of any mood episode were observed during the double-blind treatment phase: 35 occurred during Abilify Maintena treatment and 68 occurred during placebo treatment. The hazard ratio from the Cox proportional hazard model for the placebo to Abilify Maintena comparison was 2.220 (95% CI: 1.475, 3.340), thus patients receiving placebo had a 2.2-fold greater risk of experiencing a recurrence of a mood episode than patients receiving Abilify Maintena. These results support efficacy for Abilify Maintena over 52 weeks of treatment.



ARIP IMD= Aripiprazole IM depot

**Figure 4 Kaplan-Meier Curves of Time to Recurrence for Any Mood Episode**

For the key secondary endpoint, the percentage of patients meeting the criteria for recurrence of any mood episode (manic, mixed, depressive) was significantly lower (Fisher’s exact test  $p < 0.0001$ ) in the Abilify Maintena group (26.5%) compared with placebo (51.1%). The times to recurrence of manic, mixed, and mixed/manic mood episodes were statistically significantly delayed in the aripiprazole IM depot group compared with the placebo group, with the log-rank test  $p$ -values of  $< 0.0001$ , 0.0237, and

< 0.0001, respectively. The time to recurrence of a depressive mood episode showed no difference between the aripiprazole IM depot and placebo groups (p = 0.8247) (Table 8)

**Table 8 Analysis of Time to Recurrence of a Manic, Depressive, or Mixed Mood Episode (Double-blind, Placebo-controlled Phase Efficacy Sample)**

Mood Episode	Number of Subjects Treated	Number of Subjects With Recurrence	Recurrence Rate (%)	Median Time to Recurrence (Days)	Hazard Ratio <sup>a</sup>	95% CI	P-value <sup>b</sup>
<b>Manic</b>							
Aripiprazole IM Depot	132	12	9.1	NE	0.259 <sup>c</sup>	(0.136, 0.495)	<0.0001
Placebo	133	40	30.1	377	3.856 <sup>d</sup>	(2.020, 7.358)	
<b>Depressive</b>							
Aripiprazole IM Depot	132	20	15.2	NE	0.932 <sup>c</sup>	(0.497, 1.747)	0.8247
Placebo	133	19	14.3	NE	1.073 <sup>d</sup>	(0.572, 2.013)	
<b>Mixed<sup>e</sup></b>							
Aripiprazole IM Depot	132	2	1.5	NE	0.202 <sup>c</sup>	(0.044, 0.939)	0.0237
Placebo	133	9	6.8	NE	4.939 <sup>d</sup>	(1.065, 22.904)	
<b>Mixed/Manic<sup>e</sup></b>							
Aripiprazole IM Depot	132	14	10.6	NE	0.249 <sup>c</sup>	(0.137, 0.451)	<0.0001
Placebo	133	49	36.8	377	4.017 <sup>d</sup>	(2.216, 7.282)	

<sup>a</sup>The HR was derived from the Cox proportional hazard model with treatment as term.

<sup>b</sup>The p-value was derived from the log-rank test.

<sup>c</sup>HR < 1 is in favor of aripiprazole IM depot 400/300 mg group.

<sup>d</sup>HR > 1 is in favor of aripiprazole IM depot 400/300 mg group.

<sup>e</sup>Time to recurrence of a mixed mood or a mixed/manic mood episode are exploratory endpoints.

## 5.2 Pharmacokinetic properties

### *Absorption*

Aripiprazole absorption into the systemic circulation is slow and prolonged following Abilify Maintena administration due to the low solubility of aripiprazole particles.

Following a single intramuscular dose of Abilify Maintena in the deltoid or gluteal muscle, the extent of absorption (AUC) is comparable for both injection sites, but the rate of absorption (C<sub>max</sub>) was higher following administration to the deltoid muscle.

Following multiple intramuscular doses, the plasma concentrations of aripiprazole gradually rise to a maximum plasma concentration at median  $t_{\max}$  of 4 days for the deltoid muscle and 7 days for the gluteal muscle.

Dose-proportional increases in aripiprazole and dehydro-aripiprazole concentrations and AUC parameters are observed after monthly Abilify Maintena injections of 400 mg and 300 mg and steady state aripiprazole plasma concentrations were attained by the fourth monthly injection for both sites of administration.

### ***Distribution***

Based on results from trials with oral administration of aripiprazole, aripiprazole is widely distributed throughout the body with an apparent volume of distribution of 4.9 L/kg, indicating extensive extravascular distribution. At therapeutic concentrations, aripiprazole is highly bound (88-97% to >99%, as determined by polydimethylsiloxane-glass bead and equilibrium dialysis assays, respectively) to serum proteins, primarily albumin, *in vitro*. Aripiprazole did not alter the pharmacokinetics and pharmacodynamics of highly protein-bound warfarin, suggesting that protein displacement of warfarin did not occur.

### ***Metabolism***

Aripiprazole undergoes minimal pre-systemic metabolism at the site of injection. Aripiprazole is extensively metabolised by the liver primarily by three biotransformation pathways: dehydrogenation, hydroxylation and N-dealkylation. Based on *in vitro* studies, CYP3A4 and CYP2D6 enzymes are primarily responsible for dehydrogenation and hydroxylation of aripiprazole, and N-dealkylation is catalysed by CYP3A4. Aripiprazole is the predominant drug moiety in systemic circulation. After multiple dose administration of Abilify Maintena, dehydro-aripiprazole, the active metabolite represents about 29.1-32.5% of aripiprazole AUC in plasma.

### ***Excretion***

After administration of multiple doses of 400 mg or 300 mg of Abilify Maintena, the observed mean aripiprazole terminal elimination half-life is 46.5 and 29.9 days, respectively. Following a single oral dose of [<sup>14</sup>C]-labelled aripiprazole, approximately 27% of the administered radioactivity was recovered in the urine and approximately 60% in the faeces. Less than 1% of unchanged oral aripiprazole was excreted in the urine and approximately 18% was recovered unchanged in the faeces.

### ***Special Populations***

#### ***CYP2D6 Poor Metabolisers***

Based on population pharmacokinetic evaluation of Abilify Maintena, the total body clearance of aripiprazole was 3.71 L/h in extensive metabolisers of CYP2D6 (EMs) and approximately 1.88 L/h (approximately 50% lower) in poor metabolisers of CYP2D6 (PMs). For dose recommendations see 4.2 Dose and method of administration, Dosage in Special Population – Known CYP2D6 Poor Metabolisers. Subjects were entered into clinical studies without knowledge of their metaboliser status and, therefore, the safety profile reflects experience in both EMs and PMs.

#### ***Elderly Patients***

After oral administration of aripiprazole, there are no clinically relevant differences in the pharmacokinetics of aripiprazole between healthy elderly and younger adults. Similarly, there was no detectable effect of age (18-61 year age range) in a population pharmacokinetic analysis of Abilify Maintena clinical trials in patients with schizophrenia.

### ***Gender***

After oral administration of aripiprazole, there are no differences in the pharmacokinetics of aripiprazole between healthy male and female subjects when differences in body weight are considered. Population pharmacokinetic analysis of Abilify Maintena revealed a difference in the predicted mean half-lives between men (24 days) and women (32 days) as well as a gender dependent absorption rate. At steady state (model predicted) however, the parameters of  $C_{min}$ ,  $C_{max}$ , and  $AUC_{0-\tau}$  did not exhibit any trends towards gender.

### ***Race***

Population pharmacokinetic evaluation of Abilify Maintena showed no evidence of race-related differences in the pharmacokinetics of aripiprazole.

### ***Renal Impairment***

In a single-dose study with oral administration of aripiprazole, the pharmacokinetic characteristics of aripiprazole and dehydro-aripiprazole were found to be similar in patients with severe renal disease compared to those in young healthy subjects.

In patients with severe renal impairment (creatinine clearance  $<30$  mL/min),  $C_{max}$  of aripiprazole (given in a single dose of 15 mg) and dehydro-aripiprazole increased by 36% and 53%, respectively, but AUC was 15% lower for aripiprazole and 7% higher for dehydro-aripiprazole.

### ***Hepatic Impairment***

A single-dose study with oral administration of aripiprazole to subjects with varying degrees of liver cirrhosis (Child-Pugh Classes A, B, and C) did not reveal a significant effect of hepatic impairment on the pharmacokinetics of aripiprazole and dehydro-aripiprazole. The AUC of aripiprazole, compared to healthy subjects, increased 31% in mild hepatic impairment, increased 8% in moderate hepatic impairment, and decreased 20% in severe hepatic impairment. None of these differences would require dose adjustment, but the study included only three patients with Class C liver cirrhosis, which is insufficient to draw conclusions on their metabolic capacity.

### ***Smoking***

Population pharmacokinetic evaluation of oral aripiprazole has revealed no evidence of clinically relevant effects from smoking on the pharmacokinetics of aripiprazole.

## **5.3 Preclinical safety data**

### ***Animal Toxicology***

The toxicological profile for aripiprazole administered to experimental animals by intramuscular injection is generally similar to that seen following oral administration at comparable plasma levels of the drug. With intramuscular injection, however, injection-site tissue reactions are observed that consist of localised inflammation, swelling, scabbing and foreign-body reactions to deposited drug. These effects gradually resolved with discontinuation of dosing.

Choleliths observed in the bile of monkeys given aripiprazole orally at doses of 25 to 125 mg/kg/day for 4 to 52 weeks (1-3 times the oral MRHD of 30 mg/day based on plasma AUC and 15-76 times the oral MRHD based on  $mg/m^2$ ) have been attributed to precipitation of sulfate conjugates of hydroxy metabolites, which exceeded their solubility limits in bile. Human biliary concentrations of these sulfate conjugates after repeated daily administration of the oral MRHD are substantially lower (0.2-14% of their in vitro solubility limits).

Bilateral retinal degeneration was observed in albino rats given oral aripiprazole for 6 months or 2 years at exposures of 6-13 times the clinical exposure at the oral MRHD of 30 mg/day (based on plasma

AUC). The exposure at the no-effect dose was 3 times that at the MRHD. A subsequent 18-month study reported this finding in albino but not pigmented rats, possibly due to lack of photoprotective ocular melanin in the albino rats, although it is unknown whether pigmentation prevented or merely delayed retinal degeneration in the pigmented rats. The clinical relevance of this finding is uncertain.

### ***Carcinogenicity***

Lifetime carcinogenicity studies were conducted in ICR mice and in Sprague-Dawley (SD) and Fischer (F344) rats. Aripiprazole was administered for 2 years in the diet at doses of 1, 3, 10, and 30 mg/kg/day to ICR mice and 1, 3, and 10 mg/kg/day to F344 rats (0.2 to 5 and 0.3 to 3 times the oral maximum recommended human dose [MRHD] of 30 mg/day based on mg/m<sup>2</sup>, respectively). SD rats were dosed orally by gavage for 2 years at 10, 20, 40, and 60 mg/kg/day (3 to 18 times the MRHD based on mg/m<sup>2</sup>). There was no evidence of tumorigenesis in male mice. In female mice, the incidences of pituitary gland adenomas and mammary gland adenocarcinomas and adenocanthomas were increased at dietary doses of 3 to 30 mg/kg/day (0.1 to 0.9 times the MRHD based on AUC and 0.5 to 5 times the MRHD based on mg/m<sup>2</sup>). In female rats, the incidence of mammary gland fibroadenomas was increased at a dietary dose of 10 mg/kg/day (<0.1 times the MRHD based on AUC and 3 times the MRHD based on mg/m<sup>2</sup>); and the incidences of adrenocortical carcinomas and combined adrenocortical adenomas/carcinomas were increased at an oral gavage dose of 60 mg/kg/day (10 times the MRHD based on AUC and 18 times the MRHD based on mg/m<sup>2</sup>). In male rats, the incidences of benign and combined benign/malignant pheochromocytomas were also increased at an oral gavage dose of 60 mg/kg/day (10 times the MRHD based on AUC and 18 times the MRHD based on mg/m<sup>2</sup>).

Proliferative changes in the pituitary and mammary gland of rodents have been observed following chronic administration of other antipsychotic agents and are considered prolactin mediated. Serum prolactin was not measured in the aripiprazole carcinogenicity studies. At the doses associated with mammary and/or pituitary tumours, hyperprolactinaemia was observed in female mice in a 13-week dietary study but not in female rats in 4- and 13-week dietary studies. Hyperprolactinaemia was observed in female rats after 5 and 13 weeks of oral administration at doses up to that associated with adrenocortical tumours, but serum prolactin was decreased at this dose in male rats. The relationship between tumourigenic findings with aripiprazole and prolactin is unclear and the relevance for human risk of prolactin-mediated endocrine tumours is unknown. The adrenocortical response in female rats is considered a consequence of increased adrenocortical cell proliferation secondary to chronic drug-related adrenocortical cytotoxicity; the no-effect exposure (plasma AUC) was about 7 times clinical exposure at the MRHD.

### ***Genotoxicity***

Aripiprazole was tested for genotoxic potential in a standard range of assays for gene mutation, chromosomal damage, and DNA damage and repair. Aripiprazole was nongenotoxic in the *in vitro* bacterial reverse mutation assay, *in vitro* forward gene mutation assay in mouse lymphoma cells, *in vitro* bacterial DNA repair assay, and the unscheduled DNA synthesis assay in rat hepatocytes. However, aripiprazole and its minor metabolite 2,3-DCPP were clastogenic in the *in vitro* chromosomal aberration assay in Chinese hamster lung cells with and without metabolic activation. A positive response in 1 of 6 *in vivo* mouse micronucleus tests was likely due to a mechanism not relevant to humans.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Carmellose sodium, mannitol, monobasic sodium phosphate monohydrate, sodium hydroxide and water for injections.

## **6.2 Incompatibilities**

In the absence of compatibility studies, this medicine must not be mixed with other medicines. Refer to 4.5 Interaction with other medicines and other forms of interaction.

## **6.3 Shelf life**

36 months from date of manufacture stored at or below 30°C.

## **6.4 Special precautions for storage**

Lyophilised Abilify Maintena is stable up to the expiration date stamped on the vial.

Store in the original packaging below 30°C. Do not freeze.

For single use in only one patient. Discard any unused solution.

The reconstituted suspension should be used immediately, but may be stored below 25°C for up to 4 hours. Shake the vial vigorously for at least 60 seconds to re-suspend prior to injection.

## **6.5 Nature and contents of container**

Abilify Maintena is presented as a sterile, single-dose, lyophilised powder for reconstitution with water for injections to give a prolonged-release suspension for injection to deliver 400 mg of aripiprazole in 400mg/vial strength or 300 mg of aripiprazole in 300-mg/vial strength. Abilify Maintena is intended for intramuscular injection only.

Abilify Maintena powder and solvent for prolonged-release suspension for injection is available as a one-month therapeutic kit. Each therapeutic kit contains one vial of lyophilised powder (400 mg or 300 mg), one vial of water for injections, one 3 mL sterile syringe with a 21 gauge needle for reconstitution, one 3 mL sterile syringe without a needle, one 1-inch (25 mm) 23 gauge sterile safety needle for injection, one 1.5-inch (38 mm) 22 gauge sterile safety needle for injection and one 2-inch (51 mm) 21 gauge sterile safety needle for injection and one vial adapter.

## **6.6 Special precautions for disposal**

Any unused medicine or waste material should be disposed of in accordance with local requirements.

# **7 MEDICINE SCHEDULE**

Prescription medicine.

# **8 SPONSOR**

Pharmacy retailing t/a  
Healthcare Logistics  
58 Richard Pearse Drive  
Airport Oaks  
Mangere  
Auckland  
New Zealand

Telephone: +64 9 918 5100

Abilify Maintena is a registered trademark of Otsuka Pharmaceutical Company.



## 9 DATE OF FIRST APPROVAL

20 August 2015

## 10 DATE OF REVISION OF THE TEXT

30 April 2025

## SUMMARY TABLE OF CHANGES

Section Changed	Summary of new information
4.1, 4.4, 4.8 and 5.1	Extension of indication to include Acute Schizophrenia and Bipolar I disorder; deletion of information regarding the potential for cognitive and motor impairment from Section 4.4; editorial changes to Section 5.1 to align with the Australian PI.
4.2	Addition of deltoid administration site.
4.9	Added risk assessment wording to overdose information.