

Ocrevus[®]

Ocrevus intravenous (IV): ocrelizumab 300 mg in 10 mL concentrate for solution for infusion

Ocrevus subcutaneous (SC): ocrelizumab 920 mg in 23 mL solution for subcutaneous injection

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using Ocrevus?

Ocrevus contains the active ingredient ocrelizumab. Ocrevus is used to treat relapsing forms of multiple sclerosis (RMS) and primary progressive multiple sclerosis (PPMS). Ocrevus decreases the number of flare-ups (relapses) and slows the worsening of RMS. Ocrevus also slows the worsening of PPMS. For more information, see Section [1. Why am I using Ocrevus?](#) in the full CMI.

2. What should I know before I use Ocrevus?

Do not use if you have ever had an allergic reaction to Ocrevus or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding. For more information, see Section [2. What should I know before I use Ocrevus?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with Ocrevus and affect how it works or Ocrevus may affect how other medicines work. A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use Ocrevus?

Your doctor or nurse will administer your Ocrevus dose as an intravenous infusion, or as a subcutaneous injection. More information can be found in Section [4. How do I use Ocrevus?](#) in the full CMI.

5. What should I know while using Ocrevus?

Things you should do	<ul style="list-style-type: none">• Tell your doctor or nurse if you have any signs and symptoms of infection; any signs or symptoms of infusion or injection related reaction or allergic reaction; any stomach or bowel problems; or any signs or symptoms of a liver problem.• Tell your doctor if you plan to receive or have received a vaccine.• Avoid the use of Ocrevus during the second or third trimester of pregnancy unless advised otherwise by your doctor.• Avoid breastfeeding for the first few days after birth while you are using Ocrevus.• Tell your doctor immediately if your condition worsens, or if you, your partner or caregiver notice any new symptoms like changes in movement or behaviour.
Things you should not do	<ul style="list-style-type: none">• Do not stop using this medicine without talking to your doctor first.
Driving or using machines	<ul style="list-style-type: none">• Be careful before you drive or use any machines or tools until you know how Ocrevus affects you.
Looking after your medicine	<ul style="list-style-type: none">• It is unlikely that you will have to store Ocrevus at home. If necessary, refrigerate between 2°C and 8°C, do not freeze.• Keep the vial in the carton to protect it from light and keep it out of reach of children.

For more information, see Section [5. What should I know while using Ocrevus?](#) in the full CMI.

6. Are there any side effects?

Common side effects include infusion related reactions or injection reactions and some types of infections, which can be serious or less serious. Allergic reactions (swelling of the face, lips, tongue or throat, trouble breathing, chest tightness, wheezing, coughing, and asthma that becomes worse) are serious side effects. Stomach and bowel problems (diarrhoea that does not go away, stomach pain, or blood in your stools) or liver problems (new or worsening fatigue, loss of appetite, nausea, vomiting, pain in the right part of the abdomen, dark-colored urine, or yellowing of your skin or eyes) could be a serious side effect.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

Ocrevus[®] (pronounced oak-rev-us)

Active ingredient: *ocrelizumab*

Consumer Medicine Information (CMI)

This leaflet provides important information about using Ocrevus. Please read this leaflet carefully before you start using Ocrevus.

All medicines have risks and benefits. Your doctor has weighed the risks of you being given Ocrevus against the benefits he/she expects it will have for you. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using Ocrevus.**

Keep this leaflet with the medicine. You may need to read it again.

Where to find information in this leaflet:

- [1. Why am I using Ocrevus?](#)
- [2. What should I know before I use Ocrevus?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use Ocrevus?](#)
- [5. What should I know while using Ocrevus?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I using Ocrevus?

Ocrevus contains the active ingredient ocrelizumab.

Ocrevus is a monoclonal antibody. Antibodies are proteins, produced naturally in the body, which help the immune system by specifically recognising and binding to proteins found on target cells. Monoclonal antibodies are designed and manufactured to work like natural antibodies so that they can target specific proteins involved in disease.

Ocrevus is used to treat relapsing forms of multiple sclerosis (RMS) and primary progressive multiple sclerosis (PPMS). Ocrevus decreases the number of flare-ups (relapses) and slows the worsening of RMS. Ocrevus also slows the worsening of PPMS.

In multiple sclerosis (MS) your immune system mistakenly attacks the protective layer (myelin) around your nerve cells. This causes inflammation and damage which stops your nervous system working properly. Ocrevus works on cells of the immune system called B-cells, which play a role in multiple sclerosis. By targeting specific B cells, Ocrevus may reduce inflammation and attacks to the nervous system. Ocrevus can decrease the number of flare-ups (relapses) and slows the progression of your disease.

Ask your doctor if you have any questions about why Ocrevus has been prescribed for you. Your doctor may have prescribed Ocrevus for another reason.

This medicine is available only with a doctor's prescription.

2. What should I know before I use Ocrevus?

Warnings

Do not use Ocrevus if:

- you have had an allergic reaction to ocrelizumab, or any of the ingredients listed at the end of this leaflet. Some of the symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing, swelling of the face, lips, tongue or other parts of the body, rash and/or itching or hives on the skin.
- Always check the ingredients to make sure you can use this medicine.

Check with your doctor if you:

- have any signs and symptoms of infection, or a history of a recurring or long-term infection such as hepatitis B.
- If you are taking or have taken medicines which affect your immune system, you may have an increased risk of infections. You should not have your treatment until the infection is resolved. There have been reports of a rare, serious brain infection called PML (progressive multifocal leucoencephalopathy) in patients receiving medicines for MS. PML can occur in patients taking Ocrevus and mostly associated with risk factors, such as age and pre-existing use of immunosuppressants. PML can cause severe disability or even death. Symptoms of PML can be similar to those of MS.
- you or your child intend to have or have had immunisation with any vaccine. Some vaccines should not be given at the same time as Ocrevus, or in the months after you receive Ocrevus. Your doctor will check if you should have any vaccines before you receive Ocrevus. There may be changes in the blood cells (B-cells) in babies whose mothers were exposed to Ocrevus during pregnancy and it may be recommended that the babies' vaccination schedule is amended/delayed.
- are allergic to any other medicines or any other substances such as foods, preservatives or dyes.
- have any other medical conditions.
- take any medicines for any other condition.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy

Tell your doctor if you are pregnant or intend to become pregnant.

You should avoid using Ocrevus during the second and third trimester of pregnancy, unless advised otherwise by your doctor.

Breastfeeding

Talk to your doctor if you are breastfeeding or plan to breastfeed.

You should avoid breastfeeding for the first few days after birth while you are using Ocrevus.

If you have not told your doctor about any of the above, tell them before you start taking Ocrevus.

Children and Adolescents

The safety and effectiveness of Ocrevus in children and adolescents under 18 years of age have not been established.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

These medicines increase your risk of side effects with Ocrevus:

- other treatments (including those for MS, or to treat other conditions) which could affect your immune system and so could affect your ability to fight infections. If you are currently using a different MS treatment or if you have used another MS treatment in the past, your doctor may ask you to stop the other medicine in advance of starting treatment with Ocrevus.
- medicines that lower blood pressure. Ocrevus may cause a temporary drop in your blood pressure. Your doctor may advise you to temporarily stop taking your blood pressure medicine before each dose.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect Ocrevus.

4. How do I use Ocrevus?

Follow all directions given to you by your doctor or nurse carefully. They may differ from the information contained in this leaflet.

How much to use

Ocrevus intravenous (IV)

Ocrevus IV is not intended for subcutaneous administration. Ocrevus IV must be prepared by a healthcare professional and will be given in a hospital or clinic by a doctor or nurse. You will have Ocrevus as an intravenous infusion. It will go slowly into your vein.

The amount of Ocrevus you are given and how long each infusion lasts are different for the first and following doses. Before you receive Ocrevus, your doctor or nurse will give you other medicines to help reduce possible side effects such as infusion-related reactions.

- The dose is 600 mg every 6 months.
- First two infusions: your first 600 mg will be given as 2 separate intravenous infusions (300 mg each), given 2 weeks apart by a doctor or nurse. Each infusion will take about 2 hours and 30 minutes.
- Subsequent infusions: your next 600 mg doses of Ocrevus will be given as a single intravenous infusion every 6 months. These infusions will take about 3 hours and 30 minutes.
- If you did not experience a serious infusion related reaction with any previous infusion, a shorter infusion of approximately 2 hours can be given for subsequent doses.
- Your doctor or nurse will closely monitor you during each infusion and may adjust your infusion rate depending on how well each dose is tolerated.
- Follow the instructions provided and use Ocrevus until your doctor tells you to stop.

Ocrevus subcutaneous (SC)

Ocrevus SC is not intended for intravenous administration. Ocrevus SC must be administered under the supervision of a healthcare professional.

Before you receive Ocrevus you will be given other medicines to help reduce the risk of possible injection reactions.

- The first injection: you will be given 920 mg of Ocrevus by SC injection in approximately 10 minutes. You will be closely monitored for at least one hour after the first injection to manage any possible injection reactions.
- Subsequent injections: if the previous injection was well tolerated, you will be given 920 mg of Ocrevus by SC injection 6 months after the previous injection.

If you miss your dose

If you forget or miss your appointment for your treatment, do not wait until the next planned dose but make another appointment as soon as possible. Your doctor may need to change the timing of your later doses.

If you use too much Ocrevus

As Ocrevus is given under the supervision of your doctor, it is very unlikely that you will be given too much. However, if you experience any side effects after being given Ocrevus, tell your doctor immediately.

You should immediately:

- phone the National Poisons Centre **by calling 0800 POISON (0800 764766)**, or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. What should I know while using Ocrevus?

Things you should do

- Tell your partner or caregiver about your Ocrevus treatment. They might notice symptoms that you do not, such as changes in movement or behaviour, which your doctor may need to investigate.
- If you plan to become pregnant or you become pregnant during treatment with Ocrevus, or if you plan to breastfeed, tell the doctor right away. Tell your baby's doctor about your treatment with Ocrevus so that they can consider this in the timing of your baby's vaccinations.
- Tell your doctor if you have any new medical conditions or plan to take any new medicines for any other condition.
- Tell your doctor if you have signs and symptoms that may indicate a liver problem (e.g. new or worsening fatigue, loss of appetite, nausea, vomiting, pain in the right part of your abdomen, dark-colored urine, or yellowing of your skin or eyes). Your doctor will order a blood test to check your liver.

Call your doctor straight away if you:

- have any signs or symptoms of infusion or injection related reactions or allergic reactions.
- think your multiple sclerosis is getting worse or if you or your partner or caregiver notice any new or unusual symptoms such as changes in movement or behaviour. There have been very rare reports of serious brain infection, called progressive multifocal leukoencephalopathy (PML) in patients receiving medicines for MS, which can cause severe disability or be life-threatening. Symptoms of PML can be similar to those of MS.

Remind any doctor, nurse, dentist or pharmacist you visit that you are using Ocrevus.

Things you should not do

- Do not stop using this medicine without talking to your doctor first. Tell your doctor if you feel that Ocrevus is not helping your condition.
- Do not take any other medicines, whether they require a prescription or not without first telling your doctor or consulting with a pharmacist.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how Ocrevus affects you.

Looking after your medicine

- It is unlikely that you will have to store Ocrevus at home. If you have to store it: refrigerate, do not freeze. Store between 2°C and 8°C. Do not shake it.
- Keep the vial in the carton to protect it from light.

Follow the instructions in the carton on how to take care of your medicine properly.

Keep it where young children cannot reach it.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

Tell your doctor or nurse as soon as possible if you do not feel well while you are receiving Ocrevus. Ocrevus helps most people with multiple sclerosis, but it may have unwanted side effects in some people.

All medicines can have side effects. If you do experience any side effects, most of them are likely to be minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Do not be alarmed by the following lists of side effects. You may not experience any of them.

Less serious side effects

Less serious side effects	What to do
Infections such as: <ul style="list-style-type: none">• viral infection (influenza)• respiratory tract infection (runny nose, sneezing, sore throat, cough, fever, fatigue, weakness, shortness of breath)• inflammation or infection of the sinuses - sinusitis (feeling of pressure or pain in the cheeks and forehead)• inflammation or infection of the lung - bronchitis (cough, chest pain, fever)• build up of mucus in the nose and throat• infection of the stomach and bowel - gastroenteritis (nausea or vomiting, fever)• eye inflammation or infection - conjunctivitis (itchy, red or sore eyes)• skin infection - cellulitis (rash, itching or hives on the skin)	Tell your doctor or nurse straight away if you have any of these side effects, even if it has been sometime since your last infusion or injection. These side effects can sometimes be serious. If you experience signs and symptoms of infection, your doctor may need to do a blood test to check your white blood cell levels even if it has been some time (e.g. several weeks or months) since your last infusion or injection.

Serious side effects

Serious side effects	What to do
<p>Infusion related reactions, injection reactions or allergic reactions:</p> <p>Infusion related reactions or injection reactions can happen during the infusion and within 24 hours after the infusion or injection.</p> <p>Symptoms may include, but are not limited to:</p> <ul style="list-style-type: none"> itchy skin rash hives redness of the skin throat irritation or pain shortness of breath swelling of the throat, face, lips, tongue, or other parts of the body reactions in the area where the injection is given (Ocrevus SC only) redness on your face (flushing) low blood pressure fever fatigue nausea headache fast heart beat <p>Infections and signs or symptoms such as:</p> <ul style="list-style-type: none"> fever or chills cough that does not go away herpes (such as cold sore, shingles and genital sores) <p>Liver problems such as:</p> <ul style="list-style-type: none"> new or worsening fatigue loss of appetite nausea vomiting pain in the right part of your abdomen dark-colored urine yellowing of your skin or eyes 	<p>During your infusion or injection, tell your doctor or nurse straight away if you experience signs and symptoms of infusion related reactions or injection reactions.</p> <p>After your infusion or injection, call your doctor straight away or go straight to the Emergency Department at your nearest hospital if you notice any of these side effects. These may be serious side effects.</p> <p>Tell your doctor straight away if you have any of these side effects, even if it has been sometime since your last infusion or injection.</p>

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Centre for Adverse Reactions Monitoring (CARM) online at <https://pophealth.my.site.com/carmreportnz/s/>. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What Ocrevus IV contains

Active ingredient (main ingredient)	Each vial of Ocrevus IV contains 300 mg of ocrelizumab
Other ingredients (inactive ingredients)	sodium acetate trihydrate trehalose dihydrate glacial acetic acid polysorbate 20 water for injections

What Ocrevus SC contains

Active ingredient (main ingredient)	Each vial of Ocrevus SC contains 920 mg of ocrelizumab
Other ingredients (inactive ingredients)	hyaluronidase sodium acetate trihydrate trehalose dihydrate glacial acetic acid polysorbate 20 methionine water for injections

Do not use this medicine if you are allergic to any of these ingredients.

What Ocrevus IV looks like

Ocrevus IV is supplied as a single-dose glass vial containing 10 mL of solution for intravenous infusion (30 mg/mL). Ocrevus is diluted before infusion into a vein.

Ocrevus IV is a clear or slightly opalescent, and colourless to pale brown liquid.

What Ocrevus SC looks like

Ocrevus SC is supplied as a single-dose glass vial containing 23 mL of solution for subcutaneous injection (40 mg/mL).

Ocrevus SC is a clear or slightly opalescent, and colourless to pale brown liquid.

Storage

Ocrevus will be stored in the pharmacy or on the hospital ward in a refrigerator at a temperature between 2°C and 8°C.

Who distributes Ocrevus

Ocrevus is supplied in New Zealand by:
Roche Products (New Zealand) Limited
PO Box 109113
Newmarket
Auckland 1149
NEW ZEALAND

Medical enquiries: 0800 276 243

This leaflet was prepared in December 2025.