

Brevinor-1[®] 28 Day Tablets

Ethinylestradiol and Norethisterone

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Brevinor-1 28 day tablets.

It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Brevinor-1 against the benefits it is expected to have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What Brevinor-1 is used for

Brevinor-1 is a birth control pill commonly known as a “Combined Oral Contraceptive”. It contains both an estrogen (ethinylestradiol) and progestogen (norethisterone) hormone.

Oral contraceptives belonging to this group produce their birth control (or contraceptive) effect by preventing ovulation (the release of an egg from the ovary) during each menstrual cycle. Combined Oral

Contraceptives also cause changes to the mucus of the cervix and the

lining of the womb which contribute to the contraceptive action.

Ask your doctor if you have any questions about why Brevinor-1 has been prescribed for you.

Your doctor may have prescribed Brevinor-1 for another reason.

Brevinor-1 28 day is only available on a prescription from your doctor.

This medicine is not expected to affect your ability to drive a car or operate machinery.

Before you start to take Brevinor-1

When you must not take it

Do not take Brevinor-1 if you have an allergy to:

- any medicine containing ethinylestradiol or norethisterone
- any of the ingredients listed at the end of this leaflet
- any other similar medicines (such as other oral contraceptives).

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin.

Do not take it if:

- you have had a stroke, or a heart attack

- you have or have had inflammation, infection or clotting in any blood vessel(s), including a clot in the lung
- you have or have had liver disease (including tumours of any type), a history of jaundice or cholestatic jaundice of pregnancy, or severe generalised itch in the body during pregnancy, Dubin-Johnson Syndrome or Rotor Syndrome
- you have abnormal vaginal bleeding, the cause of which is unknown
- you are pregnant or suspect that you may be pregnant
- you have cancer or suspected cancer of the breast or sex organs (e.g. cervix, ovaries, endometrium, womb) and known or suspected estrogen-dependent tumours
- you have a family history of breast nodules, fibrocystic disease or have had an abnormal mammograph
- you have sickle cell anaemia
- you have a lipid metabolism disorder such as congenital hyperlipidaemia
- you have diabetes with blood vessel damage
- you have a history of herpes during pregnancy
- you have otosclerosis (an ear disorder) which worsened in past pregnancies

- you suffer from partial paralysis during migraine attacks (hemiplegic migraine).

Do not take Brevinor-1 if you have Hepatitis C and are taking the combination drug regimen ombitasvir/paritaprevir/ritonavir and dasabuvir with or without ribarivin as this may cause increases in liver function blood test results (increase in ALT liver enzyme). Brevinor-1 can be restarted approximately 2 weeks after completion of treatment with the combination drug regimen.

If you are not certain whether these may apply to you, or you are worried by anything in this list, tell your doctor.

Tell your doctor about any existing medical condition as this may be affected by taking the birth control pill.

Do not take Brevinor-1 if the packaging is torn or shows signs of tampering.

If it is expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should be taking this medicine, talk to your doctor.

Before you start to take it

Tell your doctor if you are allergic to any foods, dyes, preservatives or any other medicines.

You must tell your doctor if:

- you are a heavy smoker (15 or more cigarettes per day), especially if you are aged over 35 years.

Oral contraceptives increase your risk of having a stroke or heart attack. Smoking, while taking oral contraceptives, further increases this risk.

Tell your doctor if you have any other health problems, especially:

- uterine fibroids
- gallbladder disease
- liver, heart or kidney disease

- high blood pressure
- high cholesterol
- diabetes
- epilepsy
- asthma
- migraine or other headaches
- depression
- a disorder of calcium or phosphorus metabolism.

If you have any of these conditions you should have regular check-ups with your doctor to make sure that taking Brevinor-1 is not making these worse.

Tell your doctor if you have any of the following:

- recent surgery or trauma
- recently had a baby
- confined to bed rest for long periods.

The risk of developing blood clots in the deep veins of your body, which can break away and block a blood vessel elsewhere in your body, is increased if you take an oral contraceptive in the above situations.

Tell your doctor if you wear contact lenses and experience a change in vision or intolerance to your lenses.

Your doctor may refer you to an eye specialist.

Brevinor-1 contains lactose. If you know that you are intolerant to some sugars, or your doctor has told you so, speak to your doctor before taking it.

Tell your doctor if you plan to become pregnant or are breastfeeding.

Your doctor can discuss the risks and benefits with you.

If you have not told your doctor about any of the above, tell him/her before you start taking Brevinor-1.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other

medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may alter the effectiveness of Brevinor-1. These include medicines such as:

- rifampicin for the treatment of tuberculosis
- antibiotics such as ampicillin, oxacillin, tetracyclines, sulfamethoxazole and trimethoprim
- anti-fungal agents such as griseofulvin
- barbiturates
- medicines for epilepsy such as carbamazepine and phenytoin
- medicines used to treat or relieve the symptoms of allergy (such as hay fever)
- medicines used to treat or prevent pain
- anti-migraine preparations
- phenylbutazone, a medicine used to relieve symptoms of inflammation, including arthritis
- St John's wort, an ingredient found in medicines you can purchase without a prescription from a pharmacy, supermarket or health food shop.

While you are taking these medicines, and for seven days after stopping them, you should use a non-hormonal method of contraception (such as condoms or a diaphragm, but not the rhythm or temperature methods). If the seven days extend into the inactive orange tablet section, then you should start a new pack on the next day after having taken the last white active tablet from the current pack. Skip the 7 orange tablets.

This is particularly important if you need to take antibiotics or medicines for epilepsy.

Brevinor-1 may alter the effectiveness of some medicines. These include:

- medicines for high blood pressure such as guanethidine
- medicines to prevent blood clots.

Ask your doctor or pharmacist about how long you need to use additional non-hormonal contraception.

If you have not told your doctor or pharmacist about any of the above, tell them before you start taking Brevinor-1.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while you are taking this medicine.

Do not take Brevinor-1, and tell your doctor if you are taking the combination drug regimen ombitasvir/paritaprevir/ritonavir and dasabuvir with or without ribarivir for the treatment of Hepatitis C. Treatment with Brevinor-1 should be stopped when you are taking this combination of treatment regimen and can be restarted approximately 2 weeks after completion of this treatment.

How to take Brevinor-1

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the box, ask your doctor or pharmacist for help.

How to take it

Swallow one Brevinor-1 tablet with a glass of water.

When to take it

You must take Brevinor-1 every day, regardless of how often you have sex.

Brevinor-1 will work best if you do not miss any tablets and take it at the same time each day. Taking

Brevinor-1 at the same time each day will also help you remember when to take your tablets.

If you are concerned about this, please speak to your doctor or pharmacist.

Starting a hormonal contraceptive for the first time

- To begin Brevinor-1 28 Day, take your first tablet on the first day of your next period, that is, the day your bleeding starts.
- Take your first tablet from the top row of the green section of the strip (i.e. the section which contains all the white tablets). Take the tablet which corresponds to the appropriate day of the week. For example, if your first day of bleeding is on TUESDAY, you should take the tablet marked “TUE” from the top row of the green section of the strip.
- Continue to take one tablet every day, following the arrows around the strip, until all 21 white tablets are finished.
- You should then take one orange tablet daily for the next 7 days, following the arrows so that you are taking the correct tablet for the day of the week. Taking these orange inactive tablets helps you to remember to take a tablet every day.

You can expect your period during the week that you are taking these orange inactive tablets. Your protection continues during this week.

- On the day after your last orange tablet, begin the next strip with a white tablet from the top row of the green section that matches the day of the week. Do this even if you are still bleeding.
- You should start your tablets the same day of the week every four weeks.

- Repeat this sequence of tablet taking for as long as birth control is required.

This product is effective from the first day if taken as directed above.

Although spotting and break-through bleeding may occur in some women, these tend to disappear in the majority of patients after the first three to four cycles.

Make sure you always have a new strip of tablets available, so that you can continue to take the tablets without interruption.

Changing from a different oral contraceptive

If you are switching to Brevinor-1 28 Day from another 21 or 28 day oral contraceptive, follow the instructions below carefully.

If switching from a 21 day oral contraceptive:

- Stop taking your current oral contraceptive after you have taken the last active tablet.
- Leave 7 tablet-free days.
- Start the new Brevinor-1 28 Day pack on the eighth day by taking a white active tablet from the top row of the green section which corresponds to the day of the week.
- Continue to take one tablet every day, following the arrows around the strip until you finish all 21 white tablets in the green section of the strip.
- Then take one orange tablet daily for the next 7 days, before starting your new strip.

You must use an additional, non-hormonal method of contraception (such as condoms or a diaphragm, but not the rhythm or temperature methods) until a white tablet has been taken daily for 7 days without a break.

If switching from a 28 day oral contraceptive:

- Stop taking your current oral contraceptive after you have

taken the last inactive tablet in the strip.

- Start the new Brevinor-1 28 Day pack on the next day by taking a white active tablet from the top row which corresponds to the day of the week.
- Continue to take one tablet every day, following the arrows around the strip until you finish all 21 white tablets in the green section of the strip.
- Then take one orange tablet daily for the next 7 days, before starting your new strip.

You must use an additional, non-hormonal method of contraception (such as condoms or a diaphragm, but not the rhythm or temperature methods) until a white tablet has been taken daily for 7 days without a break.

If you vomit or have diarrhoea after taking Brevinor-1

If you suffer from a stomach upset which results in vomiting or diarrhoea, the effectiveness of Brevinor-1 may be reduced.

During any period of vomiting or diarrhoea, continue taking Brevinor-1 tablets. Also use a non-hormonal method of contraception (such as condoms or a diaphragm, but not the rhythm or temperature methods), and continue for seven days following the episode of vomiting or diarrhoea. If these seven days extend into the inactive orange tablet section, you should start a new pack on the next day after having taken the last white tablet from the green section of the current pack (i.e. skip the orange inactive tablets)

You may not have a period until you finish the second pack.

If you have vomiting or diarrhoea after taking an orange tablet, do not worry.

If you forget to take it

If you forget to take Brevinor-1 it may not work as well in protecting you from becoming pregnant.

Do not try to make up for missed doses by taking more than one tablet at a time.

If you miss a white active tablet:

If you are less than 12 hours late in taking your tablet, you should take that tablet at once and then take the next one at your usual time.

If you are more than 12 hours late in taking your tablet, do not take it.

Take the next day's tablet at the usual time and use an additional, non-hormonal method of contraception for the next seven days (such as condoms or a diaphragm, but not the rhythm or temperature methods).

If the seven days extend into the inactive orange tablet section, then start a new pack on the day after taking the last white active tablet from the green section of the current pack. Take your first tablet from the top row of the green section of the new strip, then repeat the sequence of tablet taking for as long as birth control is required. This will mean that you will not have a period until you finish the second pack.

If you miss more than one white tablet, contact your doctor for advice on what to do.

If you miss an orange (inactive) tablet:

Take it as soon as you remember and continue on as before.

An additional birth control method is not necessary in this case.

If your doctor told you to take Brevinor-1 differently, or you are unclear about the above directions, then you should discuss this with him or her.

If you have trouble remembering to take Brevinor-1, ask your pharmacist for some hints.

If you miss a period

If you have missed a period you may be pregnant.

Contact your doctor to check if you are pregnant.

If you take too much (overdose)

Immediately telephone your doctor or pharmacist or the National Poisons Centre (telephone 0800 764 766) or go to Accident and Emergency at your nearest hospital if you think you or someone else may have taken too much Brevinor-1. Do this even if there are no signs of discomfort or poisoning.

Serious ill effects have not been reported in young children who have taken large doses of birth control pills.

Overdosage may cause nausea or vomiting. This may be followed by vaginal bleeding or breast enlargement in some women.

While you are taking Brevinor-1

Things you must do

Tell all doctors, dentists and pharmacists who are treating you that you are taking Brevinor-1.

Tell the hospital doctor that you are taking Brevinor-1 birth control pills if you need to have an operation or go to hospital in an emergency.

If you are about to be started on any new medicine, tell your doctor or dentist and your pharmacist that you are taking Brevinor-1.

If you become pregnant while taking Brevinor-1, see your doctor immediately.

If you are about to have any blood tests, tell your doctor you are taking Brevinor-1.

It may interfere with the results of some tests.

Visit your doctor regularly for check-ups, including a Pap smear.

A pap smear can detect any abnormal cells from the cervix, which may develop into cancer. Cervical cancer has been reported to occur more frequently in women who use oral contraceptives.

Your doctor will advise you of the type and frequency of any tests required.

Perform regular breast self-examination.

Examining your breasts for lumps or any changes in size or shape can help you find a breast cancer early. Breast cancer has been found more frequently in women who use oral contraceptives. It is not known whether this increase is caused by the use of oral contraceptives, or if it is due to the fact that users were examined more often, and therefore the breast cancer was detected earlier.

If you are unsure, ask your doctor about breast self-examination.

If you are worried about contracting a sexually transmitted disease (STD) use a condom during sexual intercourse.

Brevinor-1 does not protect against the transmission of STDs such as HIV-AIDS, chlamydia, genital herpes and warts, gonorrhoea, hepatitis B or human papilloma virus. To protect against STDs ask your partner to wear a condom when having sexual intercourse with you.

Tell your doctor if you feel depressed, think you are retaining water, experience headaches, experience persistent or recurrent irregular bleeding, or your eyes are uncomfortable whilst wearing contact lenses.

Your doctor will make an assessment of your condition and advise whether or not you should continue to take Brevinor-1.

Things you must not do

Do not take Brevinor-1 to treat any other complaints unless your doctor tells you to.

Do not give your medicine to anyone else even if they have the same condition as you.

Do not stop taking Brevinor-1, or change the dosage, without checking with your doctor.

If you stop taking Brevinor-1 or do not take a tablet every day, without using another form of contraception, you may become pregnant.

Things to be careful of

Slight breast tenderness or a feeling of sickness may occur in the first few months of use. This usually improves or stops with continued use.

If vaginal irritation or discharge occurs, it may be an indication of a yeast infection for which treatment is available from your doctor.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Brevinor-1.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

It can be difficult to tell whether side effects are the result of taking Brevinor-1 or are side effects of another medicine you are taking.

Do not be alarmed by the list of side effects.

You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor if...

Tell your doctor if you notice any of the following and they worry you:

- break-through bleeding
- spotting
- gastric or stomach discomforts including nausea, vomiting and diarrhoea
- change in menstrual flow
- change in weight
- retention of fluids
- dark discolouration of the skin
- blotchy discolouration on the face or limbs (which may persist after the tablets have been stopped)
- absence of periods
- breast changes (tenderness, enlargement and secretion)
- headache, dizziness
- mental depression
- fatigue or tiredness
- hair growth or loss of scalp hair
- acne, rashes, itching
- leg cramps
- backache
- vaginal thrush, vaginal irritation, bladder irritation or urinary tract infection
- pre-menstrual-like symptoms
- change in cervical secretions
- suppression of milk production
- change in appetite
- contact lenses becoming difficult to wear.

The above list contains the more common side effects of your medicine.

Tell your doctor as soon as possible if...

Tell your doctor as soon as possible if you notice any worsening of conditions that you may already have such as:

- porphyria
- gallbladder disease

- kidney disease
- hereditary angioedema (swelling of the face lips, mouth tongue or throat).

Oral contraceptives are associated with inflammatory bowel diseases (ulcerative colitis, Crohn's disease). See your doctor as soon as possible if you develop bloody diarrhoea, abdominal pain, fever, rectal bleeding, feeling tired, lose your appetite or lose weight. These are some of the symptoms of inflammatory bowel diseases.

The above list includes serious side effects that may require medical attention. Serious side effects are rare.

Go to hospital if...

Tell your doctor immediately or go to Accident and Emergency at your nearest hospital, if you notice any of the following:

- unexplained or persistent pains in the head, chest, stomach or legs
- gradual or sudden, partial or complete loss of vision
- double vision, or symptoms of severe vision impairment
- eye protrusion, swelling of the eye or eye lesions
- migraine headaches for the first time
- more frequent or severe migraines if you already suffer from them
- breast lumps
- jaundice or a yellowing of the skin or eyes, often with fever, fatigue, loss of appetite, dark coloured urine, nausea and vomiting.

Taking oral contraceptives may be associated with liver disease including liver cancer.

- rise in blood pressure

You may experience headache, blurred vision or palpitations. Sometimes your blood pressure may rise without you experiencing any of

these symptoms. It is important to keep your routine doctor's appointments so that your blood pressure can be checked.

- swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or trouble breathing.

Whilst these side effects are rare, they are serious. You may need urgent medical attention or hospitalisation.

Other side effects not listed above may also occur in some patients.

Tell your doctor if you notice anything else that is making you feel unwell, even if it is not on this list.

After stopping Brevinor-1

Delays in becoming pregnant may occur after Brevinor-1 therapy is stopped. This is more likely to occur in women whose periods were irregular before using birth control pills.

See your doctor if you continue to experience difficulties in falling pregnant.

After taking Brevinor-1

Storage

Keep your Brevinor-1 tablets in a safe place away from the sight and reach of children.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Keep your Brevinor-1 tablets in a dry place, at a temperature below 25 degrees C.

Do not keep your tablets in the refrigerator.

Do not store Brevinor-1 or any other medicine in the bathroom or near a sink. Do not leave it in the car on hot days or on window sills.

Heat and dampness can destroy some medicines.

Disposal

If your doctor tells you to stop taking Brevinor-1 or if the tablets have passed their expiry date, ask your pharmacist what to do with any that are left over.

Product description

What it looks like

Brevinor-1 28 Day is available in calendar packs consisting of three strips of tablets (3 months' supply) each with 21 white active tablets marked "SEARLE" on one side and "BX" on the other, and 7 orange inactive tablets.

Ingredients

Each white tablet contains:

- norethisterone 1 milligram (mg)
- ethinylestradiol 35 micrograms (µg)
- magnesium stearate
- povidone
- maize starch
- lactose monohydrate.

Each orange tablet contains:

- magnesium stearate
- microcrystalline cellulose
- lactose and lactose monohydrate
- sunset yellow FCF (CI No. 15985).

Brevinor-1 does not contain sucrose, gluten or tartrazine.

Supplier

Brevinor-1 is supplied in New Zealand by:

Pfizer New Zealand Limited

PO Box 3998

Auckland, New Zealand

Toll Free number: 0800 736 363

Date of preparation

This leaflet was prepared in March 2017.

© Pfizer Australia Pty Ltd 2017

® Registered Trademark