

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Rocuronium bromide Medsurge.

It does not contain all the available information.

It does not take the place of talking to your doctor.

All medicines have risks and benefits. Your doctor has weighed the risks of you having Rocuronium bromide Medsurge against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor.

Keep this leaflet.

You may need to read it again.

What Rocuronium bromide Medsurge is used for

Rocuronium bromide Medsurge is one of a group of medicines called muscle relaxants.

Muscle relaxants are used during an operation as part of the general anaesthetic. When you have an operation, your muscles must be completely relaxed. This makes it easier for the surgeon to perform the operation.

Normally the nerves send messages to the muscles by impulses. Rocuronium bromide Medsurge acts by blocking these impulses so the muscles are relaxed. Because the muscles needed for breathing also become relaxed you will need help with your breathing (artificial respiration) during and after your operation until you can breathe on your own. During the operation the effect of the muscle relaxant is

constantly checked and if necessary, some more drug is given. At the end of the operation the effects of Rocuronium bromide Medsurge are allowed to wear off and you can start breathing on your own. Sometimes another drug is given to help speed this up. Rocuronium bromide Medsurge can also be used in Intensive Care to keep your muscles relaxed.

Ask your doctor if you want any more information about this medicine.

Rocuronium bromide Medsurge is not addictive.

Before you are given Rocuronium bromide Medsurge

When you must not be given it

You must not be given Rocuronium bromide Medsurge if you have an allergy to:

- any medicine containing rocuronium bromide
- any of the ingredients listed at the end of this leaflet

Some of the symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing; swelling of the face, lips, tongue or other parts of the body; rash, itching or hives on the skin.

Before you are given it

If you are going to have an operation it is important that you discuss the following points with your doctor, since it can influence the way Rocuronium bromide Medsurge is given to you.

Tell your doctor if you have or have had any medical conditions, especially the following:

- an allergy to muscle relaxants
- kidney disease
- liver or gallbladder disease
- a heart disease
- diseases affecting nerves or muscles
- oedema (local or generalised swelling due to fluid)

Certain medical conditions may affect how Rocuronium bromide Medsurge works.

- low potassium levels in the blood
- high magnesium levels in the blood
- low calcium levels in the blood
- low levels of protein in the blood
- dehydration
- too much acid in the blood
- too much carbon dioxide in the blood
- general ill-health
- overweight
- burns

If you are suffering from any of these conditions your doctor will take this into account when deciding the correct dose of Rocuronium bromide Medsurge for you.

Rocuronium bromide Medsurge can be used in term newborn infants (0-28 days), infants (28 days-23 months), children (2-11 years), adolescents (12-18 years) and elderly patients but your doctor should first assess your medical history.

Elderly (65 years and older) may be at an increased risk of residual paralysis.

Rocuronium bromide 50 mg/5 mL solution for injection

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you are pregnant or plan to become pregnant or are breast-feeding.

Your doctor will discuss the risks and benefits of using Rocuronium bromide Medsurge if you are pregnant or breast-feeding.

If you have not told your doctor about any of the above, tell them before you are given Rocuronium bromide Medsurge.

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may be affected by Rocuronium bromide Medsurge or affect how well it works. These include:

- anaesthetics, medicines to make you sleep during surgery
- long term concurrent use of corticosteroids (anti-inflammatory medicines) and Rocuronium bromide Medsurge in the Intensive Care Unit
- antibiotics
- lithium, a medicine used to treat bipolar disorder
- medicines used to treat heart disease or high blood pressure (quinidine, calcium channel blockers, beta-blockers and diuretics (fluid tablets))
- quinine, a medicine used to treat malaria
- magnesium salts

- lignocaine and bupivacaine hydrochloride (local anaesthetics)
- other muscle relaxants
- carbamazepine and phenytoin, medicines used to treat epilepsy

You may need to use different amounts of your medicines or take different medicines. Your doctor will advise you.

If you are taking magnesium sulphate to treat toxemia of pregnancy (preeclampsia), tell your doctor as the dose of Rocuronium bromide Medsurge may need to be reduced.

Your doctor will have a complete list of medicines that may cause problems when used with Rocuronium bromide Medsurge.

How Rocuronium bromide Medsurge is given

Rocuronium bromide Medsurge will be given by a doctor. It will not be given to you until you are asleep from the anaesthetic.

It will be injected into a vein before and/or during an operation. It will be given as a single injection or continuous infusion.

The usual dose is 0.6 mg rocuronium bromide per kg body weight and the effect lasts 30-40 minutes. During the operation your doctor will check whether Rocuronium bromide Medsurge is still working.

You may be given additional doses if they are needed.

Overdose

As Rocuronium bromide Medsurge doses are carefully worked out and are given by a doctor experienced in its use, it is extremely unlikely that you will be given too much Rocuronium bromide Medsurge.

However, if this does happen, your doctor will make sure that you continue breathing artificially until you can breathe on your own again. Your doctor may speed-up your recovery by giving you a drug that reverses the effects of Rocuronium bromide Medsurge.

After having Rocuronium bromide Medsurge

Things to be careful of

Your doctor will tell you when it is safe to drive and operate potentially dangerous machinery after you have been given Rocuronium bromide Medsurge.

Side effects

Tell your doctor if you notice anything that is making you feel unwell.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following list of side effects. You may not experience any of them.

Ask your doctor to answer any questions you may have.

Tell your doctor if you notice the following and they worry you:

- flushing
- pain at injection site
- irritation at injection site
- red skin rash or itchy rash

Tell your doctor immediately if you notice any of the following:

- fast heartbeat
- dizziness, light-headedness (low blood pressure)
- muscle weakness or paralysis

Rocuronium bromide 50 mg/5 mL solution for injection

- aching muscles or weakness, not caused by exercise
- wheezing, coughing
- difficulty breathing
- rapid, shallow breathing, cold, clammy skin, a rapid, weak pulse, dizziness, weakness and fainting
- swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing
- sudden signs of allergy such as rash, itching, hives on the skin, swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or trouble breathing
- Sudden fever with rapid heartbeat, rapid breathing and stiffness, pain and/or weakness in your muscles
- Severe allergic coronary blood vessels spasm (Kounis syndrome) resulting in chest pain (angina) or heart attack (myocardial infarction)
- Dilated pupils (mydriasis) or fixed pupils

These are serious side effects. You may need urgent medical attention or hospitalisation.

Storage

Rocuronium bromide Medsurge is stored in the hospital.

It should be kept in the refrigerator at 2-8°C and not be frozen. Rocuronium bromide Medsurge can be stored outside the refrigerator at a temperature of 8-25°C for a maximum of 12 weeks. After first removal from the refrigerator, the 12 weeks shelf life applies. The storage period may not exceed the labelled shelf-life.

Product description

What it looks like

A vial containing a clear, colourless to pale yellow solution for injection.

Rocuronium bromide Medsurge is available in clear glass vials in packs of 10 vials

Ingredients

Rocuronium bromide Medsurge contains 10 mg/mL of rocuronium bromide as the active ingredient. It also contains:

- sodium acetate
- sodium chloride
- glacial acetic acid
- water for injections

No preservative has been added.

Supplier

Medsurge Pharma Limited
PO Box 331054
Takapuna
Auckland 0622
Telephone: 0800 788 261

**Marketed and Distributed by
Medsurge Healthcare Pty Ltd.**
Telephone: 1300 788 261
Website: <https://medsurgehc.com>

This leaflet was revised in July 2024