

## TENOFOVIR DISOPROXIL EMTRICITABINE EFAVIRENZ MYLAN

*Tenofovir disoproxil maleate, Emtricitabine, Efavirenz  
Film coated tablet 300mg, 200mg, 600mg*



---

### What is in this leaflet

---

Please read this leaflet carefully before you start taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.

This leaflet answers some common questions about Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan against the benefits they expect it will have for you.

**If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.** You may need to read it again.

---

### What Tenofovir disoproxil Emtricitabine Efavirenz Mylan is used for

---

Tenofovir Disoproxil Emtricitabine Efavirenz Mylan is used to treat Human Immunodeficiency Virus (HIV-1) infection in adults. This medicine can be taken alone or in combination with other anti-HIV medicines.

Tenofovir Disoproxil Emtricitabine Efavirenz Mylan consists of three medicines:

- Tenofovir disoproxil maleate

- Emtricitabine
- Efavirenz

These are combined in one tablet to help control Human Immunodeficiency Virus (HIV-1) infection.

Tenofovir disoproxil and emtricitabine belong to a group of antiviral medicines known as nucleoside and nucleotide reverse transcriptase inhibitors (NRTI).

Efavirenz belongs to a group of antiviral medicines known as non-nucleoside reverse transcriptase inhibitors (NNRTI).

#### How Tenofovir Disoproxil Emtricitabine Efavirenz Mylan works

HIV-1 infection destroys CD4 T cells, which are important to the immune system. The immune system helps fight infection. After a large number of T cells are destroyed, acquired immune deficiency syndrome (AIDS) may develop.

Tenofovir Disoproxil Emtricitabine Efavirenz Mylan helps block HIV-1 reverse transcriptase, a viral chemical in your body (enzyme) that is needed for HIV-1 to multiply. It lowers the amount of HIV-1 in the blood (viral load). It may also help increase the number of T cells (CD4+ cells), allowing your immune system to improve. Lowering the amount of HIV-1 in the blood lowers the chance of death of infection that happen when your immune system is weak (opportunistic infections).

#### Use in children and elderly

Tenofovir Disoproxil Emtricitabine Efavirenz Mylan is for adults. Do not take Tenofovir Disoproxil

Emtricitabine Efavirenz Mylan if you are under the age of 18 years. Do not take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan if you are over the age of 65 before discussing with your doctor.

---

### Does Tenofovir Disoproxil Emtricitabine Efavirenz Mylan cure HIV or AIDS

---

**Tenofovir Disoproxil Emtricitabine Efavirenz Mylan does not cure HIV-1 infection or AIDS.**

The long-term effects of Tenofovir Disoproxil Emtricitabine Efavirenz Mylan are not known at this time. People taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan may still get opportunistic infections or other conditions that happen with HIV-1 infection. Opportunistic infections are infections that develop because the immune system is weak. Some of these conditions are:

- Pneumonia
- Herpes virus infections, and
- *Mycobacterium avium complex* (MAC) infection.

**This medicine is only available from a pharmacist after it has been prescribed by a doctor who specializes in the treatment of HIV-1 infection.**

**If you wish to continue receiving treatment with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan it is important you remain under the care of a hospital or doctor who specializes in the treatment of HIV-1 infection.**

---

## Does Tenofovir Disoproxil Emtricitabine Efavirenz Mylan reduce the risk of passing HIV to others

---

Tenofovir Disoproxil Emtricitabine Efavirenz Mylan has not been shown to lower your chance of passing HIV to others through sexual contact, sharing needles, or being exposed to your blood.

Do not share needles or other injection equipment.

Do not share personal items that can have blood or body fluids on them, like toothbrushes or razor blades.

Do not have any kind of sex without protection.

---

## Before you take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan

---

### *Who must not take it*

Together with your doctor, you need to decide whether Tenofovir Disoproxil Emtricitabine Efavirenz Mylan is right for you. If you are not sure whether you should be taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan, talk to your doctor.

Do not take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan if you are allergic to:

- Tenofovir
- Emtricitabine
- Efavirenz or
- any of the other ingredients of Tenofovir Disoproxil Emtricitabine Efavirenz Mylan

Some of the symptoms of an allergic reaction may include:

- Shortness of breath
- Wheezing or difficulty breathing
- Rash, itching or hives on the skin

- Swelling of the face, lips, tongue or other parts of the body.

The ingredients of Tenofovir Disoproxil Emtricitabine Efavirenz Mylan are listed in the product description section of this leaflet.

If you are not sure whether you should be taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan, talk to your doctor.

### *Before you start to take it*

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you are pregnant, or planning to become pregnant during your course of medication.

Women taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan should not become pregnant. Serious birth defects have been seen in babies of animals and women treated with efavirenz during pregnancy. It is not known whether efavirenz caused these defects.

Tell your doctor immediately if you become pregnant while taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.

Also, talk to your doctor if you want to become pregnant.

Do not breast-feed if you are taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.

Do not breast-feed if you are taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan. The active substances in this medicine (tenofovir and emtricitabine and efavirenz) have been found in breast milk at low concentrations.

It is recommended that nursing mothers do not breast-feed during treatment with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan. In general, women infected with HIV should not breast feed their infants in order to avoid transmission of HIV to their new-born infants.

Tell your doctor if you have kidney problems or are

undergoing kidney dialysis treatment.

Tell your doctor if you have bone problems

Tell your doctor if you have liver problems, including Hepatitis B Virus (HBV) infection.

Your doctor may want to do tests to check your liver while you take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.

Tell your doctor if you have ever had mental illness or are using recreational drugs or alcohols.

Tell your doctor if you have ever had seizures or are taking medicines for seizures.

Tell your doctor if you have ever had a serious allergic drug reaction involving the skin (e.g. Stevens-Johnson syndrome).

### *Taking other medicines*

Tenofovir Disoproxil Emtricitabine Efavirenz Mylan may change the effect of other medicines, including the ones for HIV-1 and may cause serious side effects.

Your doctor may change your other medicines or change their doses. Other medicines, including herbal products may affect Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.

For this reason, it is very important to let your doctor or pharmacist know what medications, herbal supplements or vitamins you are taking.

### *Medicines you should not take with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan*

The following medicines may cause serious or life-threatening side effects when taken with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.

Do not take elbasvir/grazoprevir with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.

**It may lose its effect and reduce your response to elbasvir/grazoprevir**

**Do not take glecaprevir/pibrentasvir with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan. It may lose its effect and reduce your response to glecaprevir/pibrentasvir.**

**Do not take St. John's wort (Hypericum perforatum) or products containing St. John's wort with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.**

St. John's wort is a herbal product sold as a dietary supplement. Talk to your doctor or pharmacist if you are taking or are planning to take St. John's wort. Taking St. John's wort may decrease Tenofovir Disoproxil Emtricitabine Efavirenz Mylan levels and lead to increased viral load and possible resistance to Tenofovir Disoproxil Emtricitabine Efavirenz Mylan or cross-resistance to other anti-HIV drugs.

**Voriconazole should not be taken with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.**

It may lose its effect or may increase the chance of having side effects from Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.

**Do not take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan if you are already taking any other medicines that contain the same active ingredients.**

**Do not take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan if you are taking other medicines that contain:**

- Efavirenz
- Lamivudine
- Tenofovir alafenamide

Do not take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan to treat your HIV infection if you are also taking adefovir dipivoxil.

**It is also important to tell your doctor if you are taking any of the following:**

- Bepridil, cisapride, midazolam, pimozide, triazolam, ergot medications.
- Saquinavir, clarithromycin or itraconazole; these medicines may need to be replaced with another medicine when taken with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.
- Calcium channel blockers such as diltiazem and others; indinavir; methadone, rifabutin, rifampin.
- Lipid-lowering medicines such as atorvastatin, pravastatin or simvastatin.
- Antidepressant medicines such as sertraline or bupropion.
- Immunosuppressant medicines cyclosporine, tacrolimus or sirolimus; these medicines may need to have their dose changed when taken with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.
- Didanosine; tenofovir may increase the amount of didanosine in your blood, which could result in more side effects. You may need to be monitored more carefully if you are taking didanosine and Tenofovir Disoproxil Emtricitabine Efavirenz Mylan together. Also, the dose of didanosine may need to be changed.
- Atazanavir sulphate or lopinavir/ritonavir; these medicines may increase the amount of tenofovir in your blood which could result in more side effects. Atazanavir sulphate is not recommended with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan. You may need to be monitored more carefully if you are taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan and lopinavir/ritonavir together. Also, the dose of lopinavir/ritonavir may need to be changed.
- Other HIV medications including amprenavir, ritonavir, fosamprenavir calcium, raltegravir, ritonavir or maraviroc. Also, the

dose of maraviroc may need to be changed.

- Medicines for seizures such as phenytoin, carbamazepine or phenobarbitone; your doctor may want to switch you to another medicine or check drug levels in your blood from time to time.
- Antifungal medications such as itraconazole and posaconazole; these medicines may need to be replaced with another medicine when taken with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.
- Hepatitis C antiviral agents such as ledipasvir/sofosbuvir or sofosbuvir/velpatasvir, sofosbuvir/velpatasvir/voxilaprevir, simeprevir, boceprevir, and telaprevir; these medicines may need to be replaced with another medicine when taken with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.
- Antimalarial agents such as artemether/lumefantrine.

**These are not all the medicines that may cause problems if you take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan. Be sure to tell your doctor or pharmacist about all medicines that you take.**

---

## **How to take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan**

---

**Follow all directions given to you by your doctor or pharmacist carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the bottle, ask your doctor or pharmacist for help.**

### ***How much to take***

The usual dose is one Tenofovir Disoproxil Emtricitabine Efavirenz Mylan tablet a day.

### ***How to take it***

**Always take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan on an empty stomach.**

Swallow the tablet whole with a full glass of water.

Take the tablet at bedtime may make some side effects less bothersome.

### ***When to take it***

**Take your medicine at about the same time each day.**

Taking it at the same time each day will have the best effect. It will also help you remember when to take it.

**Take your medicine on an empty stomach, for example, 1 hour before food or 2 hours after food.**

Food can interfere with the absorption of this medicine.

### ***How long to take it***

**Continue taking your medicine for as long as your doctor tells you.**

This medicine helps to control your condition but does not cure it. It is important to keep taking your medicine even if you feel well.

### ***If you forget to take it***

**Do not miss a dose of Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.**

**Otherwise, take it as soon as you remember, and then go back to taking your medicine as you would normally.**

**Do not take a double dose to make up for the dose that you missed.**

This may increase the chance of you getting an unwanted side effect.

**If you are not sure what to do, ask your doctor or pharmacist.**

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

---

## **While you are taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan**

---

### ***Things you must do***

**If you are about to be started on any new medicine, tell your doctor and pharmacist that you are taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.** Your doses may need adjustment.

**Tell any other doctors, dentists, and pharmacists who treat you that you are taking this medicine.**

**If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine.** It may affect other medicines used during surgery.

**If you become pregnant while taking this medicine, tell your doctor immediately.**

**If you are about to have any blood tests, tell your doctor that you are taking this medicine.** It may interfere with the results of some tests.

**Keep all of your doctor's appointments so that your progress can be checked.** Your doctor may do some tests from time to time to make sure the medicine is working and to prevent unwanted side effects.

**When your Tenofovir Disoproxil Emtricitabine Efavirenz Mylan supply starts to run low, get more from your doctor or pharmacy.**

This is very important because the amount of virus in your blood may increase if the medicine is stopped for even a short time. The virus may develop resistance to Tenofovir Disoproxil Emtricitabine Efavirenz Mylan and become harder to treat.

### ***Things you must not do***

**Do not take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan to treat any other complaints unless your doctor tells you to.**

**Do not give your medicine to anyone else, even if they have the same condition as you.**

**Do not stop taking your medicine or lower the dosage without checking with your doctor.** If you stop taking it suddenly, your condition may worsen or you may

have unwanted side effects.

**Do not take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan after the expiry date on the label.** If you take it after the expiry date has passed, it may not work as well.

**Do not take any other medicines, including prescription or non-prescription medicines and herbal products, without checking with your doctor.**

### ***Things to be careful of***

**Be careful driving or operating machinery until you know how Tenofovir Disoproxil Emtricitabine Efavirenz Mylan affects you.**

This medicine may cause dizziness, impaired concentration and/or drowsiness in some people. If you have any of these symptoms, do not drive, operate machinery or do anything else that could be dangerous.

**If you feel light-headed, dizzy or faint when getting out of bed or standing up, get up slowly.** Standing up slowly, especially when you get up from bed or chairs, will help your body get used to the change in position and blood pressure. If this problem continues or gets worse, talk to your doctor.

Taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan with alcohol or other medicines causing similar side effects as Tenofovir Disoproxil Emtricitabine Efavirenz Mylan, such as drowsiness, may increase those side effects.

---

## **In case of overdose**

---

### ***If you take too much (overdose)***

**Immediately telephone your doctor or the National Poisons Centre (telephone 0800 POISON or 0800 764 766), or go to accident and emergency at your nearest hospital, if you think that you or anyone else may have taken too much Tenofovir Disoproxil Emtricitabine Efavirenz Mylan. Do this even if there are no signs of discomfort or poisoning.**

You may need urgent medical attention.

---

## Side effects

---

Like all medicines, Tenofovir Disoproxil Emtricitabine Efavirenz Mylan can have side effects, although not everybody gets them. Some may be serious and need medical attention.

**Check with your doctor as soon as possible if you have any problems while taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan, even if you do not think the problem are connected with the medicine or are not listed in this leaflet.**

**Tenofovir Disoproxil Emtricitabine Efavirenz Mylan may cause the following serious side effects:**

### Lactic Acidosis

**If you have any of the following symptoms after taking your medicine, tell your doctor IMMEDIATELY or go to the accident and emergency department at your nearest hospital:**

- You feel very weak or tired
- You have unusual (not normal) muscle pain
- You have trouble breathing
- You have stomach pain with nausea and vomiting
- You feel cold, especially in your arms and legs
- You feel dizzy or light headed
- You have a fast or irregular heartbeat

These side effects may be due to a condition called lactic acidosis (build-up of an acid in the blood).

**Lactic acidosis can be a medical emergency and may need to be treated in the hospital.**

### Serious Liver problems (hepatotoxicity)

**If you have any of the following symptoms while taking your medication, tell your doctor IMMEDIATELY or go to the accident and emergency department at your nearest hospital:**

- Your skin or white part of your eyes turns yellow (jaundice)
- Your urine turns dark
- Your bowel movements (stools) turn light in colour
- You don't feel like eating food for several days or longer
- You feel sick to your stomach (nausea)
- You have lower stomach area (abdominal) pain

These side effects may be due to a condition called hepatotoxicity with liver enlargement (hepatomegaly) and fat deposits in the liver (steatosis) which sometimes occurs in patients taking anti-HIV-1 medicines.

**You may be more likely to get lactic acidosis or liver problems if you are female, very overweight (obese), or have been taking similar nucleoside analog-containing medicines, like Tenofovir Disoproxil Emtricitabine Efavirenz Mylan, for a long time.**

### Hepatic Flares

**If you have HIV-1 infection and hepatitis B virus (HBV) infection you should not stop your Tenofovir Disoproxil Emtricitabine Efavirenz Mylan treatment without first discussing this with your doctor, as some patients have had blood tests or symptoms indicating a worsening of their hepatitis ("hepatic flare") after stopping individual component (tenofovir DM, emtricitabine and efavirenz) of Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.**

You may require medical exams and blood tests for several months after stopping treatment. Tenofovir Disoproxil Emtricitabine Efavirenz Mylan is not approved for the treatment of HBV, so you must discuss your HBV therapy with your doctor.

### Serious Psychiatric problems

A small number of patients may experience severe depression, strange thoughts, or angry behaviour while taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.

Some patients have thoughts of suicide and a few have actually committed suicide.

These problems may occur more often in patients who have had mental illness.

**Contact your doctor right away if you think you are having these psychiatric problems, so your doctor can decide if you should continue to take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.**

### Kidney problems

If you have had kidney problems in the past or take other medicines that can cause kidney problems, your doctor should do regular blood tests to check your kidneys.

### Change in Bone Mineral Density (thinning bones)

It is not known whether long-term use of Tenofovir Disoproxil Emtricitabine Efavirenz Mylan will cause damage to your bones. If you have had bone problems in the past, your doctor may need to do tests to check your bone mineral density or may prescribe medicines to help your bone mineral density.

### Allergy

Some people are allergic to medicines.

**If you have any of the following symptoms soon after taking your medicine, DO NOT TAKE ANY MORE Tenofovir Disoproxil**

## **Emtricitabine Efavirenz Mylan and tell your doctor**

**IMMEDIATELY or go to the accident and emergency department at your nearest hospital.**

- Skin troubles such as lumpy skin rash or “hives”
- Swelling of the face, lips, mouth or throat which may cause difficulty in swallowing or breathing
- Wheezing, chest pain or tightness
- Fainting

**These are very serious effects. If you have them, you may have a serious allergic reaction. You may need urgent medical attention or hospitalisation. Hypersensitivity reactions are very rare.**

### **Pancreatitis**

**If you have any of the following symptoms after starting your medication, tell your doctor IMMEDIATELY or go to the Accident and Emergency department at your nearest hospital:**

- Severe stomach pain or cramps
- Nausea
- Vomiting

These side effects may be due to a condition called pancreatitis which sometimes occurs in patients taking anti-HIV-1 medicines.

### **Common side effects:**

Patients may have the following during treatment with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan:

- Dizziness
- Headache
- Trouble sleeping
- Drowsiness
- Trouble concentrating

- Unusual dreams

These side effects may be reduced if you take Tenofovir Disoproxil Emtricitabine Efavirenz Mylan at bedtime on an empty stomach. They also tend to go away after you have taken the medicine for a few weeks. If you have these common side effects, such as dizziness, it does not mean that you will also have serious psychiatric problems, such as severe depression, strange thoughts or angry behaviour.

**Tell your doctor right away if any of these side effects continue or if they bother you.**

It is possible that these symptoms may be more severe if Tenofovir Disoproxil Emtricitabine Efavirenz Mylan is used with alcohol or mood altering (recreational) drugs.

Other common side effects may include:

- Rash

Rashes usually go away without any change in treatment. In a small number of patients, rash may be serious. If you develop a rash, call your doctor immediately.

- Tiredness
- Upset stomach
- Vomiting
- Gas and diarrhoea

Other possible side effects:

- Change in body fat. Changes in body fat develop in some people receiving anti-HIV-1 therapy. These changes may include an increased amount of fat in the upper back and neck ('buffalo hump'), in the breasts and around the trunk. Loss of fat from the legs, arms and face may also happen. The cause and long-term health effects of these fat changes are not known.
- Sleeping problems (including difficulty to fall asleep or sleepiness)
- Skin discolouration (small spots or freckles) may also occur with

Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.

**Tell your doctor or pharmacist if you notice any side effects while taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.**

**Contact your doctor before stopping Tenofovir Disoproxil Emtricitabine Efavirenz Mylan because of side effects or for any other reason.**

**Ask your doctor or pharmacist if you don't understand anything in this list.**

This is not a complete list of side effects possible with Tenofovir Disoproxil Emtricitabine Efavirenz Mylan.

**Ask your doctor or pharmacist for a more complete list of side effects of Tenofovir Disoproxil Emtricitabine Efavirenz Mylan and all the medicines you will take.**

**Do not be alarmed by this list of possible side effects. You may not experience any of them.**

---

## **After taking Tenofovir Disoproxil Emtricitabine Efavirenz Mylan**

---

### **Storage**

**Keep your tablets in the bottle until it is time to take them.**

If you take the tablets out of the bottle they may not keep well.

**Keep your tablets in a cool dry place where the temperature stays below 25°C.**

Do not store Tenofovir Disoproxil Emtricitabine Efavirenz Mylan or any other medicine in the bathroom or near a sink.

Do not leave it on a window sill or in the car.

Heat and dampness can destroy some medicines.

**Keep it where children cannot reach it.**

A locked cupboard at least one-and-

a half metres above the ground is a good place to store medicines.

## **Disposal**

**If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.**

---

## **General advice**

**Talk to your doctor or pharmacist if you have any questions about this medicine or your condition.**

Medicines are sometimes prescribed for conditions that are not mentioned in this leaflet.

**Do not use Tenofovir Disoproxil Emtricitabine Efavirenz Mylan for a condition for which it was not prescribed.**

**Do not give Tenofovir Disoproxil Emtricitabine Efavirenz Mylan to other people, even if they have the same symptoms that you have. It may harm them.**

This leaflet summarises the most important information about Tenofovir Disoproxil Emtricitabine Efavirenz Mylan. If you would like more information, ask your doctor or pharmacist. Your doctor or pharmacist can give you information about this medicine that was written doctors or pharmacists (Data sheet).

---

## **Product description**

### ***What it looks like***

Tenofovir Disoproxil Emtricitabine Efavirenz Mylan tablets are capsule shaped and pink in colour.

Each tablet is debossed with "M" on one side of the tablet and "TEE" on the other side.

Tenofovir Disoproxil Emtricitabine Efavirenz Mylan tablets are supplied in bottles containing 30 tablets.

## **Ingredients**

### Active ingredient(s):

Tenofovir Disoproxil Emtricitabine Efavirenz Mylan contains 300 mg of

tenofovir disoproxil maleate, 200 mg of emtricitabine and 600 mg of efavirenz as the active ingredient.

### Inactive ingredient(s):

Tenofovir Disoproxil Emtricitabine Efavirenz Mylan also contains:

- Croscarmellose sodium
- Hypolose
- Hydroxypropylcellulose
- Magnesium stearate
- Microcrystalline cellulose
- Sodium metabisulfite
- Lactose monohydrate
- Silica colloidal anhydrous
- Iron oxide red
- Opadry pink 85F540043

---

## **If you want to know more**

Should you have any questions regarding this product, please contact your pharmacist or doctor.

### ***Who supplies this medicine***

Tenofovir Disoproxil Emtricitabine Efavirenz Mylan is supplied in New Zealand by:

Mylan New Zealand Ltd,  
PO Box 11183,  
Ellerslie,  
Auckland  
NEW ZEALAND

Telephone: 0800 579 811

---

## **Date of Preparation**

1 June 2021  
(Based on datasheet dated 1 June 2021)