

CONSUMER MEDICINE INFORMATION

XAREMED (rivaroxaban)

Rivaroxaban 10 mg, 15 mg, 20 mg film coated tablets

What is in this leaflet

Please read this leaflet carefully before you start using XAREMED (rivaroxaban).

This leaflet answers some common questions about XAREMED (rivaroxaban). It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you using XAREMED (rivaroxaban) against the benefits they expect it will have for you.

If you have any concerns about using this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine. You may need to read it again.

What XAREMED (rivaroxaban) is used for

The active substance is rivaroxaban. It belongs to a group of medicines called anticoagulants. It works by inhibiting the blood clotting protein called Factor Xa, thus reducing the tendency of blood to form clots.

XAREMED (rivaroxaban) has been prescribed to you for one of the following uses:

- Prevention of blood clots in your veins after a hip or knee replacement operation because after an operation you are at an increased risk of getting blood clots.
- Prevention of blood clots in your brain (stroke) and/or other blood vessels in your body if you have a form of irregular heart rhythm called non-valvular atrial fibrillation
- Treatment of blood clots in the veins of your legs (deep vein thrombosis, DVT) and clots in your lung (pulmonary embolism, PE) and to prevent blood clots from re-occurring in your legs and/or lungs.

XAREMED (rivaroxaban) is a prescription medicine. It should only be used in adults under medical supervision.

Before you take XAREMED (rivaroxaban)

When you must not take it

Do not take XAREMED (rivaroxaban) if you have an allergy to:

- any medicine containing rivaroxaban
- any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin.

Do not take XAREMED (rivaroxaban):

- if you are bleeding excessively or at an increased risk of bleeding
- if you have liver disease which leads to an increased risk of bleeding
- if you have end stage kidney disease or if you are undergoing dialysis (a procedure used to remove waste products from the blood). Your doctor will know how to determine your kidney function.
- If you are taking medicines for fungal infections e.g. ketoconazole, or itraconazole, voriconazole, or posaconazole, unless they are only applied to the skin
- if you are taking anti-viral medicines for HIV/AIDS e.g. ritonavir
- if you had bleeding in the brain within the last 6 months

If you are not sure whether you should start using XAREMED (rivaroxaban), talk to your doctor.

Do not use this medicine if you are pregnant or intend to become pregnant.

Women should use a reliable contraceptive while taking XAREMED (rivaroxaban).

Do not use XAREMED (rivaroxaban) if you are breast-feeding.

It is not known whether rivaroxaban passes into human breast milk.

Tell your doctor if you have a mechanical heart valve

- XAREMED (rivaroxaban) may not be suitable for you because it has not been studied in patients with a mechanical heart valve.

This medicine should not be used in a child under the age of 18 years.

Do not take this medicine after the expiry date printed on the pack and blister.

The expiry date is printed on the carton and on each blister after “EXP” (e.g. 11 18 refers to November 2018). The expiry date refers to the last day of that month. If it has expired return it to your pharmacist for disposal.

Do not take this medicine if the packaging is torn or shows signs of tampering.

If the packaging is damaged, return it to your pharmacist for disposal.

Before you start to use it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have had any of the following medical conditions:

- if you have kidney disease. Special care should be taken in patients with moderate or severe kidney disease. Your doctor will know how to determine your kidney function.
- if you have kidney disease and undergoing dialysis (a procedure used to remove waste products from the blood)
- if you have liver disease
- if you have a prosthetic heart valve
- if you have an increased risk of bleeding such as:
 - bleeding disorders
 - very high blood pressure, not controlled by medical treatment
 - an active ulcer or a recent ulcer of your stomach or bowel
 - a problem with the blood vessels in the back of your eyes
 - recent bleeding in your brain
 - a recent operation on your brain, spinal column or eye
 - abnormalities in blood vessels of your spine or brain
 - a lung disease where your bronchi are widened (bronchiectasis), or history of bleeding in the lungs.
 - if a doctor has told you that you have a severe form of antiphospholipid syndrome (a disease which can cause blood clots).
 - you have a coronary artery disease with severe symptomatic heart failure
 - If you have CAD and/or PAD and had the following:
 - a bleed in your brain
 - a blood clot in your brain (ischaemic, nonlacunar stroke) in the previous month or
 - a blockage of the small arteries in the brain (lacunar stroke).
- If you have an active cancer – this may also mean you have an increased risk of bleeding. An active cancer means that in the last 6 months you:
 - have been diagnosed with cancer
 - had a relapse of cancer
 - were being treated for cancer

Your doctor may decide to keep you under closer observation.

In the event of a surgery

Tell your doctor, dentist or pharmacist if you need to have an operation (including dental work) while you are taking XAREMED (rivaroxaban). It is very important to take XAREMED (rivaroxaban) and any other medications you might be on, before and after the operation exactly at the times you have been told by your doctor.

During any invasive procedure or operation, if it involves a catheter or injection into your spinal column (e.g. for epidural or spinal anaesthesia or pain reduction):

- it is very important to take XAREMED (rivaroxaban) before and after the injection or removal of the catheter exactly at the times you have been told by your doctor
- tell your doctor immediately if you get numbness or weakness of your legs or problems with your bowel or bladder after the end of anaesthesia, because urgent care is necessary.

If you have not told your doctor about any of the above, tell them before you start taking XAREMED (rivaroxaban).

XAREMED (rivaroxaban) contains lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking it.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including medicines that you buy without a prescription from your pharmacy, supermarket or health food shop. You should also tell any health professional who is prescribing a new medication for you that you are taking XAREMED (rivaroxaban).

Some medicines may interfere with XAREMED (rivaroxaban). These include:

- other medicines to reduce blood clotting e.g. enoxaparin, clopidogrel or warfarin
- some medicines to treat depression (selective serotonin reuptake inhibitors (SSRIs) or serotonin norepinephrine reuptake inhibitors (SNRIs)).

These medicines may be affected by RIVAROXABAN, may increase the effect of RIVAROXABAN or may affect how well RIVAROXABAN works. You may need different amounts of your medicine, or you may need to take different medicines.

Your doctor or pharmacist has more information on medicines to be careful with or avoid while taking XAREMED (rivaroxaban).

Tell your doctor if you are taking anti-inflammatory and pain relieving medicines e.g. naproxen, or medicines used for the protection of heart disease e.g. acetylsalicylic acid (aspirin).

Your doctor may decide to keep you under closer observation. If your doctor thinks that you are at increased risk of developing stomach or bowel ulcers, he may also use a preventative ulcer treatment.

If you are taking

- medicines for treatment of epilepsy (phenytoin, carbamazepine)
- St John's Wort, an herbal product used for depression
- Rifampicin, an antibiotic

Tell your doctor before taking XAREMED (rivaroxaban), because its effect may be reduced.

How to take XAREMED (rivaroxaban)

Follow all directions given to you by your doctor or pharmacist carefully. They may differ from the information contained in this leaflet.

If you do not understand the instructions on the pharmacist label, ask your doctor or pharmacist for help.

How much to take

[Provide the necessary and usual instructions for proper use of the medicine]

To prevent clots in your veins after a hip or knee replacement operation, the dose is one rivaroxaban 10 mg tablet once a day with or without food.

To prevent blood clots in brain (stroke) and other blood vessels, the usual dose is one rivaroxaban 20 mg tablet once a day. If your kidneys are not working properly, your doctor may reduce your dose to one rivaroxaban 15 mg tablet once a day. The tablet packs are marked with days of the week to help you remember if you have taken your daily dose. rivaroxaban 15 mg and 20 mg tablets are to be taken with food.

To treat blood clots in your legs and clots in your lungs and for preventing blood clots from re-occurring, the usual dose is one rivaroxaban 15 mg tablet TWICE daily for the first three weeks. The initial treatment pack (42 tablet pack) is marked with days of the week and “am” for the morning dose and “pm” for the evening dose. This will help you remember if you have taken the required dose. After the first three weeks, take one rivaroxaban 20 mg tablet ONCE daily. After 6 to 12 months treatment your doctor may decide to continue treatment with either one rivaroxaban 20 mg tablet ONCE a day or one rivaroxaban 10 mg tablet ONCE a day. Rivaroxaban 15 mg and 20 mg tablets are to be taken with food.

If you have difficulty swallowing the tablet whole, talk to your doctor about other ways to take XAREMED (rivaroxaban). The tablet may be crushed and mixed with water or apple puree immediately before you take it. This drink should be immediately followed by food.

If necessary, the crushed XAREMED (rivaroxaban) tablet may be given to you through a stomach tube.

After giving the crushed XAREMED (rivaroxaban) tablet via the stomach tube, you or your carer should flush the tube with water. If you are taking the 15 or 20mg XAREMED (rivaroxaban) tablet, you should be fed via the stomach tube straight after your dose of XAREMED (rivaroxaban).

When to take it

Following hip or knee replacement operation: Take the first tablet 6 to 10 hours after your operation or as advised by your doctor. Then take a tablet every day for the duration prescribed, unless your doctor tells you to stop.

If you have had a hip replacement you will usually take the tablets for 5 weeks. If you have had a knee replacement you will usually take the tablets for 2 weeks. Your doctor will advise you about the exact duration.

For prevention of stroke or treatment or prevention of blood clots in your legs and/or lungs, take the tablet(s) every day until your doctor tells you to stop. Your doctor will decide how long you must continue your treatment.

If your heartbeat needs to be restored to normal by a procedure called cardioversion, take rivaroxaban according to your doctor's instructions.

If you need a procedure to treat blocked blood vessels in your heart (called a percutaneous coronary intervention – PCI with an insertion of a stent), the dose may be changed by your doctor. Your doctor will advise you about any changes to the amount of rivaroxaban you should take.

It is important that you follow instructions from your doctor and not to miss or stop taking your medicine or lower the dosage without checking with your doctor. rivaroxaban has been prescribed to you by your doctor to treat and/or prevent a serious medical condition.

Try to take the tablet(s) at the same time every day to help you remember.

If you forget to take it

If you are taking one rivaroxaban 10 mg, or one 15 mg, or one 20 mg tablet ONCE a day: If you have missed a dose, take it as soon as you remember. Do not take more than one tablet in a single day to make up for a forgotten dose.

Take the next tablet on the following day and then carry on taking a tablet once a day as normal. **Do not take a double dose to make up for a forgotten tablet.**

If you are taking one rivaroxaban 15 mg tablet TWICE a day and have missed a dose, take it as soon as you remember. If you forget to take a dose; you can take two rivaroxaban 15 mg tablets at the same time to get a total dose of 30 mg in one day. The following day onwards, you should take one rivaroxaban 15 mg tablet twice a day as normal, until required. you can take the next dose at the usual time. Do not take a double dose to make up for a missed dose.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

While you are taking XAREMED (rivaroxaban)

Things you must do

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking rivaroxaban.

Take rivaroxaban exactly as your doctor has prescribed.

Remember to carry your patient alert card in your wallet with you at all times.

Tell all doctors, dentists and pharmacists who are treating you that you are taking rivaroxaban.

Tell your doctor if you need to have a surgical or dental procedure.

Tell your doctor that you are using rivaroxaban, if your doctor is planning for you to have an anaesthetic injection in your back (spinal or epidural injection).

Tell your doctor if other medications are prescribed to you during the course of therapy with rivaroxaban.

If you become pregnant while you are taking rivaroxaban, immediately tell your doctor.

Things you must not do

Do not take XAREMED (rivaroxaban) to treat any other complaints unless your doctor tells you to.

Do not give XAREMED (rivaroxaban) to anyone else, even if they have the same condition as you.

Do not stop taking your medicine or lower the dosage without checking with your doctor first because XAREMED (rivaroxaban) treats and prevents serious conditions.

Things to be careful of

If this medicine makes you feel faint or dizzy, do not drive or use machinery.

In case of overdose

If you take too much (overdose)

Immediately telephone your doctor or the National Poisons Centre (telephone 0800 POISON or 0800 764 766) or go to accident and emergency at your nearest hospital, if you think that you or anyone else may have taken too much XAREMED (rivaroxaban).

Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

Taking too much XAREMED (rivaroxaban) increases the risk of bleeding.

Side Effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are using XAREMED (rivaroxaban).

All medicines can have side effects. Sometimes they are serious, most of the time they are not. In serious cases, you may need medical attention.

Do not be alarmed by the following lists of side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Your doctor may need to monitor and conduct blood tests, as XAREMED (rivaroxaban) can affect your liver or pancreatic enzymes. You may not experience any specific symptoms.

Like other similar medicines (anticoagulants), XAREMED (rivaroxaban) may cause bleeding, which may potentially be life threatening. In some cases this bleeding may not be obvious. There is no antidote available to reverse the effects of XAREMED (rivaroxaban), however there are measures your health professional can take to control/stop the bleeding. Please see your doctor if you experience any symptoms of bleeding.

If any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital:

- exceptional weakness, unexplained swelling
- breathlessness, chest pain
- signs of allergy such as rash, itching or hives on skin, swelling of the face, lips, tongue, or other parts of the body, shortness of breath, wheezing or trouble breathing
- signs of liver problems such as yellowing of the skin and/or eyes (jaundice)
- prolonged or excessive bleeding from gums, nose etc
- numbness in the arms and legs
- dizziness, fainting
- oozing from a surgical wound
- vomiting or coughing up blood
- blood in the urine or stool
- heavy menstrual bleeding
- skin condition with severe blisters and bleeding in the lips, eyes, mouth, nose and genitals
- extensive skin rash associated with fever
- blood in urine, reduced urine output, swelling of the ankles, feet and legs, increased time for blood to clot, and heavy bleeding. These symptoms may be related to a condition called anticoagulant-related nephropathy.

Your doctor may decide to keep you under observation or change how you should be treated.

Tell your doctor as soon as possible if you experience any of the following side effects and they worry you.

- tiredness, pale skin and breathlessness
- bruising
- feeling sick (nausea)
- headache
- diarrhoea, indigestion, or stomach pain
- pain in the arms or legs
- constipation
- fever
- frequent infections such as severe chills, sore throat and mouth ulcers

- mild rash, itchy skin
- fast heart beat

These side effects are usually mild.

Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

Other side effects not listed above may also occur in some people. If you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

After using XAREMED (rivaroxaban)

Storage

Keep your tablets in their blister pack until it is time to take them.

If you take the tablets out of the box or blister pack they may not keep well.

Store the tablets in a cool dry place where the temperature stays below 25°C.

Do not store it or any other medicine in the bathroom, near a sink, or on a window-sill.

Do not leave it in the car.

Heat and damp can destroy some medicines.

Keep out of the reach and sight of children. A locked cupboard at least one and a half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking the tablets, or the tablets have passed their expiry date, ask your pharmacist what to do with any that are left over.

Return any unused medicine to your pharmacist.

Product description

What it looks like

XAREMED (rivaroxaban) rivaroxaban 10 mg film-coated tablets

Light red colored, round, biconvex, film coated tablets, with engraving 10 on one face and other face plain with approximate dimension of 6.10 mm in diameter.

XAREMED (rivaroxaban) rivaroxaban 15 mg film-coated tablets

Red colored, round, biconvex, film coated tablets, with engraving 15 on one face and other face plain with approximate dimension of 7.10 mm in diameter.

XAREMED (rivaroxaban) rivaroxaban 20 mg film-coated tablets

Brown red colored, round, biconvex, film coated tablets, with engraving 20 on one face and other face plain with approximate dimension of 7.60 mm in diameter.

The tablets are packed in PVC/PCTFE (Aclar)/Aluminium blisters.

XAREMED (rivaroxaban) rivaroxaban 10 mg film-coated tablets

The tablets are supplied in packs of 15, 28 and 30 tablets*.

XAREMED (rivaroxaban) rivaroxaban 15 mg film-coated tablets

The tablets are supplied in packs of 28 and 42 tablets*.

XAREMED (rivaroxaban) rivaroxaban 20 mg film-coated tablets

The tablets are supplied in packs of 28 tablets*.

*Not all pack sizes may be marketed.

Ingredients

Active ingredient per tablet:

- XAREMED (rivaroxaban) 10 mg contains 10 mg rivaroxaban
- XAREMED (rivaroxaban) 15 mg contains 15 mg rivaroxaban
- XAREMED (rivaroxaban) 20 mg contains 20 mg rivaroxaban

Inactive ingredients:

- microcrystalline cellulose
- croscarmellose sodium
- hypromellose
- lactose monohydrate
- magnesium stearate
- sodium lauryl sulfate
- iron oxide red (10 mg, 15 mg and 20 mg tablets)

Sponsor Details

XAREMED (rivaroxaban) is supplied in New Zealand by:

Medsurge Pharma Limited
PO Box 331054
Takapuna
Auckland 0622
Telephone: 0800 788 261

Marketed and Distributed by Medsurge Healthcare.



Telephone: 1300 788 261
Website: www.medsurgehc.com

Date of Preparation

This leaflet was June 2025