

NEW ZEALAND DATA SHEET

1. PRODUCT NAME

DBL™ Irinotecan 20 mg/mL Injection Concentrate

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

The 2 mL, 5 mL and 25mL injections contain 40 mg, 100 mg and 500mg of irinotecan hydrochloride trihydrate respectively.

Excipient(s) with known effect

- Lactic acid
- Sodium hydroxide
- Sorbitol

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for injection

Irinotecan Injection Concentrate is supplied as a sterile, pale yellow, clear, aqueous solution of pH 3.5. It is intended for dilution with 5% Glucose Injection or 0.9% Sodium Chloride Injection prior to infusion.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

DBL Irinotecan Injection Concentrate is indicated as a component of first line treatment of patients with metastatic carcinoma of the colon or rectum. Irinotecan Injection Concentrate is also indicated for patients with metastatic carcinoma of the colon or rectum whose disease has recurred or progressed following initial therapy.

4.2 Dose and method of administration

It is recommended that patients receive premedication with anti-emetic agents. Prophylactic or therapeutic administration of atropine should be considered in patients experiencing cholinergic symptoms (see section 4.4).

Combination therapy in first line treatment of metastatic colorectal cancer

Irinotecan Injection Concentrate should be administered as an intravenous infusion (IV) over 90 minutes (see section 6.6). For all regimens, the dose of LV should be administered immediately after Irinotecan Injection Concentrate, with the administration of 5-FU to follow

immediately after the administration of LV. The recommended regimens are shown in Table 1.

Table 1: Combination Agent Dosage Regimens and Dose Modifications^a

Regimen 1 6 week cycle Treatment resumes Day 43	Irinotecan hydrochloride	125 mg/m ² IV over 90 min on Day 1, 8, 15, 22 then 2 wk rest		
	LV	20 mg/m ² IV bolus injection Day 1, 8, 15, 22 then 2 wk rest		
	5-FU	500 mg/m ² IV bolus injection Day 1, 8, 15, 22 then 2 wk rest		
		Starting dose and modified dose levels^b		
		Starting dose (mg/m ²)	Dose level -1 (mg/m ²)	Dose Level -2 (mg/m ²)
	Irinotecan hydrochloride	125	100	75
	LV	20	20	20
	5-FU	500	400	300
Regimen 2 6 week cycle Treatment resumes Day 43	Irinotecan hydrochloride	180 mg/m ² IV over 90 min on Day 1, 15, 29 then 1 wk rest		
	LV	200 mg/m ² IV over 2 h on Day 1, 2, 15, 16, 29, 30 then 1 wk rest		
	5-FU Bolus	400 mg/m ² IV on Day 1, 2, 15, 16, 29, 30 then 1 wk rest		
	5-FU Infusion ^c	600 mg/m ² IV over 22h on Day 1, 2, 15, 16, 29, 30 then 1 wk rest		
		Starting dose and modified dose levels^b		
		Starting dose (mg/m ²)	Dose level -1 (mg/m ²)	Dose Level -2 (mg/m ²)
	Irinotecan hydrochloride	180	150	120
	LV	200	200	200
	5-FU Bolus	400	320	240
	5-FU Infusion ^c	600	480	360

^a Dose reductions beyond dose level -2 by decrements of ~20% may be warranted for patients continuing to experience toxicity. Provided intolerable toxicity does not develop, treatment with additional cycles may be continued indefinitely as long as patients continue to experience clinical benefit.

^b Refer to Table 2

^c Infusion follows bolus administration

Dosing for patients with bilirubin >34 mmol/L cannot be recommended since such patients were not included in clinical trials.

Irinotecan in combination with Cetuximab

For dosage and administration of concomitant cetuximab, refer to the full prescribing information for cetuximab. Normally, the same dose of irinotecan is used as administered in the last cycles of the prior irinotecan-containing regimen. Irinotecan must not be administered earlier than 1 hour after the end of the cetuximab infusion.

Dose modifications

Patients should be carefully monitored for toxicity and assessed prior to each treatment, especially during the first cycle of therapy. Doses of Irinotecan Injection Concentrate and 5-FU should be modified as necessary to accommodate individual patient tolerance to treatment. Based on the recommended dose levels described in Table 1, subsequent doses should be

adjusted as suggested in Table 2, which shows the recommended dose modifications for combination schedules. All dose modifications should be based on the worst preceding toxicity. Patients should be diarrhoea free (return to pre treatment bowel function) without requiring antidiarrhoeal medications for at least 24 hours before receiving the next chemotherapy administration.

A new cycle of therapy should not begin until the toxicity has recovered to NCI grade 1 or less, the granulocyte count has recovered to $\geq 1.5 \times 10^9/L$, the platelet count has recovered to $\geq 100 \times 10^9/L$ and treatment-related diarrhoea is fully resolved. Treatment should be delayed for 1 to 2 weeks to allow recovery from treatment-related toxicity. If the patient has not recovered after a 2 week delay, consideration should be given to discontinuing therapy. Provided intolerable toxicity does not develop, treatment with additional cycles of Irinotecan Injection Concentrate/5-FU/LV may be continued indefinitely as long as patients continue to experience clinical benefit.

Table 2: Recommended dose modifications during a cycle of therapy with the Irinotecan Injection Concentrate/5-FU/LV combination and at the start of each subsequent cycle of therapy

Toxicity NCI CTC Grade ^a	During a Cycle of Therapy	At the Start of Subsequent Cycles of Therapy ^b
No toxicity	Maintain dose level	Maintain dose level
Neutropenia		
1	Maintain dose level	Maintain dose level
2	Decrease by 1 dose level	Maintain dose level
3	Omit dose until resolved to \leq grade 2, then decrease by 1 dose level	Decrease by 1 dose level
4	Omit dose until resolved to \leq grade 2, then decrease by 2 dose levels	Decrease by 2 dose levels
Neutropenic fever	Omit dose until resolved, then decrease by 2 dose levels	
Other haematological toxicities	Dose modifications for leucopenia or thrombocytopenia during a cycle of therapy and at the start of subsequent cycles of therapy are also based on NCI toxicity criteria, and are the same as recommended for neutropenia above.	
Diarrhoea		
1	Delay dose until resolved to baseline, then give same dose	Maintain dose level
2	Omit dose until resolved to baseline, then decrease by 1 dose level	Maintain dose level
3	Omit dose until resolved to baseline, then decrease by 1 dose level	Decrease by 1 dose level
4	Omit dose until resolved to baseline, then decrease by 2 dose levels	Decrease by 2 dose levels
Other nonhaematological toxicities^c		
1	Maintain dose level	Maintain dose level
2	Omit dose until resolved to \leq grade 1, then decrease by 1 dose level	Maintain dose level

Toxicity NCI CTC Grade ^a	During a Cycle of Therapy	At the Start of Subsequent Cycles of Therapy ^b
3	Omit dose until resolved to \leq grade 2, then decrease by 1 dose level	Decrease by 1 dose level
4	Omit dose until resolved to \leq grade 2, then decrease by 2 dose levels	Decrease by 2 dose levels

^a Severity of adverse events based on NCI CTC (version 2.0)

^b Relative to the starting dose used in the previous cycle

^c For mucositis/stomatitis decrease only 5-FU, not Irinotecan Injection Concentrate

Single agent therapy in recurrent or progressive metastatic colorectal cancer

Irinotecan Injection Concentrate should be administered as an intravenous infusion (see section 6.6) over 90 minutes in a recommended weekly or once every 3 week dosage schedule as shown below in Table 3.

Table 3: Single-Agent Regimens of Irinotecan Injection Concentrate and Dose Modifications

Weekly Regimen^a 6 week cycle Treatment resumes Day 43	125 mg/m ² IV over 90 mins Day 1, 8, 15, 22 then 2 week rest		
	Starting dose and modified dose levels^c		
	Starting Dose (mg/m ²)	Dose Level –1 (mg/m ²)	Dose Level –2 (mg/m ²)
	125	100	75
Once every 3 week regimen^b	350 mg/m ² IV over 90 mins once every 3 weeks		
	Starting dose and modified dose levels^c		
	Starting Dose (mg/m ²)	Dose Level –1 (mg/m ²)	Dose Level –2 (mg/m ²)
	350	300	250

^a Subsequent doses may be adjusted as high as 150 mg/m² or as low as 50 mg/m² in 25 to 50 mg/m² decrements depending on individual patient tolerance

^b Subsequent doses may be adjusted as low as 200 mg/m² in 50 mg/m² decrements depending on individual patient tolerance

^c Refer to Table 5

A reduction in the starting dose by one level of Irinotecan Injection Concentrate may be considered for patients with any of the following circumstances: over 65 years, prior pelvic/abdominal radiotherapy, performance status of 2 or moderately increased bilirubin levels (17 - 34 μ mol/L). Dosing for patients with bilirubin >34 mmol/L cannot be recommended since such patients were not included in clinical trials.

Patients with Hepatic Impairment (Single Agent)

In patients with hepatic dysfunction, the following starting doses are recommended:

Table 4: Starting Doses in Patients with Hepatic Dysfunction – Single Agent Regimens

Regimen	Serum Total Bilirubin Concentration	Serum ALT/AST Concentration	Starting Dose, mg/m ²
Single Agent Weekly	1.5–3.0 x IULN	≤5.0 x IULN	60
	3.1–5.0 x IULN	≤5.0 x IULN	50
	<1.5 x IULN	5.1–20.0 x IULN	60
	1.5–5.0 x IULN	5.1–20.0 x IULN	40
Single Agent Once Every 3 Weeks	1.5–3.0 x IULN	-	200
	>3.0 x IULN	-	Not recommended ^a

^a The safety and pharmacokinetics of IRINOTECAN given once every 3 weeks have not been defined in patients with bilirubin >3.0 x IULN and this schedule cannot be recommended in these patients.

Dose modifications

Patients should be carefully monitored for toxicity and doses of Irinotecan Injection Concentrate should be modified as necessary to accommodate individual patient tolerance to treatment. Based on recommended dose-levels described in Tables 3 and 4, subsequent doses of Irinotecan Injection Concentrate should be adjusted as suggested in Table 5. All dose modifications should be based on the worst preceding toxicity.

A new cycle of therapy should not begin until the toxicity has recovered to NCI grade 1 or less, the granulocyte count has recovered to $\geq 1.5 \times 10^9/L$, the platelet count has recovered to $\geq 100 \times 10^9/L$ and treatment-related diarrhoea is fully resolved. Treatment may be delayed for 1 to 2 weeks to allow recovery from treatment-related toxicity. If the patient has not recovered, consideration should be given to discontinuing Irinotecan Injection Concentrate therapy. Provided intolerable toxicity does not develop, treatment with additional cycles of Irinotecan Injection Concentrate may be continued indefinitely as long as patients continue to experience clinical benefit.

Table 5: Recommended dose modifications during a cycle of therapy with the weekly dosage schedule and at the start of each subsequent cycle of therapy with both dosage schedules

Toxicity NCI ^a Grade	During a cycle of therapy	At the start of subsequent cycles of therapy	
	Weekly	Weekly	Once every 3 weeks
No toxicity	Maintain dose level	Increase by 1 dose level up to a maximum dose of 150 mg/m ²	Maintain dose level
Neutropenia			
1	Maintain dose level	Maintain dose level	Maintain dose level
2	Decrease by 1 dose level	Maintain dose level	Maintain dose level
3	Omit dose until resolved to ≤ grade 2, then decrease by 1 dose level	Decrease by 1 dose level	Decrease by 1 dose level
4	Omit dose until resolved to ≤ grade 2, then decrease by 2 dose levels	Decrease by 2 dose levels	Decrease by 1 dose level

Neutropenic fever	Omit dose until resolved, then decrease by 2 dose levels	Decrease by 2 dose levels	Decrease by 1 dose level
Other haematological toxicities	Dose modifications for leucopenia, thrombocytopenia and anaemia during a cycle of therapy and at the start of subsequent cycles of therapy are also based on NCI toxicity criteria and are the same as recommended for neutropenia above.		
Diarrhoea			
1	Maintain dose level	Maintain dose level	Maintain dose level
2	Decrease by 1 dose level	Maintain dose level	Maintain dose level
3	Omit dose until resolved to \leq grade 2, then decrease by 1 dose level	Decrease by 1 dose level	Decrease by 1 dose level
4	Omit dose until resolved to \leq grade 2, then decrease by 2 dose levels	Decrease by 2 dose levels	Decrease by 1 dose level
Other non-haematological toxicities			
1	Maintain dose level	Maintain dose level	Maintain dose level
2	Decrease by 1 dose level	Decrease by 1 dose level	Decrease by 1 dose level
3	Omit dose until resolved to \leq grade 2, then decrease by 1 dose level	Decrease by 1 dose level	Decrease by 1 dose level
4	Omit dose until resolved to \leq grade 2, then decrease by 2 dose levels	Decrease by 2 dose levels	Decrease by 1 dose level

^a Severity of adverse events based on NCI CTC (version 2.0)

4.3 Contraindications

Irinotecan Injection Concentrate is contraindicated in patients with a known hypersensitivity to the drug or its excipients. Irinotecan Injection Concentrate antigenicity has not been observed in clinical trials, but irinotecan hydrochloride antigenicity occurred in tests for passive cutaneous anaphylaxis in guinea pigs and rabbits, and in tests for active systemic anaphylaxis in guinea pigs. In these tests, both animal species produced antibodies against irinotecan hydrochloride, and some deaths occurred in guinea pigs sensitised to irinotecan hydrochloride.

Irinotecan Injection Concentrate is contraindicated in women who intend to become pregnant (see section 4.6 and section 5.3).

Irinotecan Injection Concentrate is contraindicated in pregnancy and lactation (see section 4.6).

4.4 Special warnings and precautions for use

Administration: Irinotecan Injection Concentrate should be administered only under the supervision of a physician who is experienced in the use of cancer chemotherapeutic agents. Appropriate management of complications is possible only when adequate diagnostic and treatment facilities are readily available.

Extravasation: Irinotecan Injection Concentrate is administered by intravenous infusion. Care should be taken to avoid extravasation and the infusion site should be monitored for signs of inflammation. Should extravasation occur, flushing the site with sterile water and application of ice are recommended.

Mayo clinic regimen: Except in a well-designed clinical study, Irinotecan Injection Concentrate should not be used in combination with the "Mayo Clinic" regimen of 5-FU/LV (administration for 4-5 consecutive days every 4 weeks; refer to Table 9) because of reports of increased toxicity, including toxic deaths. Irinotecan Injection Concentrate should be used as recommended in section 4.2.

Immunosuppressant effects/Increased susceptibility to infections: Administration of live or live-attenuated vaccines in patients immunocompromised by chemotherapeutic agents, including irinotecan, may result in serious or fatal infections. Vaccination with a live vaccine should be avoided in patients receiving irinotecan. Killed or inactivated vaccines may be administered; however, the response to such vaccines may be diminished.

Cardiovascular: Thromboembolic events including angina pectoris, arterial thrombosis, cerebral infarct, cerebrovascular accident, deep thrombophlebitis, embolus lower extremity, heart arrest, myocardial infarct, myocardial ischaemia, peripheral vascular disorder, pulmonary embolus, sudden death, thrombophlebitis, thrombosis and vascular disorder have been observed rarely in patients receiving irinotecan hydrochloride. The specific cause of these events has not been determined (see section 4.8).

Use with caution in the following circumstances

Patients at particular risk: Physicians should exercise particular caution in monitoring the effects of irinotecan hydrochloride in patients with poor performance status. Patients with poor performance status are at increased risk of irinotecan-related adverse events. In patients receiving either irinotecan hydrochloride/ 5-FU/LV or 5-FU/LV in clinical trials comparing these agents, higher rates of hospitalisation, neutropenic fever, thromboembolism, first-cycle treatment discontinuation and early deaths were observed in patients with a baseline performance status of 2 than in patients with a baseline performance of 0 or 1. Patients with performance status of 3 or 4 should not receive irinotecan hydrochloride.

Physicians should exercise particular caution in monitoring the effects of irinotecan hydrochloride in elderly patients and in patients who have previously received pelvic/abdominal irradiation (see section 4.8).

Irradiation therapy: Patients who have previously received pelvic/abdominal irradiation are at increased risk of severe myelosuppression following the administration of irinotecan hydrochloride. The concurrent administration with irradiation has not been adequately studied and is not recommended.

Cholinergic effects: Irinotecan hydrochloride has cholinergic effects and should be used with caution in patients with asthma or cardiovascular diseases, and in patients with mechanical intestinal or urinary obstruction.

Respiratory: Interstitial pulmonary disease presenting as pulmonary infiltrates is uncommon during irinotecan therapy. Interstitial pulmonary disease can be fatal. Risk factors possibly associated with the development of interstitial pulmonary disease include pre-existing lung disease, use of pneumotoxic drugs, radiation therapy and colony stimulating factors. Patients with risk factors should be closely monitored for respiratory symptoms before and during irinotecan therapy.

Before administration

Monitoring: Careful monitoring of the white blood cell count with differential, haemoglobin and platelet count is recommended before each dose of irinotecan hydrochloride. Liver function should be monitored before initiation of treatment and monthly or as clinically indicated.

Nausea and vomiting: Irinotecan hydrochloride is emetogenic. It is recommended that patients receive premedication with anti-emetic agents. In clinical studies with the weekly dosage schedule, the majority of patients received 10 mg of dexamethasone given in conjunction with another type of anti-emetic agent, such as a 5-HT₃ blocker (e.g. ondansetron or granisetron). Anti-emetic agents should be given on the day of treatment, starting at least 30 minutes before administration of Irinotecan Injection Concentrate. Physicians should also consider providing patients with an anti-emetic regimen (e.g. prochlorperazine) for subsequent use as needed. Patients with vomiting associated with delayed (i.e. late) diarrhoea should be hospitalized as soon as possible for treatment.

Advice to patients: Patients should be advised of the expected toxic effects of irinotecan hydrochloride, particularly of gastrointestinal complications such as nausea, vomiting, abdominal cramping, diarrhoea and infection. Each patient should be instructed to have loperamide readily available and to begin treatment for late diarrhoea (generally occurring more than 24 hours after administration of irinotecan hydrochloride) at the first episode of poorly formed or loose stools, or the earliest onset of bowel movements more frequent than normally expected for the patient (see section 4.4).

Patients should be advised to consult their physician if any of the following occur after treatment with Irinotecan Injection Concentrate: diarrhoea for the first time; inability to control diarrhoea within 24 hours; vomiting; fever or evidence of infection; symptoms of dehydration, such as faintness, light-headedness or dizziness; bloody or black stools; inability to take fluids by mouth due to nausea or vomiting. Patients should also be alerted to the possibility of alopecia. Laxatives should be avoided (see section 4.5) and patients should contact their physician to discuss any laxative use.

Others

As this product contains sorbitol, it is unsuitable in hereditary fructose intolerance.

Diarrhoea and its Management

Irinotecan hydrochloride can induce both an early and a late form of diarrhoea that appear to be mediated by different mechanisms. Both forms of diarrhoea may be severe.

Early diarrhoea (occurring during or shortly after infusion of irinotecan hydrochloride) is cholinergic in nature. It is usually transient and only infrequently is severe. It may be accompanied by symptoms of rhinitis, increased salivation, miosis, lacrimation, diaphoresis, flushing, bradycardia and intestinal hyperperistalsis that can cause abdominal cramping. Administration of 0.25 to 1 mg of intravenous or subcutaneous atropine should be considered (unless clinically contraindicated) in patients experiencing cholinergic symptoms occurring during or shortly after infusion of irinotecan hydrochloride. Patients ≥ 65 years of age should be closely monitored due to a greater risk of early diarrhoea observed in this population.

Late diarrhoea (generally occurring more than 24 hours after administration of irinotecan hydrochloride) can be prolonged, may lead to dehydration, electrolyte imbalance or infection and can be life-threatening. Late diarrhoea should be treated promptly with loperamide. Patients should be instructed to have loperamide readily available and begin treatment at the first episode of poorly formed or loose stools, or the earliest onset of bowel movements more frequent than normally expected for the patient. One dosage regimen for loperamide used in clinical trials consisted of 4 mg at the first onset of late diarrhoea, and then 2 mg every 2 hours until the patient was diarrhoea-free for at least 12 hours. During the night, the patient may take 4 mg of loperamide every 4 hours. Loperamide is not recommended to be used for more than 48 consecutive hours at these doses, because of the risk of paralytic ileus, nor for less than 12 hours. Premedication with loperamide is not recommended.

Patients with diarrhoea should be carefully monitored and given fluid and electrolyte replacement if they become dehydrated and should be given antibiotics if they develop ileus, fever or severe neutropenia. After the first treatment, subsequent chemotherapy should be delayed until patients are diarrhoea-free (return to pre-treatment bowel function) for at least 24 hours without the need for antidiarrhoea medication. If NCI grade 2, 3 or 4 diarrhoea occurs, subsequent doses of irinotecan hydrochloride should be reduced within the current cycle (see section 4.2).

In addition to antibiotic treatment, hospitalization is recommended for management of the diarrhoea, in the following cases: diarrhoea associated with fever, severe diarrhoea (requiring intravenous hydration), patients with vomiting associated with delayed (i.e., late) diarrhoea and diarrhoea persisting beyond 48 hours following the initiation of high-dose loperamide therapy and in the few rare instances where patients are deemed unlikely to observe recommendations regarding management of adverse events (need for immediate and prolonged antidiarrhoeal treatment combined with high fluid intake at onset of delayed diarrhoea).

Haematology

Irinotecan commonly causes neutropenia, leucopenia and anaemia, any of which may be severe and therefore should not be used in patients with severe bone marrow failure (refer to section 4.8). Serious thrombocytopenia is uncommon.

Neutropenia: Deaths due to sepsis following severe neutropenia have been reported in patients treated with irinotecan hydrochloride. Neutropenic complications should be managed promptly with antibiotic support. Therapy with irinotecan hydrochloride should be temporarily omitted if neutropenic fever occurs or if the absolute neutrophil count drops below $1.5 \times 10^9/L$. A new cycle of therapy should not begin until the granulocyte count has recovered to $\geq 1.5 \times 10^9/L$. After the patient recovers, subsequent doses of irinotecan hydrochloride should be reduced depending upon the level of neutropenia observed (see section 4.2). Routine administration of a colony-stimulating factor (CSF) is not necessary but physicians may consider CSF use in individual patients experiencing problems related to neutropenia.

Hypersensitivity

Hypersensitivity reactions including severe anaphylactic and anaphylactoid reactions have been observed.

Colitis/Ileus

Cases of colitis have been reported. In some cases, colitis was complicated by ulceration, bleeding, ileus and infection. Cases of ileus without preceding colitis have also been reported. Patients experiencing ileus should receive prompt antibiotic support.

Chronic inflammatory bowel disease and/or bowel obstruction

Patients must not be treated with irinotecan hydrochloride until resolution of the bowel obstruction.

Patients with Reduced UGT1A1 Activity

Uridine diphosphate-glucuronosyl transferase 1A1 (UGT1A1), which mediates the conjugation of the active metabolite SN-38 (see section 5.2) is encoded by the UGT1A1 gene. This gene is highly polymorphic resulting in variable metabolic capacities among individuals. One specific variation of the UGT1A1 gene includes a polymorphism in the promoter region known as the UGT1A1*28 variant allele. This variant and other congenital deficiencies in UGT1A1 expression (such as Crigler-Najjar and Gilbert's syndrome) are associated with reduced enzyme activity and increased systemic exposure to SN-38. Higher plasma concentrations of SN-38 are observed in individuals who are homozygous for the UGT1A1*28 allele (also referred to as UGT1A1 7/7 genotype) compared to patients who have one or two wild-type alleles.

Individuals with Crigler-Najjar syndrome (types 1 and 2) or those homozygous for the UGT1A1*28 allele (Gilbert's syndrome) are at increased risk of haematological toxicity (grades 3 and 4) following administration of irinotecan hydrochloride at moderate or high doses (>150 mg/m²). A relationship between UGT1A1 genotype and the occurrence of irinotecan hydrochloride induced diarrhoea has not been established.

Patients known to be homozygous for UGT1A1*28 should be administered the normally-indicated irinotecan starting dose and monitored for haematologic toxicities. A reduced irinotecan hydrochloride starting dose should be considered for patients who have experienced haematologic toxicity with previous treatment. The exact reduction in starting dose in this patient population has not been established and any subsequent dose modifications should be based on individual patient tolerance to treatment.

Renal impairment

Studies in patients with impaired renal function have not been conducted (refer to section 5.2). Therefore, caution should be undertaken in patients with impaired renal function. Irinotecan is not recommended for use in patients on dialysis.

Hepatic impairment

In patients with hyperbilirubinemia, the clearance of irinotecan is decreased and therefore the risk of haematotoxicity is increased (refer to section 5.2).

The use of irinotecan hydrochloride in patients with a serum bilirubin concentration of >3.0 x institutional upper limit of normal (IULN) given as a single agent on the once every 3 weeks schedule has not been established. In clinical trials of the single agent weekly dosage schedule, it has been noted that patients with even modest elevations in total baseline serum bilirubin levels (17-34 µmol/L) have had a significantly greater likelihood of experiencing first-cycle grade 3 or 4 neutropenia than those with bilirubin levels that were less than 17 µmol/L (50%

versus 17.7%; $p < 0.001$) (refer to section 5.2 and section 4.2). Patients with deficient glucuronidation of bilirubin, such as those with Gilbert's syndrome, may also be at greater risk of myelosuppression when receiving therapy with irinotecan hydrochloride.

Use in the elderly

Physicians should exercise particular caution in monitoring the effects of irinotecan hydrochloride in elderly patients. A reduction in the starting dose of Irinotecan Injection Concentrate may be considered for patients over 65 years of age (see section 4.2).

Paediatric population

The safety and effectiveness of irinotecan hydrochloride in children have not been established.

Effects on laboratory tests

There are no known interactions between irinotecan hydrochloride and laboratory tests.

4.5 Interaction with other medicines and other forms of interaction

CYP3A4 and/or UGT1A1 Inhibitors

Irinotecan and its active metabolite SN-38 are metabolised via the human cytochrome P450 3A4 isoenzyme (CYP3A4) and uridine diphosphate-glucuronosyl transferase 1A1 (UGT1A1) (see section 5.2). Coadministration of irinotecan with inhibitors of CYP3A4 and/or UGT1A1 may result in increased systemic exposure to irinotecan and the active metabolite SN-38. Physicians should take this into consideration when administering irinotecan with these drugs.

Ketoconazole: Irinotecan clearance is greatly reduced in patients receiving concomitant ketoconazole, leading to increased exposure to the active metabolite SN-38. Ketoconazole should be discontinued at least 1 week prior to starting irinotecan therapy and should not be administered during irinotecan therapy.

Atazanavir sulfate: Co-administration of atazanavir sulfate, a CYP3A4 and UGT1A1 inhibitor has the potential to increase systemic exposure to SN-38, the active metabolite of irinotecan. Physicians should take this into consideration when co-administering these medicines.

CYP3A4 Inducers

Anticonvulsants: Concomitant administration of CYP3A-inducing anticonvulsant drugs (e.g. carbamazepine, Phenobarbital or phenytoin) leads to reduced exposure to SN-38. Consideration should be given to starting or substituting non-enzyme –inducing anticonvulsants at least one week prior to initiation of irinotecan therapy in patients requiring anticonvulsant treatment.

St. John's Wort (Hypericum perforatum): Exposure to the active metabolite SN-38 is reduced in patients taking concomitant St. John's Wort. St John's Wort should be discontinued at least 1 week prior to the first cycle of irinotecan, and should not be administered during irinotecan therapy.

Other Interactions

Neuromuscular blocking agents: Interaction between irinotecan and neuromuscular blocking agents cannot be ruled out. Since irinotecan has anticholinesterase activity, drugs with anticholinesterase activity may prolong the neuromuscular blocking effects of suxamethonium and the neuromuscular blockade of non-depolarizing drugs may be antagonized.

Antineoplastic agents: Possible pharmacokinetic interactions of irinotecan hydrochloride with other concomitantly administered medications have not been formally investigated. The adverse effects of irinotecan hydrochloride, such as myelosuppression and diarrhoea, would be expected to be exacerbated by other antineoplastic agents having similar adverse events.

Dexamethasone: Lymphocytopenia has been reported in patients receiving irinotecan hydrochloride and it is possible that the administration of dexamethasone as anti-emetic prophylaxis may have enhanced the likelihood of this effect. However, serious opportunistic infections have not been observed and no complications have specifically been attributed to the lymphocytopenia.

Hyperglycaemia has also been reported in patients receiving irinotecan hydrochloride. Usually this has been observed in patients with a history of diabetes mellitus or evidence of glucose intolerance prior to administration of irinotecan hydrochloride. It is probable that the administration of dexamethasone contributed to hyperglycaemia in some patients.

Prochlorperazine: The incidence of akathisia in clinical trials of the single agent weekly dosage schedule was greater (8.5%, 4/47 patients) when prochlorperazine was administered on the same day as irinotecan hydrochloride than when these drugs were given on separate days (1.3%, 1/80 patients). However, the 8.5% incidence of akathisia is within the range reported for use of prochlorperazine when given as a premedication for other chemotherapies.

Laxatives: It would be expected that the incidence or severity of diarrhoea would be worsened by laxative use during therapy with irinotecan hydrochloride, but this has not been studied.

Diuretics: In view of the potential risk of dehydration secondary to vomiting and/or diarrhoea induced by irinotecan hydrochloride, the physician may wish to withhold diuretics during dosing with irinotecan hydrochloride and, certainly, during periods of active vomiting or diarrhoea.

4.6 Fertility, pregnancy and lactation

Fertility

No significant adverse effects on fertility and general reproductive performance were observed after intravenous administration of irinotecan hydrochloride in doses of up to 6 mg/kg/day to rats. Atrophy of male reproductive organs was observed after multiple daily irinotecan hydrochloride doses both in rodents at 20 mg/kg (AUC approximately the same value as in patients administered 125 mg/m² weekly) and dogs at 0.4 mg/kg (AUC about 1/15th the value in patients administered 125 mg/m² weekly).

Pregnancy - Category D

There are no adequate and well-controlled studies of irinotecan in pregnant women. Irinotecan hydrochloride may cause foetal harm when administered to a pregnant woman. Administration of 6 mg/kg/day intravenous irinotecan hydrochloride to rats (AUC about 0.2 times the

corresponding values in patients administered 125 mg/m²) and rabbits (about one-half the recommended human weekly starting dose on a mg/m² basis) during the period of organogenesis, is embryotoxic as characterised by increased post-implantation loss and decreased numbers of live foetuses. Irinotecan hydrochloride was teratogenic in rats at doses greater than 1.2 mg/kg/day (AUC about 1/40th the corresponding values in patients administered 125 mg/m²) and in rabbits at 6.0 mg/kg/day. Teratogenic effects included a variety of external, visceral, and skeletal abnormalities.

Women of childbearing potential should not be started on irinotecan until pregnancy is excluded. Pregnancy should be avoided if either partner is receiving irinotecan.

Due to the potential for genotoxicity, advise female patients of reproductive potential to use highly effective contraception during treatment and for 6 months after the last dose of irinotecan.

Due to the potential for genotoxicity, advise male patients with female partners of reproductive potential to use effective contraception during treatment and for 3 months after the last dose of irinotecan.

Lactation

The available data are limited to one patient only. Irinotecan and its active metabolite SN-38 were measured in the milk of one lactating patient. The effect on newborn/infants is unknown. Because of the potential for serious adverse reactions in nursing infants, it is recommended not to breastfeed when receiving therapy with irinotecan.

Radioactivity appeared in rat milk within 5 minutes of intravenous administration of radiolabelled irinotecan hydrochloride and was concentrated up to 65-fold at 4 hours after administration relative to plasma concentrations. Irinotecan hydrochloride has been shown to impair learning ability and cause a delay in postnatal development in rats.

4.7 Effects on ability to drive and use machinery

The effect of irinotecan on the ability to drive or use machinery has not been evaluated. However, patients should be warned about the potential for dizziness or visual disturbances which may occur within 24 hours following the administration of irinotecan, and advised not to drive or operate machinery if these symptoms occur.

4.8 Undesirable effects

Combination therapy in first line treatment of metastatic colorectal cancer

In the two phase III studies, a total of 955 patients with metastatic colorectal cancer received irinotecan hydrochloride in combination with 5-FU/LV, 5-FU/LV alone, or irinotecan hydrochloride alone (see Table 9, section 5.1). In these studies, 370 patients received irinotecan hydrochloride in combination with 5-FU/LV, 362 patients received 5-FU/LV alone, and 223 patients received irinotecan hydrochloride alone.

Fifty-nine (6.1%) patients died within 30 days of last study treatment: 27 (7.3%) received irinotecan hydrochloride in combination with 5-FU/LV, 19 (5.3%) received 5-FU/LV alone,

and 13 (5.8%) received irinotecan hydrochloride alone. Deaths potentially related to treatment occurred in 3 (0.7%) patients who received irinotecan hydrochloride in combination with 5-FU/LV (2 neutropenic fever/sepsis, 1 treatment toxicity), 3 (0.7%) patients who received 5-FU/LV alone (1 neutropenic fever/sepsis, 1 CNS bleeding during thrombocytopenia, 1 unknown) and 2 (0.9%) patients who received irinotecan hydrochloride alone (2 neutropenic fever). Deaths within 60 days of study treatment were reported for 18 (4.9%) patients who received irinotecan hydrochloride in combination with 5-FU/LV, 18 (5.0%) patients who received 5-FU/LV alone and 15 (6.7%) patients who received irinotecan hydrochloride alone. Discontinuations due to adverse events were reported for 26 (7.0%) patients who received irinotecan hydrochloride in combination with 5-FU/LV, 15 (4.1%) patients who received 5-FU/LV alone, and 26 (11.7%) patients who received irinotecan hydrochloride alone.

Table 6 lists the grade 3 and 4 clinically relevant adverse events reported in the combination treatment arms of the two phase III studies.

Table 6: Percent (%) of Patients Experiencing Clinically Relevant Grade 3 and 4 Adverse Events in Phase III Studies of Combination Therapies^a

Adverse Event		Study 1			Study 2	
		Irinotecan HCl 5-FU/LV N=225 ^b	5-FU/LV N=219 ^b	Irinotecan HCl N=223 ^b	Irinotecan HCl 5-FU/LV N=145 ^c	5-FU/LV N=143 ^c
TOTAL Grade 3/4 Adverse Events		53.3	45.7	45.7	72.4	39.2
GASTROINTESTINAL						
Diarrhoea	late	22.7	13.2	31.0	14.4	6.3
	grade 3	15.1	5.9	18.4	10.3	4.2
	grade 4	7.6	7.3	12.6	4.1	2.1
	early	4.9	1.4	6.7	--	--
Nausea		15.6	8.2	16.1	2.1	3.5
Abdominal pain		14.6	11.5	13.0	2.1	0.7
Vomiting		9.7	4.1	12.1	3.5	2.8
Anorexia		5.8	3.7	7.2	2.1	0.7
Constipation		3.1	1.8	0.4	0.7	1.4
Mucositis		2.2	16.9	2.2	4.1	2.8
HAEMATOLOGICAL						
Neutropenia		53.8	66.7	31.0	46.2	13.4
	grade 3	29.8	23.7	19.3	36.4	12.7
	grade 4	24.0	42.5	12.1	9.8	0.7
Leucopenia		37.8	23.3	21.5	17.4	3.5
Anaemia		8.4	5.5	4.5	2.1	2.1
Neutropenic fever		7.1	14.6	5.8	3.4	0.7

Thrombocytopenia	2.6	2.7	1.7	0	0
Neutropenic infection	1.8	0	2.2	2.1	0
BODY AS A WHOLE					
Asthenia	19.5	11.9	13.9	9.0	4.2
Pain	3.1	3.6	2.2	9.7	8.4
Fever	1.7	3.6	0.4	0.7	0.7
Infection	0	1.4	0.4	7.6	3.5
METABOLIC AND NUTRITIONAL					
Increased bilirubin	7.1	8.2	7.2	3.5	10.6
DERMATOLOGICAL					
Exfoliative dermatitis	0	0.5	0	--	--
Rash	0	0.9	0.4	--	--
Hand and foot syndrome	--	--	--	0.7	0.7
Cutaneous signs	--	--	--	0.7	0
RESPIRATORY					
Dyspnoea	6.3	0.5	2.2	1.4	0
Cough	1.3	0	0.4	--	--
Pneumonia	2.7	1.0	1.3	--	--
NEUROLOGICAL					
Dizziness	1.3	0	1.8	--	--
Somnolence	1.8	1.8	1.3	--	--
Confusion	1.8	0	0	--	--
CARDIOVASCULAR					
Vasodilation	0.9	0	0	--	--
Hypotension	1.3	0.5	1.7	1.4	0
Thrombophlebitis	2.7	3.2	1.8	--	--
Pulmonary embolus	2.7	1.4	0.4	--	--
Myocardial infarction	1.3	0	0.4	--	--

^a Severity of adverse events based on National Cancer Institute's Common Toxicity Criteria (NCI CTC) (version 1.0)

^b Number of patients in the as-treated population for each group

^c Number of patients treated in the de Gramont regimen (B2/C2 treatment arms of Table 9)

The most clinically significant adverse events for patients receiving irinotecan hydrochloride-based therapy were diarrhoea, nausea, vomiting, neutropenia, and alopecia (complete hair loss = grade 2). The most clinically significant adverse events for patients receiving 5-FU/LV therapy were diarrhoea, neutropenia, neutropenic fever, and mucositis. In Study 1, grade 4 neutropenia, neutropenic fever (defined as \geq grade 2 fever and grade 4 neutropenia), and mucositis were observed less often with irinotecan hydrochloride/5-FU/LV than with administration of 5-FU/LV.

There is no evidence that the safety profile of irinotecan is influenced by cetuximab or *vice versa*. In combination with cetumaxib, additional reported undesirable effects were those expected with cetuximab (such as acneform rash).

Single agent therapy in recurrent or progressive metastatic colorectal cancer

Information on adverse reactions for irinotecan hydrochloride as single agent therapy is available from 304 patients with metastatic carcinoma of the colon or rectum treated in phase II trials with the once weekly dosage schedule, 316 patients treated with the once-every-3-week dosage schedule and over 1100 patients with a variety of tumour types treated in Japan. In general the types of toxicities observed were similar. 4.3% of patients treated with the weekly dosage schedule and 8% of patients treated with the once-every-3-week dosage schedule discontinued treatment with irinotecan hydrochloride because of medical events. Seventeen of the 304 patients treated with the weekly dosage schedule died within 30 days of the administration of irinotecan hydrochloride and in five cases (1.6%), the deaths were potentially drug-related. Eleven patients treated with irinotecan hydrochloride in the once-every-3-week dosage schedule died within 30 days of treatment and in three cases (1%), the deaths were potentially related to treatment with irinotecan hydrochloride. The main causes of the deaths potentially related to treatment were neutropenic infection, grade 4 diarrhoea and asthenia.

The frequency of the most common adverse events reported from the single agent second line studies is presented in Table 7 below. Additional information on adverse events follows the table, organised by body system category.

Table 7: Adverse events reported from the second line single agent therapy in 304 patients^a

Event	Weekly dosage schedule		3 weekly dosage schedule (NCI Grade 3 and 4 only)	
	% of Patients	% NCI Grade 3 & 4	Study 1 (%)	Study 2 (%)
GASTROINTESTINAL				
Diarrhoea (late)	87.8	30.6	21.7	22.0
Nausea	86.2	16.8	13.8	11.0
Vomiting	66.8	12.5	13.8	14.2
Abdominal cramping / pain	56.9	16.4	13.8	8.7
Anorexia	54.9	5.9	5.3	5.5
Diarrhoea (early)	50.7	7.9	12.2	1.6
Constipation	29.9	2.0	9.5	7.9
Flatulence	12.2	--	--	--
Stomatitis	11.8	0.7	--	--
Dyspepsia	10.5	--	--	--
HAEMATOLOGICAL				
Leucopenia ^b	63.2	28.0	22.2	14.2
Anaemia	60.5	6.9	7.4	6.3
Neutropenia ^b	53.9	26.3	22.2	14.2
Thrombocytopenia	--	--	1.1	3.9

BODY AS A WHOLE				
Asthenia	75.7	12.2	14.8	13.4
Fever	45.4	0.7	--	--
Pain	23.7	2.3	18.5 ^c	16.5 ^d
Headache	16.8	0.7	--	--
Back pain	14.5	1.6	--	--
Chills	13.8	0.3	--	--
Minor infection	14.5	0	--	--
Oedema	10.2	1.3	--	--
Abdominal enlargement	10.2	0.3	--	--
METABOLIC AND NUTRITIONAL				
Weight reduction	30.3	0.7		
Dehydration	14.8	4.3		
Increased alkaline phosphatase	13.2	3.9		
Increased SGOT	10.5	1.3		
DERMATOLOGICAL				
Alopecia	60.5	Not applicable ^e	Not applicable ^e	Not applicable ^e
Sweating	16.4	0	--	--
Rash	12.8	0.7	1.6	0.8
RESPIRATORY				
Dyspnoea	22.0	3.6		
Increased coughing	17.4	0.3		
Rhinitis	15.5	0		

^a Severity of adverse events based on NCI CTC (version 1.0)

^b Combined results for leucopenia/neutropenia are presented for the once-every-3-week dosage schedule

^c In this study, 22.2% of patients treated with best supportive care experienced NCI grade 3/4 pain

^d In this study, 13.2% of patients treated with infusional 5-FU experienced NCI grade 3/4 pain

^e Complete hair loss = NCI grade 2

Gastrointestinal

Nausea, vomiting and diarrhoea are common adverse events following treatment with irinotecan hydrochloride and can be severe. Among those patients treated at the 125 mg/m² single agent weekly dose, the median duration of any grade of late diarrhoea was 3 days, and for grade 3 or 4 late diarrhoea was 7 days. The frequency of grade 3 and 4 late diarrhoea was significantly greater in patients 65 years or older (39.8% versus 23.4%, p=0.0025).

Abdominal pain and cramping are associated with early-onset diarrhoea (diarrhoea which occurs within 24 hours of drug administration). In studies it has been found that atropine is useful in ameliorating these events. Colonic ulceration, sometimes with gastrointestinal bleeding, ileus and infection, has been observed in association with administration of irinotecan hydrochloride.

Haematological

Irinotecan hydrochloride commonly causes neutropenia, leucopenia (including lymphocytopenia) and anaemia. Serious thrombocytopenia is uncommon. In clinical studies with the single agent weekly dosage schedule, one death due to neutropenic sepsis without fever was judged to be potentially drug-related (0.3%, 1/304). Blood transfusions were given to 9.9% of patients. When evaluated in the trials of single agent weekly administration, the frequency of grade 3 or 4 neutropenia was significantly higher in patients who had received previous pelvic/abdominal irradiation than in those who had not received such irradiation (48.1% versus 24.1%, $p=0.0356$). In these same studies, patients with total baseline serum bilirubin levels of 17 $\mu\text{mol/L}$ or more also had a significantly greater likelihood of experiencing first-cycle grade 3 or 4 neutropenia than those with bilirubin levels that were less than 17 $\mu\text{mol/L}$ (50% versus 17.7%, $p<0.001$).

Cholinergic symptoms

Patients may have cholinergic symptoms of rhinitis, increased salivation, miosis, lacrimation, diaphoresis, flushing and intestinal hyperperistalsis that can cause abdominal cramping and early diarrhoea. If these symptoms occur, they manifest during or shortly after drug infusion. They are thought to be related to the anticholinesterase activity of the irinotecan parent compound and are more likely to occur at higher doses. The timing of the symptoms is most consistent with the occurrence of peak irinotecan hydrochloride serum levels during parenteral administration.

Metabolic and nutritional

The dehydration observed in 14.8% of patients in clinical studies was as a consequence of diarrhoea, nausea and vomiting.

Hepatic

For the once-every-3-week dosage schedule, hepatic events, such as ascites and jaundice of NCI grade 3/4 severity occurred in 8.5% of patients in one study and 8.7% of patients in another study. In the clinical studies evaluating the single agent weekly dosage schedule, NCI grade 3 or 4 liver enzyme abnormalities were observed in fewer than 10% of patients. These events typically occur in patients with known hepatic metastases. The dehydration observed in 14.8% of patients in these studies was as a consequence of diarrhoea, nausea and vomiting. Increases in serum creatinine or blood urea nitrogen, generally attributable to complications of infection or to dehydration related to nausea, vomiting or diarrhoea have been observed.

Renal

There have been cases of acute renal failure. Rare instances of renal dysfunction due to tumour lysis syndrome have also been reported.

Dermatological

Alopecia has been reported during treatment with irinotecan hydrochloride. Rashes have also been reported but did not result in discontinuation of treatment.

Respiratory

Severe pulmonary events are infrequent. Over half the patients with dyspnoea in the clinical studies evaluating the single agent weekly dosage schedule had lung metastases; the extent to which malignant pulmonary involvement or other pre-existing lung disease may have contributed to dyspnoea in these patients is unknown. For the once-every-3-week dosage schedule, respiratory events, such as dyspnoea and cough of NCI grade 3/4 severity occurred in 10.1% of patients in one study and 4.7% of patients in another study.

A potentially life-threatening pulmonary syndrome, consisting of dyspnoea, fever and a reticulonodular pattern on chest x-ray was observed in a small percentage of patients in early Japanese studies. The contribution of irinotecan hydrochloride to these preliminary events was difficult to assess because these patients also had lung tumours and some had pre-existing nonmalignant pulmonary disease. As a result of these observations, however, clinical studies in the USA enrolled few patients with compromised pulmonary function, significant ascites, or pleural effusions.

Neurological

Insomnia and dizziness were observed in 19.4% and 14.8% respectively of patients studied in clinical trials of the single agent weekly dosage schedule but were not usually considered to be directly related to the administration of irinotecan hydrochloride. Dizziness may sometimes represent symptomatic evidence of orthostatic hypotension in patients with dehydration.

Cardiovascular

Vasodilation (flushing) may occur during administration of irinotecan hydrochloride. Irinotecan hydrochloride has anti-cholinesterase activity. As such, there are possible cardiovascular effects due to its administration. These include sudden death, blackout and bradycardia. Patients should be monitored for cholinergic effects during administration of irinotecan hydrochloride, and atropine should be readily available for treatment of these effects. There were no cases of sudden death reported in the Phase II clinical studies of the single agent weekly dosage schedule involving 304 patients. In these studies, two patients (0.7%) suffered syncope and one patient (0.3%) suffered bradycardia.

Thromboembolic events including angina pectoris, arterial thrombosis, cerebral infarct, cerebrovascular accident, deep thrombophlebitis, embolus lower extremity, heart arrest, myocardial infarct, myocardial ischemia, peripheral vascular disorder, pulmonary embolus, sudden death, thrombophlebitis, thrombosis and vascular disorder have been observed rarely in patients receiving irinotecan hydrochloride. The specific cause of these events has not been determined.

Other

Other NCI grade 3 or 4 drug-related adverse events observed in 1-10% of patients in clinical trials included mucositis, bilirubinaemia and hypovolaemia. In fewer than 1% of patients, NCI grade 3 or 4 rectal disorder, gastrointestinal monilia, hypokalaemia, hypomagnesaemia, increased GGTP, malaise, sepsis and abnormal gait were observed.

Post-marketing Experience

Cardiac disorders: Myocardial ischemic events have been observed following irinotecan therapy predominantly in patients with underlying cardiac disease, other known risk factors for cardiac disease or previous cytotoxic chemotherapy.

Gastrointestinal disorders: Infrequent cases of intestinal obstruction, ileus, megacolon or gastrointestinal hemorrhagic and rare cases of colitis, including typhilitis (ileocecal syndrome), ischaemic and ulcerative colitis have been reported. In some cases, colitis was complicated by ulceration, bleeding, ileus or infection. Cases of ileus without preceding colitis have also been reported. Rare cases of intestinal perforation have been reported. Rare cases of symptomatic pancreatitis or asymptomatic elevated pancreatic enzymes have been observed.

Hypovolaemia: There have been rare cases of renal impairment and acute renal failure, generally in patients who became infected and/or volume depleted from severe gastrointestinal toxicities. Infrequent cases of renal insufficiency, hypotension or circulatory failure have been observed in patients who experienced episodes of dehydration associated with diarrhoea and/or vomiting, or sepsis.

Infections and infestations: Bacterial, fungal and viral infections have been reported.

Immune system disorders: Hypersensitivity reactions including severe anaphylactic or anaphylactoid reactions have also been reported.

Investigations: Rare cases of hyponatraemia mostly related with diarrhoea and vomiting have been reported. Transient and mild to moderate increases in serum levels of transaminases (i.e. AST and ALT) in the absence of progressive liver metastasis, transient increase of amylase and occasionally transient increase of lipase have been very rarely reported.

Musculoskeletal and connective tissue disorders: Early effects such as muscular contraction or cramps and paresthesia have been reported.

Nervous system disorders: Speech disorders, generally transient in nature, have been reported in patients treated with irinotecan; in some cases, the event was attributed to the cholinergic syndrome observed during or shortly after infusion of irinotecan.

Respiratory disorders: Interstitial pulmonary disease presenting as pulmonary infiltrates is uncommon during irinotecan therapy. Early effects such as dyspnea have been reported (refer to section 4.4). Hiccups have been reported.

Renal and cardiovascular disorders: Infrequent cases of renal insufficiency including acute renal failure, hypotension or circulatory failure have been observed in patients who experienced episodes of dehydration associated with diarrhoea and/or vomiting, or sepsis (see section 4.4).

Reporting of Suspected Adverse Reactions

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare professionals are asked to report any suspected adverse reactions <https://nzphvc.otago.ac.nz/reporting/>.

4.9 Overdose

Symptoms

In humans, at single doses up to 750 mg/m², adverse events were similar to those reported with the recommended dosage regimens. There have been reports of overdosage at doses up to approximately twice the recommended therapeutic dose, which may be fatal. The most significant adverse effects reported were severe neutropenia and severe diarrhoea.

Treatment

There is no known antidote for overdosage of irinotecan hydrochloride. Support respiratory and cardiovascular function. Maximum supportive care should be instituted to prevent dehydration due to diarrhoea and to treat any infectious complications.

For advice on the management of overdose please contact the National Poisons Centre on 0800 POISON (0800 764766).

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Mechanism of action

Irinotecan hydrochloride is a derivative of camptothecin. Camptothecins interact specifically with the enzyme topoisomerase I, which relieves torsional strain in DNA by inducing reversible single-strand breaks. Irinotecan hydrochloride and its active metabolite SN-38 bind to the topoisomerase I - DNA complex and prevent religation of these single-strand breaks. Current research suggests that the cytotoxicity of irinotecan hydrochloride is due to double-strand DNA damage produced during DNA synthesis when replication enzymes interact with the ternary complex formed by topoisomerase I, DNA, and either irinotecan hydrochloride or SN-38. Mammalian cells cannot efficiently repair these double-strand breaks.

Irinotecan hydrochloride serves as a water-soluble precursor of the lipophilic metabolite SN-38 which is approximately 1000 times as potent as irinotecan hydrochloride as an inhibitor of topoisomerase I purified from human and rodent tumour cell lines. However, the precise contribution of SN-38 to the activity of irinotecan hydrochloride is unknown. Both irinotecan hydrochloride and SN-38 exist in an active lactone form and an inactive hydroxy acid anion form. An acidic pH promotes the formation of the lactone whereas a basic pH favours the hydroxy acid anion form.

Administration of irinotecan hydrochloride has resulted in antitumour activity in mice bearing cancers of rodent origin and human carcinoma xenografts of various histological types.

Irinotecan hydrochloride is a non-competitive inhibitor of acetylcholinesterase and a cholinergic syndrome is associated with its administration (see section 4.8).

Clinical trials

Irinotecan hydrochloride has been studied in clinical trials in combination with 5-FU and LV as a first line agent in metastatic colorectal cancer and as a single agent used after failure of

initial therapy. Weekly and once every 3 weeks dosage schedules were studied using irinotecan hydrochloride as the single agent. Weekly and once every 2 week schedules were studied with irinotecan hydrochloride used in combination treatment. Patients with a WHO performance status of 3 or 4 have not been studied in clinical trials (refer to Table 8).

Table 8: WHO scale for performance status

WHO scale	Performance status
0	Fully active; able to carry on all pre-disease performance without restriction
1	Restricted in physically strenuous activity but ambulatory and able to carry out work of a light or sedentary nature
2	Ambulatory and capable of self-care but unable to carry out any work activities; up and about more than 50% of waking hours
3	Capable of only limited self-care; confined to bed or chair more than 50% of waking hours
4	Completely disabled; cannot carry out any self-care; totally confined to bed or chair

Combination therapy for first-line treatment of metastatic colorectal cancer

Two randomised, open-label, controlled, multinational, phase III clinical trials support the use of irinotecan hydrochloride as first-line treatment of patients with metastatic carcinoma of the colon or rectum. The dosing regimens of these studies are given in Table 9.

Table 9: Dosage regimen of the studies evaluating the first line treatment of metastatic colorectal cancer

Arm	Agent	Study 1 Dosing Regimen	Study 2 Dosing Regimen
A	Irinotecan HCl	125 mg/m ² irinotecan HCl IV infusion over 90 mins. Treatment was administered once weekly for four weeks with treatment resuming on Day 43.	N/A
B1	Irinotecan HCl LV 5-FU	125 mg/m ² irinotecan HCl IV infusion over 90 mins followed immediately by 20 mg/m ² LV administered as an IV bolus injection and then 500 mg/m ² 5-FU as an IV bolus injection. Treatment was administered once weekly for four weeks with treatment resuming on Day 43 (Saltz regimen) ^a .	80 mg/m ² IV infusion over 90 mins of irinotecan HCl plus a 500 mg/m ² LV IV infusion over two hours followed immediately by an 2300 mg/m ² 5-FU IV infusion over 24 hours. Treatment was administered once weekly for six weeks with treatment resuming on Day 50 (AIO regimen) ^a

B2	Irinotecan HCl LV 5-FU	N/A	180 mg/m ² IV infusion over 90 mins of irinotecan HCl on Day 1, plus one hour later a 200 mg/m ² LV IV infusion over two hours followed immediately by a 400 mg/m ² 5-FU IV bolus injection and a 600 mg/m ² 5-FU IV infusion over 22 hours on Days 1 and 2. Treatment was administered every two weeks (de Gramont regimen) ^a
C1	LV 5-FU	20 mg/m ² LV administered as an IV bolus injection followed immediately by 425 mg/m ² 5-FU as an IV bolus injection. Treatment was given for 5 consecutive days with the treatment repeating on Day 29 (Mayo Clinic regimen) ^a .	500 mg/m ² LV IV infusion over two hours followed immediately by a 2600 mg/m ² 5-FU IV infusion over 24 hours. Administration was weekly for six weeks with treatment resuming on Day 50 (AIO regimen) ^a
C2	LV 5-FU	N/A	200 mg/m ² LV IV infusion over two hours followed immediately by a 400 mg/m ² 5-FU IV bolus injection and a 600 mg/m ² 5-FU IV infusion over 22 hours on Days 1 and 2. Treatment was administered every two weeks (de Gramont regimen) ^a

^aBased on the Saltz, Mayo Clinic, de Gramont and Association of Medical Oncology of the German Cancer Society (AIO) dosing regimens

In both studies, concomitant medications such as anti-emetics, atropine and loperamide were given to patients for prophylaxis and/or management of symptoms from treatment. In study 2, if late diarrhoea persisted for greater than 24 hours despite loperamide, a 7 day course of fluoroquinolone antibiotic prophylaxis was given. Treatment with oral fluoroquinolone was initiated in patients whose diarrhoea persisted for greater than 24 hours despite loperamide or if they developed a fever in addition to diarrhoea. Treatment with oral fluoroquinolone was also initiated in patients who developed an absolute neutrophil count (ANC) < 0.5 x 10⁹/L, even in the absence of fever or diarrhoea. Patients also received treatment with intravenous antibiotics if they had persistent diarrhoea or fever or if ileus developed.

In both studies the combination of irinotecan hydrochloride/5-FU/LV therapy resulted in significant improvements in objective tumour response rate, time to tumour progression (TTP) and survival when compared with 5-FU/LV alone. These differences in survival were observed despite the use of post-study second-line therapy, including irinotecan-containing regimens in patients in the control arm. Patient characteristics and major efficacy results are shown in Table 10.

Table 10: Combination therapy in first line treatment of metastatic colorectal cancer: Study Results

	Study 1	Study 2
--	---------	---------

Demographics and Treatment Administration	Irinotecan HCl 5-FU/LV	5-FU/LV	Irinotecan HCl	Irinotecan HCl 5-FU/LV	5-FU/LV
Number of Patients	231	226	226	198	187
Female/Male (%)	34/65	45/54	35/64	33/67	47/53
Median Age in Years (range)	62 (25-85)	61 (19-85)	61 (30-87)	62 (27-75)	59 (24-75)
Performance Status (%) ^a					
0	39	41	46	51	51
1	46	45	46	42	41
2	15	13	8	7	8
Median Primary Tumour (%)					
Colon	81	85	84	55	65
Rectum	17	14	15	45	35
Median Time from Diagnosis to Randomisation (months, range)	1.9 (0-161)	1.7 (0-203)	1.8 (0.1-185)	4.5 (0-88)	2.7 (0-104)
Prior Adjuvant 5-FU Therapy(%)					
No	89	92	90	74	76
Yes	11	8	10	26	24
Median Duration of Study Treatment (months)	5.5	4.1	3.9	5.6	4.5
Median Relative Dose Intensity (%)					
Irinotecan	72	--	75	87	--
5-FU	71	86	--	86	93
Efficacy Results					
Confirmed Objective Tumour Response Rate ^b (%) [95% CI]	39 [33-46]	21 [16-27]	18 [13-24]	35 [28-42]	22 [16-29]
Median Time to Tumour Progression (months) [95% CI]	7.0 [5.4-8.0]	4.3 [3.7-4.6]	4.2 [3.9-5.0]	6.7 [5.7-8.0]	4.4 [3.2-5.5]
Median Survival (months) [95% CI]	14.8 [12.3-17.1]	12.6 [11.1-14.6]	12.0 [11.3-13.5]	17.4 [15.2-20.2]	14.1 [12.6-17.4]

^a Refer to Table 8

^b Confirmed \geq 4 to 6 weeks after first evidence of objective response

Improvement was noted when response rates and time to tumour progression were examined across all demographic and disease-related subgroups (as categorised by age, gender, ethnic origin, performance status, extent of organ involvement with cancer, time from diagnosis of cancer, prior adjuvant therapy, and baseline laboratory abnormalities), with irinotecan hydrochloride-based combination therapy relative to 5-FU/LV.

The European Organisation of Research and Treatment of Cancer Quality of Life Questionnaire (EORTC QLQ-C30) was used in both studies. While there was no statistical evidence that there were significant differences between irinotecan hydrochloride/5-FU/LV combination and 5-FU/LV alone with regard to quality of life (QOL) improvement, descriptive evidence suggested a general trend favouring QOL improvement or less-worsening in favour of the irinotecan hydrochloride combination regimen.

Single Agent Treatment in Recurrent or Progressive Metastatic Colorectal Cancer After 5-FU Based Treatment

Weekly Dosage Schedule

Three multicentre, open-label, phase II studies, all utilising repeated cycles of once weekly treatment with irinotecan hydrochloride for 4 consecutive weeks, followed by a two week rest period were conducted in a total of 304 patients in the United States. These studies were designed to evaluate tumour response rate and toxicity with irinotecan hydrochloride in patients with metastatic colorectal cancer that recurred or progressed following a prior 5-FU based chemotherapeutic regimen. Starting doses of irinotecan hydrochloride in these trials were 100, 125 or 150 mg/m², with 150 mg/m² proving to be poorly tolerated due to unacceptably high rates of grade 4 late diarrhoea and febrile neutropenia. The results of the studies are shown in Table 11.

Table 11: Phase II clinical studies with the once weekly dosage schedule

	Study			
	A	B	C ^a	C ^a
Number of patients	48	90	64	102
Dose (mg/m ² /wk x 4)	125 ^b	125	125	100
Prior 5-FU therapy (%)				
For metastatic disease	81.3	65.5	73.4	67.7
7.6 months after adjuvant	14.6	6.7	26.6	27.5
> 6 months after adjuvant	2.1	15.6	0.0	2.0
Classification unknown	2.1	12.2	0.0	2.9
Duration of treatment (median, months)	5.4	3.5	3.9	3.3
Median relative dose intensity (%) ^c	74	67	73	81
Objective response rate (%) ^d	20.8	13.3	14.1	8.8
[95% CI]	[9.3 - 32.3]	[6.3 - 20.4]	[5.5 - 22.6]	[3.3 - 14.3]
Time to response (median, months)	2.6	1.5	2.8	2.8
Response duration (median, months)	6.4	5.9	5.6	6.4
Survival (median, months)	10.4	8.1	10.7	9.3

^a The initial dose in Study C was 125 mg/m² but was reduced to 100 mg/m² because the toxicity at the starting dose was perceived to be greater than seen in previous studies. Results are analysed separately for the two starting doses

^b Nine patients received 150 mg/m² as a starting dose; 2 (22.2%) responded to irinotecan hydrochloride

^c Relative dose intensity for irinotecan hydrochloride based on planned dose intensity of 100, 83.3 and 66.7 mg/m²/wk corresponding with 150, 125 and 100 mg/m² starting doses respectively

^d There were 2 complete responses and 38 partial responses

Of the 304 patients treated in the phase II studies, response rates to irinotecan hydrochloride were similar in males and females and among patients younger than 65 years. Rates were also similar in patients with cancer of the colon or cancer of the rectum, and in patients with single and multiple metastatic sites. Response rate was 18.5% in patients with a WHO performance status of 0, and 8.2% in patients with a performance status of 1 or 2.

The response rates with irinotecan hydrochloride were unaffected by whether or not patients had responded to prior 5-FU based treatment given for metastatic disease. Patients who had received previous irradiation to the pelvis also responded to irinotecan hydrochloride at approximately the same rate as those who had not previously received irradiation.

Overall, across the pivotal studies, stable disease was documented in 148 (48.7%) of the 304 patients in the intent to treat population and in 145 (55.6%) of the 261 patients in the evaluable population. Consistent with the results in Study C, a somewhat greater percentage of patients who were treated with the 125 mg/m² starting dose (53.4%; 103/193) than with the 100 mg/m² starting dose (39.2%; 40/102) had stable disease during therapy.

Once Every 3 Week Dosage Schedule

Two phase III, multicentre, randomised studies were conducted with a three weekly dosage regimen in patients with metastatic colorectal cancer whose disease had recurred or progressed following 5-FU therapy (n = 535). Second-line irinotecan hydrochloride was compared with best supportive care in one study and with infusional 5-FU-based therapy in the second study. The primary endpoint in both studies was survival. Parameters of clinical benefit and quality of life were also assessed. The starting dose was 350 mg/m² infused intravenously over 90 minutes to a maximum total dose of 700 mg. For patients 70 years or older and for patients with a WHO performance status of 2 the starting dose was reduced to 300 mg/m². Anti-emetics, atropine and loperamide were provided as supportive care and late diarrhoea persisting for greater than 24 hours despite loperamide was treated with a 7-day course of a fluoroquinolone antibiotic.

A significant survival advantage for irinotecan hydrochloride over best supportive care or infusional 5-FU-based therapy was demonstrated. When adjusted for baseline patient characteristics (e.g. performance status), survival among patients treated with irinotecan hydrochloride remained significantly longer than in the control populations (p = 0.001 for Study 1 and p = 0.017 for Study 2). Clinical benefit in Study 1, as measured by pain-free survival and survival without weight loss were significantly longer for patients treated with irinotecan hydrochloride than for patients in the best supportive care group (p, 0.01 and p, 0.05 respectively). The results are summarised in Table 12.

Table 12: Phase III clinical studies with the once every 3 week dosage schedule

	Study 1		Study 2	
	Irinotecan hydrochloride	Best supportive care	Irinotecan hydrochloride	5-FU ^a
Number of patients	189	90	127	129
Prior 5-FU therapy (%)				
For metastatic disease	70	63	58	68
≤ 3/6 months after adjuvant ^b	27	36	38	23
> 3/6 months after adjuvant ^b	3	0	5	9
Duration of treatment (mean, months) [95% CI]	4.6 [4.2 – 5.0]	--	4.4 [3.8 – 5.0]	3.7 [3.3 – 4.1]
Median relative dose intensity (%) ^c	94	--	95	81-99
Survival (median, months) [95% CI]	9.2 [8.4 – 10.7]	6.5 [5.0 – 7.6]	10.8 [9.5 – 12.8]	8.5 [7.7 – 10.5]
1-year survival (%) [95% CI]	36.2 [29.3 – 43.1]	13.8 [6.7 – 20.9]	44.8 [36.2 – 53.4]	32.4 [24.3 – 40.5]
Progression-free survival (median, months) [95% CI]	--	--	4.2 [3.8 – 4.8]	2.9 [2.6 – 3.7]
Symptom-free survival (median, months) [95% CI]	5.9 [3.8 - 7.6]	4.1 [2.2 - 6.9]	8.1 [6.1 - 10.7]	7.0 [4.4 - 8.7]
Pain-free survival (median, months) [95% CI]	6.9 [5.8 – 8.4]	2.0 [1.8 – 5.1]	10.3 [7.8 -**]	8.5 [6.2 – 10.2]
Median survival without performance status deterioration (%) [95% CI]	5.7 [4.3 – 6.6]	3.3 [1.9 – 3.7]	6.4 [5.2 – 7.6]	5.1 [4.2 – 6.2]
Time to weight loss ≥ 5% (median, months) [95% CI]	6.4 [5.5 – 7.6]	4.2 [3.4 – 5.1]	8.9 [6.7 – 12.3]	7.4 [4.7 – 11.6]

^a One of the following 5-FU regimens was used:

(i) Leucovorin 200 mg/m² iv over 2 hours; followed by 5-FU 400 mg/m² iv bolus; followed by 5-FU 600 mg/m² continuous iv infusion over 22 hours on Days 1 and 2 every 2 weeks.

(ii) 5-FU 250-300 mg/m²/day protracted continuous iv infusion until toxicity.

(iii) 5-FU 2.6-3 g/m²/day iv over 24 hours every week for 6 weeks with or without leucovorin 20-500 mg/m²/day every week iv for 6 weeks with a 2 week rest between cycles

^b Study 1 ≤ 6 months; Study 2 ≤ 3 months

^c Relative dose intensity for irinotecan hydrochloride based on planned dose intensity of 116.7 mg/m²/week. Dose intensity in patients receiving 5-FU in Study 2 varied depending upon type of regimen

** Cannot be estimated due to small sample size

In the two phase III studies, quality of life was assessed using the European Organisation on Research and Treatment of Cancer (EORTC) QLQ-C30 questionnaire. In Study 1, the global quality of life scores were significantly higher for patients treated with irinotecan hydrochloride than for those who received best supportive care (p=0.0013). In Study 2, the global quality of life scores were similar for patients who received either irinotecan hydrochloride or infusional 5-FU.

Other studies

A Japanese open-label, uncontrolled, late phase II study in patients with non-small-cell lung cancer enrolled a total of 153 patients. In this study, pneumonitis occurred in 6.2% (9/146) of the patients. One patient died of interstitial pneumonitis. Irinotecan hydrochloride was given at a dose of 100 mg/m² intravenously once weekly. Dosage adjustments were made according to toxicity and the duration of treatment was until disease progression or unacceptable toxicity occurred (with each patient to receive at least three doses).

5.2 Pharmacokinetic properties

Absorption

Over the recommended dose range of 50 to 350 mg/m², the AUC of irinotecan hydrochloride increases linearly with dose; the AUC of SN-38 increases less than proportionally with dose. Maximum concentrations of the active metabolite SN-38 are generally seen within 1 hour following the end of a 90 minute infusion of irinotecan hydrochloride.

Pharmacokinetic parameters for irinotecan hydrochloride and SN-38 following a 90-minute infusion of irinotecan hydrochloride at dose levels of 125 and 340 mg/m² determined in two clinical studies in patients with solid tumours are summarised in Table 13.

Table 13: Summary of mean (± standard deviation) irinotecan hydrochloride and SN-38 pharmacokinetic parameters in patients with solid tumors

Dose (mg/m ²)	Irinotecan hydrochloride					SN-38		
	C _{max} (ng/mL)	AUC ₀₋₂₄ (ng.hr/mL)	t _{1/2} (hr)	V _{area} (L/m ²)	CL (L/hr/m ²)	C _{max} (ng/mL)	AUC ₀₋₂₄ (ng.hr/mL)	T _{1/2} (hr)
125 (n=64)	1,660 ± 797	10,200 ± 3,270	5.8 ^a ± 0.7	110 ± 48.5	13.3 ± 6.01	26.3 ± 11.9	229 ± 108	10.4 ^a ± 3.1
340 (n=6)	3,392 ± 874	20,604 ± 6,027	11.7 ^b ± 1.0	234 ± 69.6	13.9 ± 4.00	56.0 ± 28.2	474 ± 245	21.0 ^b ± 4.3

C_{max}: Maximum plasma concentration

AUC₀₋₂₄: Area under the plasma concentration-time curve from time 0 to 24 hours after the end of the 90-minute infusion

t_{1/2}: Terminal elimination half-life

V_{area}: Volume of distribution of terminal elimination phase

CL: Total systemic clearance

^a Plasma specimens collected for 24 hours following the end of the 90-minute infusion

^b Plasma specimens collected for 48 hours following the end of the 90-minute infusion. Because of the longer collection period, these values provide a more accurate reflection of the terminal elimination half-lives of irinotecan hydrochloride and SN-38

In vitro studies indicate that irinotecan hydrochloride exhibits moderate plasma protein binding (30% to 68% bound). SN-38 is highly bound to human plasma proteins (approximately 95% bound). The plasma protein to which irinotecan hydrochloride and SN-38 predominantly bind is albumin.

Distribution

After intravenous infusion of irinotecan hydrochloride in humans with various cancers, irinotecan hydrochloride plasma concentrations decline in a multi-exponential manner, with a mean terminal elimination half-life of about 6 to 12 hours. The mean terminal elimination half-life of the active metabolite SN-38 is about 10 to 20 hours. In a study where irinotecan hydrochloride was administered at doses of 100-750 mg/m² by 30 minute intravenous infusion every three weeks, the plasma terminal elimination half-life was 14.2 ± 7.7 hours for irinotecan hydrochloride and 13.8 ± 1.4 hours for SN-38.

Biotransformation

The complete disposition of irinotecan hydrochloride has not been fully elucidated in humans. Irinotecan hydrochloride is subject to extensive metabolic conversion by various enzyme systems, including esterases to form the active metabolite SN-38, and uridine diphosphate-glucuronosyl transferase 1A1 (UGT1A1) mediating glucuronidation of SN-38 to form the inactive metabolite SN-38 glucuronide (SN-38G). The metabolic conversion of irinotecan hydrochloride occurs primarily in the liver. Irinotecan can also undergo CYP3A4-mediated oxidative metabolism to several pharmacologically inactive oxidation products, one of which can be hydrolysed by carboxylesterase to release SN-38. Irinotecan is oxidised by cytochrome P450 isozyme 3A4 (CYP3A4) to yield two relatively inactive metabolites, APC (7-ethyl-10-[4-N-(5-aminopentanoic acid)-1-piperidino]-carbonyloxycamptothecin) and the minor metabolite, NPC (7-ethyl-10-(4-amino-1-piperidino)carbonyloxycamptothecin).

Elimination

The disposition of irinotecan hydrochloride has not been fully elucidated in humans. In studies of patients with various cancers the urinary excretion of irinotecan hydrochloride was 11% to 20% of the administered dose; SN-38, <1%; and SN-38 glucuronide, 3%. The cumulative biliary and urinary excretion of irinotecan hydrochloride and its metabolites (SN-38 and SN-38 glucuronide) over a period of 48 hours following administration of irinotecan hydrochloride in two patients ranged from approximately 25% (100 mg/m²) to 50% (300 mg/m²).

Special Populations

Geriatric (≥ 65 years): In studies where irinotecan hydrochloride was administered weekly, the terminal half-life of irinotecan hydrochloride was 6.0 hours in patients who were 65 years or older, and 5.5 hours in patients younger than 65 years. Dose-normalised AUC₀₋₂₄ for SN-38 in patients who were at least 65 years of age was 11% higher than in patients younger than 65 years. There are no kinetic data on the use of the once-every-three-week dosage schedule in elderly patients. A lower starting dose is recommended in patients 65 years and older based on clinical toxicity experienced with this dosage regimen (see section 4.2).

Hepatic impairment: The influence of severe hepatic insufficiency on the pharmacokinetic characteristics of irinotecan hydrochloride and its metabolites has not been formally studied. Among patients with metastatic colorectal cancer and known hepatic tumour involvement (a

majority of patients), irinotecan hydrochloride and SN-38 AUC values were somewhat higher than values for patients without liver metastases (see section 4.4). Irinotecan clearance is diminished in patients with hepatic dysfunction while relative exposure to the active metabolite SN-38 is increased. The magnitude of these effects is proportional to the degree of liver impairment as measured by elevations in serum total bilirubin and transaminase concentrations (see section 4.2).

Renal impairment: The influence of renal insufficiency on the pharmacokinetics of irinotecan hydrochloride has not been evaluated.

Pharmacokinetics in Combination Therapy

In a phase I clinical study involving irinotecan hydrochloride, fluorouracil (5-FU), and leucovorin (LV) in 26 patients with solid tumours the disposition of irinotecan hydrochloride was not substantially altered when the drugs were co-administered. However, C_{max} and AUC₀₋₂₄ of SN-38, the active metabolite, were reduced (by 14% and 8%, respectively) when irinotecan hydrochloride was followed by 5-FU and LV administration compared with when irinotecan hydrochloride was given alone. Formal *in vivo* or *in vitro* drug interaction studies to evaluate the influence of irinotecan hydrochloride on the disposition of 5-FU and LV have not been conducted.

5.3 Preclinical safety data

Genotoxicity

Irinotecan hydrochloride was clastogenic both *in vitro* (Chinese hamster ovary cells) and *in vivo* (micronucleus test in mice). Neither irinotecan hydrochloride or SN-38 was mutagenic in the *in vitro* Ames assay.

Carcinogenicity

Long-term carcinogenicity studies with irinotecan hydrochloride were not conducted. Rats were, however, administered intravenous doses of 2 mg/kg or 25 mg/kg irinotecan hydrochloride once per week for 13 weeks (AUC about 1.3 times the values of patients administered 125 mg/m²) and were then allowed to recover for 91 weeks. Under these conditions, there was a significant linear trend with dose for the incidence of combined uterine horn endometrial stromal polyps and endometrial stromal sarcomas.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Hydrochloric acid (pH adjustment),
Lactic acid,
Sodium hydroxide (pH adjustment),
Sorbitol,
Water for injection.

6.2 Incompatibilities

Other drugs should not be added to the infusion solution.

6.3 Shelf life

3 years

6.4 Special precautions for storage

Store at or below 30°C. Protect from light. Do not freeze.

6.5 Nature and contents of container

Irinotecan Injection Concentrate is supplied in single use, amber glass vials containing 20 mg/mL of irinotecan hydrochloride trihydrate.

Product	Pack Size
----------------	------------------

40 mg/2 mL	1 vial
------------	--------

100 mg/5 mL	1 vial
-------------	--------

500 mg/25 mL	1 vial
--------------	--------

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

As with other potentially toxic anticancer agents, care should be exercised in the handling and preparation of infusion solutions prepared from Irinotecan Injection Concentrate. The use of gloves is recommended. If a solution of Irinotecan Injection Concentrate contacts the skin, wash the skin immediately and thoroughly with soap and water. If Irinotecan Injection Concentrate contacts the mucous membranes, flush thoroughly with water.

Irinotecan Injection Concentrate contains no antimicrobial agent. It is for single use in one patient only. Discard any residue.

Preparation of infusion solution

The vial should be inspected for damage and visible signs of leaks. If damaged, incinerate the unopened package.

Inspect vial contents for particulate matter and repeat inspection when drug product is withdrawn from vial into syringe. If particulate matter is seen, do not use the contents.

Irinotecan Injection Concentrate must be diluted prior to infusion. Irinotecan Injection Concentrate should be diluted in 5% Glucose Injection (preferred) or 0.9% Sodium Chloride Injection to a final concentration range of 0.12 to 2.8 mg/mL. Other drugs should not be added to the infusion solution. Parenteral drug products should be inspected visually for particulate matter and discolouration prior to administration whenever solution and container permit.

Solutions diluted in 0.9% Sodium Chloride Injection in infusion bags are physically and chemically stable for up to 24 hours at 25°C when exposed to light, and up to 7 days when stored refrigerated (2 to 8°C) in the dark. Refrigeration of admixtures using 0.9% sodium chloride injection is not recommended due to a low and sporadic incidence of visible particles. Solutions diluted in 5% Glucose Injection in infusion bags are physically and chemically stable for 48 hours at 25°C when exposed to light, and to 7 days when stored refrigerated (2 to 8°C) in the dark. To reduce microbiological hazard, use as soon as practicable after preparation. If storage is necessary, hold at 2° to 8°C for not more than 24 hours.

Do not freeze admixtures of Irinotecan Injection Concentrate as this may result in precipitation of the drug.

Pharmaceutical precautions

The following protective recommendations are given due to the toxic nature of this substance:

- Personnel should be trained in good technique for handling.
- Pregnant staff should be excluded from working with this drug.
- Personnel handling Irinotecan Injection Concentrate should wear protective clothing: goggles, gowns and disposable gloves and masks.
- A designated area should be defined for reconstitution (preferably under a laminar flow containment system). The work surface should be protected by disposable plastic backed absorbent paper.
- All items used for administration of cleaning, including gloves, should be placed in high-risk, waste-disposal bags for high temperature incineration.
- Spillage or leakage should be treated with dilute sodium hypochlorite (1% available chlorine) solution, preferably by soaking, and then water.
- All cleaning materials should be disposed of as indicated previously.
- Accidental contact with the eyes or skin should be treated immediately. Copious lavage with water is appropriate treatment for contact with the eyes, whereas water or soap and water, or sodium bicarbonate solution may be used on the skin; medical attention should be sought.

7. MEDICINE SCHEDULE

Prescription Medicine

8. SPONSOR

Pfizer New Zealand Limited
P O Box 3998
Auckland, New Zealand, 1140
Toll Free Number: 0800 736 363

9. DATE OF FIRST APPROVAL

14 July 2005

10. DATE OF REVISION OF THE TEXT

16 January 2020

™ = Trademark

SUMMARY TABLE OF CHANGES

Section changed	Summary of new information
All	Minor editorial changes.
4.6	Use in pregnancy – Pregnancy Category D Additional information regarding pregnancy and genotoxicity. Use in lactation Additional information regarding lactation.
4.8	Additional information regarding the safety profile of combination therapy with irinotecan and cetumaxib.