

NEW ZEALAND DATA SHEET

1. PRODUCT NAME

JESPECT®. Inactivated Japanese Encephalitis Vaccine (adsorbed), suspension for injection.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

JESPECT® Japanese Encephalitis (JE) Virus, purified inactivated vaccine is a sterile, ready to use suspension for intramuscular (IM) injection. The vaccine is prepared by propagating Japanese encephalitis virus strain (SA₁₄₋₁₄₋₂) in Vero cells. No preservatives or antibiotics are added to the formulation.

JESPECT® is a clear liquid with white precipitate and when shaken before use a white/cloudy suspension forms. JESPECT® is supplied in pre-filled syringes without needles. Each 0.5 mL dose of vaccine contains 6 antigen units (AgU) of purified, inactivated JE virus. Each dose of vaccine also contains the following excipients: 0.1% aluminium hydroxide hydrate corresponding to 0.25 mg aluminium, and Phosphate Buffered Saline. Refer section 6.1 – List of excipients.

3. PHARMACEUTICAL FORM

Suspension for injection.

Clear liquid with a white precipitate.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

JESPECT® is indicated for active immunisation against Japanese Encephalitis (JE) virus for persons 18 years of age and older.

JESPECT® should be considered for use in persons who plan to reside in or travel to areas where JE is endemic (common) or epidemic (seasonal), especially during the transmission season.

JESPECT® is indicated for persons who work with JE virus in laboratories and in industry.

4.2 Dose and method of administration

Primary vaccination

The primary vaccination series consists of a total of two doses of 0.5 mL each according to the following schedule:

- First dose at Day 0 (elected date).
- Second dose: 28 days after first dose.

It is recommended that persons who receive the first dose of JESPECT® complete the 2-dose primary vaccination course.

If the primary immunisation schedule of 2 doses is not completed, sufficient protection against Japanese encephalitis virus might not be achieved. Clinical trial data indicates that administration of the second dose up to 11 months after the first dose results in high seroconversion rates (see section **5.1 Pharmacodynamic properties**).

Booster dose

Refer to section **5.1 Pharmacodynamic properties** for clinical trial data for potential booster doses.

Method of administration

The vaccine should be inspected visually for the particulate matter and discolouration prior to administration. Discard the product if particulates are present, if it appears discoloured or if the syringe appears to be physically damaged. Any unusual product or waste material should be disposed of in accordance with local requirements.

JESPECT® should be well shaken before administration to obtain a homogenous suspension. Once shaken, the vaccine should appear as a white cloudy suspension.

JESPECT® should be administered by IM injection into the deltoid muscle. It should never be injected IV.

In exceptional circumstances, for those patients with thrombocytopenia or bleeding disorders where bleeding may occur following IM administration, JESPECT® may be administered subcutaneously. Subcutaneous administration could lead to a suboptimal response to the vaccine.

JESPECT® pre-filled syringe is for single use only in one individual only. Inject the entire contents of the syringe.

4.3 Contraindications

JESPECT® should not be administered to individuals who have previously experienced a serious reaction (e.g. anaphylaxis) to this vaccine or who are known to be hypersensitive to any of the vaccine components or to the residues protamine sulphate, formaldehyde, bovine serum albumin, host cell DNA, sodium metabisulphite, host cell protein. Individuals who show hypersensitivity reactions after receiving the first dose of the vaccine should not be given the second dose.

4.4 Special warnings and precautions for use

As with other injectable vaccines, appropriate medical treatment and supervision should always be available in the event of anaphylactic reaction. Adrenaline should always be readily available whenever the injection is given.

JESPECT® will not protect against encephalitis caused by other organisms. As with any other vaccine, vaccination with JESPECT® may not result in protection in all cases. The primary vaccination series (2 doses) should be completed at least one week prior to potential exposure to Japanese Encephalitis virus. Seroconversion rates after one dose are limited (see section **5.1 Pharmacodynamic properties**, Table 3).

JESPECT® should be administered by IM injection into the deltoid muscle. It should never be injected IV. In exceptional circumstances, for those patients with thrombocytopenia or bleeding disorders where bleeding may occur following IM administration, JESPECT® may be administered subcutaneously. Subcutaneous administration could lead to a suboptimal response to the vaccine (see section **4.2 Dose and method of administration**).

As with other vaccines, JESPECT® should not be administered in persons with acute severe febrile illness.

Safety and efficacy of JESPECT® have not been established in persons with a history of flavivirus (including JE virus, Yellow Fever virus, Dengue virus and Murray Valley encephalitis virus) infection

and/or vaccination against JE or Yellow fever (see section **4.5 Interactions with other forms of medicines and other forms of interaction**).

Use in children

The safety and efficacy of JESPECT® in persons under 18 years of age have not been established.

Use in elderly patients

Special studies in the geriatric population have not been performed; however, JESPECT® has been administered to 118 subjects ≥65 years of age. No overall differences in safety and effectiveness were observed between these subjects and younger subjects.

Carcinogenicity

JESPECT® has not been evaluated for carcinogenic potential.

Genotoxicity

JESPECT® has not been evaluated for genotoxic potential.

4.5 Interactions with other forms of medicines and other forms of interaction

Concomitant administration of JESPECT® with inactivated hepatitis A vaccine (HAVRIX® 1440) has been explored in one clinical trial (study IC51-308). There was no interference with the immune response to the JE virus and HAV, respectively (see section **5.1 Pharmacodynamic properties**).

Vaccination with JESPECT® in persons with vaccine induced tick born encephalitis (TBE) antibodies resulted in comparable SCR and GMT at Day 56 compared to TBE naive persons.

Vaccination with JESPECT® concurrently with or after administration of other flavivirus vaccines (including yellow fever vaccine) has not been assessed.

If co-administration with other vaccines is indicated, injections should be given into separate limbs. It should be noted that adverse reactions might be intensified.

In patients receiving immunosuppressive therapy or patients with immunodeficiency an adequate immune response may be diminished.

4.6 Fertility, pregnancy and lactation

Use in pregnancy – (Category B1)

In a three phase reproductive study of female rats administered approximately the clinical IM dose of JESPECT® (5 µg) prior to mating (3 weeks (regimen 1 only) and 1 week) and during gestation (day 6), there were no significant toxicological effects in the dams. High JEV-specific antibody titres were detected in maternal blood during gestation (day 5 and 20) in fetal cord blood and at the end of gestation (day 20), and in pups at weaning (lactation day 21).

In a reproductive and pre-/post-natal toxicity study, no vaccine-related effects were detected on reproduction, foetal weight, survival and development of the off-springs. However, incomplete ossification of parts of the skeleton was observed in the group receiving 2 doses, but not in the group receiving 3 doses. There is currently no evidence that this phenomenon is treatment related.

There are limited data from the use of JESPECT® in pregnant women. The vaccine should be used during pregnancy only when clearly needed and the possible advantages outweigh the possible risks for the fetus.

Use in lactation

It is not known whether vaccine antigens or antibodies induced by the vaccine are excreted in human milk. Therefore, caution should be exercised when JESPECT® is administered to a nursing woman.

In a three phase reproductive study of maternal treatment of rats with IM JESPECT® prior to mating and during gestation, assessed to lactation day 21, high JEV-specific antibody titres were detected in maternal blood during gestation, in fetal cord blood at the end of gestation and in pups at weaning (see also section **4.6 Fertility, pregnancy and lactation**).

Adequate human data on JESPECT® use during lactation are not available. JESPECT® should be administered to women who are breastfeeding only when clearly needed.

Effects on fertility

No significant effects on mating performance or fertility were observed in vaccine treated rats administered with IM JESPECT® prior to mating (3 weeks (regimen 1 only) and 1 week) during gestation (day 6). The effect of JESPECT® administration on male fertility has not been evaluated (see also section **4.6 Fertility, pregnancy and lactation**).

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed.

4.8 Undesirable effects

The safety of JESPECT® has been assessed in different controlled clinical trials in which more than 5,405 healthy adults were included of which 4,248 healthy adults received the vaccine.

Approximately 40% of treated subjects can be expected to experience adverse reactions. They usually occur within the first three days after vaccination, are usually mild and disappear within a few days. No increase in the number of undesirable reactions was noted between first and second doses. The incidence of systemic and local symptoms was less prominent after the second vaccination.

Most commonly reported adverse reactions included headache and myalgia occurring in approximately 20% and 13% of subjects, respectively. The treatment-related, treatment-emergent adverse events seen in clinical trials are described in Table 1.

Table 1: Treatment-Related Treatment-Emergent Adverse Events: Pooled Safety Population

TEAE system organ class and preferred term	JESPECT® N=4043 ¹		Placebo N=657 ²	
	n	(%)	n	(%)
Any treatment-related TEAE³	1557	(38.5)	255	(38.8)
Nervous system disorders	806	(19.9)	135	(20.5)
Headache	779	(19.3)	132	(20.1)
Dizziness	16	(0.4)	4	(0.6)
Migraine	11	(0.3)	0	(0)
Paraesthesia	6	0.2	0	(0)
General disorders and administration site conditions⁴	738	(18.3)	119	(18.1)
Fatigue	395	(9.8)	65	(9.9)
Influenza-like illness	363	(9.0)	57	(8.7)
Pyrexia	84	(2.1)	15	(2.3)
Chills	5	(0.1)	3	(0.5)
Malaise	5	(0.1)	0	(0)

Oedema peripheral	1	(0.02)	0	(0)
Injection site reactions				
Injection site haemorrhage	12	(0.3)	0	(0)
Injection site bruising	27	(0.7)	1	(0.2)
Musculoskeletal and connective tissue disorders	541	(13.4)	101	(15.4)
Myalgia	524	(13.0)	94	(14.3)
Musculoskeletal stiffness	5	(0.1)	0	(0)
Pain in extremity	3	(0.1)	0	(0)
Arthralgia	2	(0.1)	0	(0)
Gastrointestinal disorders	228	(5.6)	40	(6.1)
Nausea	193	(4.8)	37	(5.6)
Vomiting	27	(0.7)	7	(1.1)
Diarrhoea	23	(0.6)	4	(0.6)
Abdominal pain	4	(0.1)	0	(0)
Infections and infestations	76	(1.9)	10	(1.5)
Nasopharyngitis	33	(0.8)	4	(0.6)
Rhinitis	15	(0.4)	1	(0.2)
Skin and subcutaneous tissue disorders	59	(1.5)	8	(1.2)
Rash	39	(1.0)	4	(0.6)
Pruritus	4	(0.1)	1	(0.2)
Erythema	3	(0.1)	0	(0)
Urticaria	1	(0.02)	0	(0)
Respiratory, thoracic and mediastinal disorders	32	(0.8)	8	(1.2)
Pharyngolaryngeal pain	14	(0.4)	5	(0.8)
Dyspnoea	2	(0.1)	0	(0)
Investigations	34	(0.8)	3	(0.5)
Increased hepatic enzymes	9	(0.2)	0	(0)
Ear and labyrinth disorders	18	(0.4)	1	(0.2)
Vertigo	16	(0.4)	1	(0.2)
Cardiac disorders	6	(0.2)	0	(0)
Palpitations	2	(0.1)	0	(0)
Tachycardia	2	(0.1)	0	(0)
Blood and lymphatic system disorders	19	(0.5)	5	(0.8)
Lymphadenitis	2	(0.1)	0	(0)
Lymphadenopathy	8	(0.2)	0	(0)
Thrombocytopaenia	1	(0.02)	0	(0)

¹Data taken from different pooled analyses of clinical trials

²Placebo was Phosphate Buffered Saline +aluminum hydroxide adjuvant

³Events with a causality reported as probable or possible or with a missing classification were considered related to study medication.

⁴Only injection site reactions not solicited in subject diaries are displayed here. For solicited local reactions see Table2.

Abbreviations: N=number of subjects in group; n=number of subjects with event; %=percentage of subjects based on number of subjects in the group; TEAE=treatment-emergent adverse event.

The incidence of local symptoms from a pooled safety population is described in Table 2. Please note the results are from subject diary cards on the day of first and second vaccination.

Table 2: Subject Diary Local Tolerability on Day of Vaccination: Pooled Safety Population

Symptom	Vaccination	JESPECT® N=4043 ¹		Placebo N=657 ²	
		n	(%)	n	(%)
Any symptom	First (Day 0)	1342	(35.3)	219	(34.1)
	Second (Day 28)	823	(25.1)	192	(27.6)
Local pain	First	768	(20.2)	129	(20.1)
	Second	469	(13.0)	112	(16.1)
Local tenderness ³	First	693	(19.3)	120	(18.7)
	Second	570	(16.2)	110	(15.8)
Redness	First	174	(4.6)	17	(2.6)
	Second	116	(3.2)	24	(3.5)
Hardening	First	144	(3.8)	19	(3.0)
	Second	114	(3.2)	17	(2.4)
Swelling	First	82	(2.2)	11	(1.7)
	Second	67	(1.8)	11	(1.6)
Itching	First	36	(1.0)	10	(1.6)
	Second	33	(1.0)	11	(1.6)

¹Data taken from different pooled analyses of clinical trials

²Placebo was Phosphate Buffered Saline +aluminum hydroxide adjuvant

³Tenderness was not assessed in study IC51-304, included in the pooled safety population.

Abbreviations: N=number of subjects in group; n=number of subjects with event; %=percentage of subjects based on number of subjects in the respective treatment group with a response (yes/no).

Post marketing experience

The following adverse events have been reported during post marketing use of JESPECT®. Because these events are reported voluntarily, it is not possible to estimate frequency.

Nervous system disorders: paraesthesia, neuritis, syncope.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare professionals are asked to report any suspected adverse reactions <https://nzphvc.otago.ac.nz/reporting/>

4.9 Overdose

There have been no cases of overdosage reported.

Contact the Poisons Information Centre on 0800 POISON (0800 764 766) in New Zealand for advice on overdosage management.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Encephalitis vaccines.

ATC Code: J07BA02

Clinical efficacy and safety

Efficacy (immunogenicity) of JESPECT® has been studied in approximately 2,228 healthy, adult subjects in 10 multi-centre clinical trials conducted in the US and Europe.

Immunogenicity of the vaccine was evaluated in a randomized, active-controlled, observer-blinded, multicenter Phase 3 clinical trial (trial IC51-301) conducted in the US, Germany and Austria in healthy male and female subjects ≥ 18 years of age (mean age: 41.3 years; 60.8% female; race: Caucasian 80.8%, Asian 0.8%, Black/African American 13.1%, and Other 5.3%). Subjects were randomised to receive either two separate doses of 6 $\mu\text{g}/0.5$ mL of JESPECT[®] (on a 0, 28-day schedule by IM injection) (n = 430) or three separate doses of 1.0 mL of the JE vaccine JE-VAX[®] (on a 0, 7 and 28 day schedule by subcutaneous injection) (n = 437). The co-primary endpoint was seroconversion rate (anti-JE virus antibody titer $\geq 1:10$) and geometric mean titer (GMT) at Day 56 as assessed by a Plaque Reduction Neutralisation Test (PRNT) for the entire study population.

By Day 56, the proportion of subjects who had seroconverted was similar for both treatment groups (96.4% vs. 93.8% for JESPECT[®] and JE-VAX[®], respectively) (Table 3). Geometric mean titers increased by Day 56 to 243.6 for JESPECT[®] and to 102.0 for JE-VAX[®], respectively. The immune responses elicited by JESPECT[®] were non-inferior to those induced by JE-VAX[®] (Table 3).

Table 3: Seroconversion Rates and Geometric Mean Titers of JESPECT[®] and JE-VAX[®] in per protocol population (trial IC51-301)

Seroconversion Rate			
Timepoint	JESPECT[®] N=365 n (%)	JE-VAX[®] N=370 n (%)	Risk difference estimator [95% CI]
Visit 0 (Screening)	0	0	
Visit 3 (Day 28)	197 (54)	321 (86.8)	
Visit 4 (Day 56)	352 (96.4)	347 (93.8)	1.05 [-1.33, 3.43]
Geometric Mean Titer (GMT) (by Plaque Reduction Neutralisation Test)			
Timepoint	JESPECT[®] N=365 n (GMT)	JE-VAX[®] N=370 n (GMT)	GMT ratio estimator [95% CI]
Visit 0 (Screening)	365 (5.0)	370 (5.0)	
Visit 3 (Day 28)	363 (17.4)	367 (76.9)	
Visit 4 (Day 56)	361 (243.6)	364 (102.0)	2.3257 [1.9666, 2.7505]

The effect of age on the immune response to JESPECT[®] and JE-VAX[®] was assessed as a secondary endpoint (trial IC51-301), comparing subjects 50 years of age or older (N = 262, mean age 59.8) with those below 50 years of age (N = 605, mean age 33.9). There was no significant difference between seroconversion rates of JESPECT[®] and JE-VAX[®] in subjects aged <50 years compared to those aged ≥ 50 years at Day 28 or Day 56 following vaccination. Geometric mean titers were significantly higher at Day 28 in subjects aged <50 years than those aged ≥ 50 years in the JE-VAX[®] group (80.9 vs. 46.9, $p=0.0236$) but there was no significant difference at Day 56 for this treatment group. In addition there were no significant effects of age on GMT in the group receiving JESPECT[®] and no significant difference between seroconversion rates in subjects aged <50 years compared to those aged ≥ 50 years at Day 28 or Day 56 for either treatment group.

Antibody persistence

Antibody persistence was assessed in an uncontrolled Phase 3 follow-up clinical trial (trial IC51-303) of subjects who completed the pivotal immunogenicity study (trial IC51-301) or the pivotal safety study (trial IC51-302), and who received at least one dose of JESPECT[®]. The primary objective of the study was the evaluation of the immune response (seroconversion rate) to JESPECT[®] 24 months after the first vaccination. Secondary objectives included the evaluation of the immune response to

JESPECT® 6, 12, 24 and 36 months after the first vaccination and to evaluate the safety of JESPECT® during the respective study period (see section **4.8 Undesirable effects**). A total of 3,258 healthy male and female subjects were enrolled (mean age: 35.6 years, 56.5% female, race: Caucasian 90.2%, Asian 1.7%, Black/African American 4.9%, Other 3.2%) of which 2,283 subjects received JESPECT®, 338 subjects received JE-VAX®, and 637 subjects received placebo in the respective previous study. Long-term immunogenicity to JESPECT® was assessed in a subset of 181 subjects up to 24 months, and a subset of 152 subjects at 36 months (ITT population). Immunogenicity data covering a period up to 36 months after the first vaccination are described below:

Seroconversion rates for anti-JE virus antibodies at Months 2, 6, 12, 24 and 36 are summarized in Table 4 for the ITT population. At Month 2, 98.9% of subjects had seroconverted (95% CI: 96.06, 99.70), at Month 6 95.0% of subjects had seroconverted (95% CI: 90.82, 97.36). By Month 12, the percentage of subjects who had seroconverted was 83.4% (95% CI: 77.33, 88.14). Seroconversion at 24 months and 36 months was 81.8% (95% CI: 75.50, 86.71) and 84.9% (95% CI: 78.32, 89.70) respectively.

In the ITT2 population, which is a subset of the ITT population with positive PRNT results at Month 2, per definition, 100% of the subjects (N=179) had seroconverted at Month 2 (95% CI: 97.90%, 100.00%). By Month 6 and Month 12, the number of subjects of the ITT2 population who had seroconverted was 95.5% (95% CI: 91.43, 97.72) and 83.8% (95% CI: 77.70, 88.48) respectively (Table 4). The number of subjects of the ITT2 population who had seroconverted at 24 months was 82.1% ((95% CI: 75.85, 87.04) and at 36 months was 84.8% (95% CI: 78.18, 89.63).

Table 4: Seroconversion Rates at Month 2, 6, 12, 24 and 36 after Vaccination with JESPECT® (ITT and ITT2 Populations) (trial IC51-303)

Time		ITT Population		ITT2 Population	
		N=181 n (%)	95% Confidence Interval	N=179 n (%)	95% Confidence Interval
Month 2	Seroconverted	179 (98.9)	[96.06, 99.70]	179 (100)	[97.90, 100.00]
	Not seroconverted	1 (0.6)			
	Missing	1 (0.6)			
Month 6	Seroconverted	172 (95.0)	[90.82, 97.36]	171 (95.5)	[91.43, 97.72]
	Not seroconverted	9 (5.0)			
Month 12	Seroconverted	151 (83.4)	[77.33, 88.14]	150 (83.8)	[77.70, 88.48]
	Not seroconverted	30 (16.6)			
Month 24	Seroconverted	148 (81.8)	[75.50, 86.71]	147 (82.1)	[75.85, 87.04]
	Not seroconverted	33 (18.2)			
		N=152 n (%)	95% Confidence Interval	N=151 n (%)	95% Confidence Interval
Month 36	Seroconverted	129 (84.9)	[78.32, 89.70]	128 (84.8)	[78.18, 89.63]
	Not seroconverted	23 (15.1)			

Geometric mean titers at Months 2, 6, 12, 24 and 36 after vaccination with JESPECT® are summarized in Table 5. At Month 2, the GMT was 310.8 (95% CI: 268.76, 359.44) which decreased to 83.5 (95% CI: 70.89, 98.38) at Month 6 and to 41.2 (95% CI: 34.39, 49.33) at Month 12 after vaccination with JESPECT®. At Month 24 and 36, the GMT was 44.3 (95% CI: 36.72, 53.44) and 43.8 (95% CI: 36.49, 52.56) respectively.

Table 5: Geometric Mean Titers at Months 2, 6, 12, 24 and 36 after Vaccination with JESPECT® (ITT and ITT2 Populations) (trial IC51-303)

Time	ITT Population		ITT2 Population	
	N=181	95% Confidence Interval	N=179	95% Confidence Interval
Month 2	310.8	[268.76, 359.44]	318.1	[276.83, 365.43]
Month 6	83.5	[70.89, 98.38]	84.9	[72.18, 99.95]
Month 12	41.2	[34.39, 49.33]	41.6	[34.73, 49.87]
Month 24	44.3	[36.72, 53.44]	44.9	[37.15, 54.14]
	N=152	95% Confidence Interval	N=151	95% Confidence Interval
Month 36	43.8	[36.49, 52.56]	43.5	[36.21, 52.23]

The results were confirmed in the ITT2 population.

Vaccination with inactivated JE vaccine is known to produce antibodies that decline with time in a proportion of the vaccinated population. The observed decline in GMT is therefore as expected and compares well with data from other inactivated JE vaccines.

Results of another Phase 3 follow-up clinical trial (trial IC51-305) support these findings. In this study, 58.3% and 48.3% of subjects who received the recommended primary vaccination series (see section **4.2 Dose and method of administration**) showed persisting protective antibodies 12 and 24 months (respectively) after completion of primary vaccination. See also section **5.1**

Pharmacodynamic properties.

Booster immunisation

The effect of booster vaccination of JESPECT® on long term immunogenicity was investigated in an uncontrolled, open label, Phase 3 clinical trial in healthy male and female adults (trial IC51-311). Subjects (n=198) were administered one 6 µg/0.5 mL dose of JESPECT® 15 months after their first dose (Day 0) of the recommended primary vaccination series (see section **4.2 Dose and method of administration**). The primary endpoint was assessment of the effect of booster vaccination on immunogenicity in terms of seroconversion rate 12 months after administration of the booster dose (equivalent to 27 months after first dose (Day 0) of the recommended primary vaccination series). Secondary endpoints included assessment of seroconversion 28 days and 6 months after administration of the booster dose, and GMT 28 days, 6 months and 12 months after booster vaccination (equivalent to 16, 21 and 27 months after first dose (Day 0) of the recommended primary vaccination series).

Pre-booster vaccination, 69.2% of subjects had seroconverted and GMT was 22.5 (95% CI: 19.0, 26.7).

Seroconversion was 100% at 28 days after administration of the booster vaccination and remained high (98.5%) to 12 months after administration of the booster vaccination (equivalent to 27 months after first dose (Day 0) of the recommended primary vaccination series) (Table 6).

Geometric mean titers were highest 28 days after receiving the booster vaccination and declined to 361.4 after 12 months (equivalent to 27 months after first dose (Day 0) of the recommended primary vaccination series) (Table 6).

Table 6: Seroconversion rates and Geometric Mean Titers at Day 28 and Months 6 and 12 after Booster Vaccination with JESPECT® (ITT Population) (trial IC51-311)

	Seroconversion rate		GMT	
	%	95% CI		95% CI
Pre-booster (n=198)	69.2	[62.4, 75.2]	22.5	[19.0, 26.7]
Day 28 (n=198)	100.0	[98.1, 100.0]	900.1	[742.4, 1091.3]
Month 6 (n=197)	98.5	[95.6, 99.5]	487.4	[390.7, 608.1]
Month 12 (n=194)	98.5	[95.6, 99.5]	361.4	[294.5, 443.5]

Incomplete primary immunisation

Immunogenicity following incomplete primary vaccination was investigated in a non-randomized, open label, multicenter Phase 3 clinical trial (trial IC51-305) conducted in healthy male and female adults (n=349). Subjects received either the recommended primary vaccination schedule (see section **4.2 Dose and method of administration**; one 6 µg/0.5 mL dose of JESPECT® on Day 0 and a second 6 µg/0.5 mL dose of JESPECT® on Day 28) (n=116), or one single dose of 6 µg/0.5 mL dose of JESPECT® (on Day 0) (n=117), or one 12 µg/0.5 mL dose of JESPECT® (on Day 0) (n=116). Those subjects that were seronegative (PRNT₅₀ titers < 1:10) at Month 6 received a single 6 µg/0.5 mL booster dose at 11 months after the first dose (Day 0). Those subjects that were seronegative at Month 12 received a single 6 µg/0.5 mL booster dose at 23 months after the first dose (Day 0).

Administration of the second injection of the primary immunisation series up to 11 months after the first dose results in 99% seroconversion and GMT of 504.3 (95% CI 367.3, 692.3) (Table 7).

Table 7: Seroconversion rates and Geometric Mean Titers 4 weeks after a single 6 µg/0.5mL booster dose is administered to subjects that are seronegative at Month 11 or Month 23 after complete primary immunisation (2 x 6 µg/0.5 mL) or incomplete primary immunisation (1 x 6 µg/0.5 mL) with JESPECT® (ITT Population) (trial IC51-305)

	SCR (%)	GMT	
			95% CI
Complete primary immunisation (2 x 6 µg/0.5 mL dose)			
+Booster at Month 11 (n=17)	100	673.6	[378.7, 1198.2]
+ Booster at Month 23 (n=27)	100	2536.7	[1467.7, 4384.4]
Incomplete primary immunisation (1 x 6 µg/0.5 mL dose)			
+Booster at Month 11(n=100)	99	504.3	[367.3, 692.3]
+Booster at Month 23 (n=5)	100	571.4	[88.2, 3702.9]

Concomitant use

The concomitant use of JESPECT® with inactivated Hepatitis A Virus (HAV) vaccine (HAVRIX® 1440) has been explored in one clinical trial (study IC51-308). There was no interference with the immune response to the JE virus and HAV, respectively. Concomitant administration of JESPECT® and inactivated hepatitis A vaccine was shown to be non-inferior to single vaccinations with regard to GMT of anti-JE virus neutralising antibody and HAV antibody, and for seroconversion rates of both antibody types.

5.2 Pharmacokinetic properties

Evaluation of pharmacokinetic properties is not required for vaccines.

5.3 Preclinical safety data

Animal studies in mice, rats and rabbits have shown that the JESPECT[®] vaccine induces the immune system to produce neutralising antibodies against Japanese encephalitis virus which correlate with protection and survival. Mice treated with JESPECT[®] vaccine or human JESPECT[®] antisera demonstrated protection against lethal Japanese encephalitis challenge from several different (SA₁₄, Beijing and KE-093) strains in a generally dose- or titre-dependent manner, respectively.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Aluminium hydroxide hydrate

Phosphate Buffered Saline consisting of:

Sodium chloride – 9mg/mL

Monobasic potassium phosphate 0.144mg/mL

Dibasic sodium phosphate 0.795 mg/mL

Water for injections

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

3 years

6.4 Special precautions for storage

Store in a refrigerator at 2° to 8° C. **Do not freeze.** Store in the original package in order to protect from light.

6.5 Nature and contents of container

JESPECT[®] is supplied as a 0.5 mL suspension in a pre-filled syringe (Type I glass) with a plunger stopper (chlorobutyl elastomer) in a pack size of 1 needle-less syringe.

6.6 Special precautions for disposal and other handling

JESPECT[®] pre-filled syringe is for single use only in one individual only. Inject the entire contents of the syringe.

Do not use if the package is torn or damaged.

Do not use the vaccine after the expiration date shown on the label.

Upon storage, a fine white deposit with a clear colourless supernatant can be observed. Shake well before use.

7. MEDICINE SCHEDULE

Prescription Only Medicine

8. SPONSOR

Seqirus (NZ) Ltd

PO Box 62590

Greenlane

Auckland 1546

Telephone: 0800 502 757

9. DATE OF FIRST APPROVAL

18 October 2012

10. DATE OF PREPARATION

9 February 2021

JESPECT® is a registered trademark of Valneva Scotland Ltd.

Summary table of changes

Section changed	Summary of new information
2	Update to excipient names
6.2	Update to excipient names inclusion of formulation details.
4.8	Spelling correction