

NEW ZEALAND DATA SHEET

1. PRODUCT NAME

Linezolid Juno 2 mg/mL solution for infusion

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 1 mL of Linezolid solution for infusion contains 2 mg of linezolid. 300ml infusion bags contain 600mg linezolid.

For full list of excipients, see **section 6.1**.

3. PHARMACEUTICAL FORM

Solution for infusion.

Isotonic, clear, colourless to yellow solution.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Linezolid solution for infusion is indicated for the treatment of infections when known or suspected to be caused by susceptible organisms including those associated with concurrent bacteraemia such as:

- Pneumonia - community acquired and nosocomial pneumonia
- Severe skin and soft tissue infections
- Enterococcal infections.

Linezolid is active against Gram-positive bacteria only. Linezolid has no clinical activity against Gram-negative pathogens. Specific Gram-negative therapy is required if a concomitant Gram-negative pathogen is documented or suspected.

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Dose and method of administration

Patients who commence treatment on the parenteral formulation may be switched to an oral presentation when clinically indicated. In such circumstances, no dose adjustment is required as linezolid has an oral bioavailability of approximately 100%.

The injection should be administered over a period of 30 to 120 minutes.

Adults and children 12 years or older

The recommended dosage should be administered intravenously (IV) as shown in Table 1. Duration of treatment is variable. It is dependent on the pathogen, the site of infection and its severity, and on the patient's clinical response. To date, the maximum treatment duration in controlled clinical trials has been 28 days. No increase in the recommended dosage or duration of treatment is required for infections associated with concurrent bacteraemia.

Table 1. Dosage guidelines for Linezolid IV Injection

Infections (including those associated with concurrent bacteraemia)	Twice daily dosage and route of administration	Duration of treatment
Community acquired pneumonia	600 mg IV	10-14 consecutive days
Nosocomial pneumonia		
Severe skin and soft tissue infections	600 mg IV	
Enterococcal infections	600 mg IV	14-28 consecutive days

Children less than 12 years old

The recommended dosage should be administered intravenously (IV) as shown in Table 2. As for adults and adolescents, the duration of treatment is variable. It is dependent on the pathogen, the site of infection and its severity, and on the patient's clinical response.

Table 2. Dosage guidelines for Linezolid for paediatric patients from birth through 11 years of age

Infections (including those associated with concurrent bacteraemia)	Dosage for paediatric patients from birth through 11 years of age [§]	Duration of treatment
Community acquired pneumonia	10 mg/kg IV once every 8 hours	10 to 14 consecutive days
Nosocomial pneumonia		
Skin and soft tissue infections		
Enterococcal infections	10 mg/kg IV every 8 hours	14 to 28 consecutive days

[§] Neonates <7 days: most pre-term neonates < 7 days of age (gestational age < 34 weeks) have low systemic linezolid clearance values and large AUC values than many full-term neonates and older infants. These neonates should be initiated with dosing regimes of 10 mg/kg every 12 hours. Consideration may be given to the use of 10 mg/kg every 8 hours regime in neonates with sub-optimal clinical response. All neonatal patients should receive 10 mg/kg every 8 hours by 7 days of life (see **section 5.2**).

Dosage adjustments in special populations

No dose adjustment is required in the elderly, in patients with impaired hepatic function or impaired renal function. However, linezolid should be administered after haemodialysis in patients receiving such treatment (refer **section 5.2**).

4.3 Contraindications

Hypersensitivity to linezolid or to any of the excipients in the relevant pharmaceutical form (refer **section 2**).

Monoamine Oxidase Inhibitors

Linezolid should not be used in patients taking any medicinal product which inhibits monoamine oxidases A or B (*e.g.*, phenelzine) or within two weeks of taking any such medicinal product.

Potential Interactions Producing Elevation of Blood Pressure

Unless patients are monitored for potential increases in blood pressure, linezolid should not be administered to patients with uncontrolled hypertension, pheochromocytoma, thyrotoxicosis and/or patients taking any of the following types of medications: directly and indirectly acting

sympathomimetic agents (e.g., pseudoephedrine), vasopressive agents (e.g., adrenalin, noradrenalin), dopaminergic agents (e.g., dopamine, dobutamine) (see **section 4.4**).

Potential Serotonergic Interactions

Unless patients are carefully observed for signs and/or symptoms of serotonin syndrome, linezolid should not be administered to patients with carcinoid syndrome and/or patients taking any of the following medications: serotonin re-uptake inhibitors, tricyclic antidepressants, serotonin 5-HT₁ receptor agonists (triptans), pethidine or buspirone (see **section 4.5**).

4.4 Special warning and precautions for use

It is recommended that therapy with linezolid should be initiated in a hospital environment following guidance from appropriate specialists.

Myelosuppression (including anaemia, leucopenia, pancytopenia and thrombocytopenia) has been reported in patients receiving linezolid. In cases where the outcome is known, the affected haematological parameters have risen towards pre-treatment levels when linezolid was discontinued. Complete blood counts should be monitored weekly in patients who receive linezolid for longer than two weeks, particularly those with pre-existing myelosuppression, those receiving concomitant drugs that produce bone marrow suppression, or those with a chronic infection who have received previous antibiotic therapy. Discontinuation of therapy should be considered in patients who develop or who have a worsening of myelosuppression.

Controlled clinical trials did not include patients with diabetic foot lesions, decubitus, or ischaemic lesions, severe burns or gangrene. Therefore, experience in the use of linezolid in the treatment of these conditions is limited.

Peripheral neuropathy and optic neuropathy have been reported in patients treated with linezolid, primarily those patients treated for longer than the maximum recommended duration of 28 days. In cases of optic neuropathy that progressed to loss of vision, patients were treated for extended periods beyond the maximum recommended duration.

If symptoms of visual impairment appear, such as changes in visual acuity, changes in colour vision, blurred vision, or visual field defect, prompt ophthalmic evaluation is recommended.

Visual function should be monitored in all patients taking linezolid for extended periods (greater than or equal to 3 months) and in all patients reporting new visual symptoms regardless of length of therapy with linezolid. If peripheral or optic neuropathy occurs, the continued use of linezolid in these patients should be weighed against the potential risks.

Lactic acidosis has been reported with the use of linezolid. Patients who develop recurrent nausea or vomiting, unexplained acidosis, or a low bicarbonate level while receiving linezolid should receive immediate medical attention. Convulsions have been reported to occur rarely in patients when treated with linezolid. In most of these cases, a history of seizures or risk factors for seizures were reported.

Antibiotic associated pseudomembranous colitis has been reported with nearly all antibacterial agents including linezolid. A toxin produced with *Clostridium difficile* appears to be the primary cause. The severity of the colitis may range from mild to life threatening. It is important to consider this diagnosis in patients who develop diarrhoea or colitis in association with antibiotic use (this may occur up to several weeks after cessation of antibiotic therapy). Mild cases usually respond to drug discontinuation alone. However, in moderate to severe cases appropriate therapy with a suitable oral

antibacterial agent effective against *Clostridium difficile* should be considered. Fluids, electrolytes and protein replacement should be provided when indicated. Drugs which delay peristalsis, *e.g.*, opiates and diphenoxylate with atropine may prolong and/or worsen the condition and should not be used. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. The use of antibiotics may occasionally result in an overgrowth of non-susceptible organisms. Should superinfection occur during therapy, appropriate measures should be taken.

The safety and effectiveness of linezolid when administered for periods longer than 28 days have not been established. In healthy volunteers, coadministration of rifampin with linezolid resulted in a 21% decrease in linezolid C_{max} and a 32% decrease in linezolid AUC (see **section 4.5**). The clinical significance of this interaction is unknown.

Linezolid has no clinical activity against Gram-negative pathogens and is not indicated for the treatment of Gram-negative infections. Specific Gram-negative therapy is required if a concomitant Gram-negative pathogen is documented or suspected. Linezolid should be used with special caution in patients at high risk for life threatening systemic infections, such as those with infections related to central venous catheters in intensive care units. Linezolid is not approved for the treatment of patients with catheter-related bloodstream infections.

Linezolid should be used with special caution in patients with severe renal insufficiency and only when the anticipated benefit is considered to outweigh the theoretical risk. It is recommended that linezolid should be used in patients with severe hepatic insufficiency only when the anticipated benefit is considered to outweigh the theoretical risk.

Mortality in subjects with catheter-related infections. An open-label, randomized clinical trial was conducted in adult patients with catheter-related Gram-positive bloodstream infections comparing linezolid (600 mg q12h IV/PO) to vancomycin 1 g IV q12h or oxacillin 2 g IV q6h/dicloxacillin 500 mg PO q6h with a treatment duration of 7 to 28 days. The mortality rates in this study were 78/363 (21.5%) and 58/363 (16.0%) on linezolid and the comparator, respectively. Based on results from a logistic regression, the estimated odds ratio is 1.426 [95%CI 0.970, 2.098]. While causality has not been established, this observed imbalance occurred primarily in linezolid-treated patients in whom either Gram-negative pathogens, mixed Gram-negative and Gram-positive pathogens, or no pathogen were identified at baseline.

Patients randomized to linezolid who had only a Gram-positive infection at baseline, including the subgroup of patients with Gram-positive bacteremia experienced a survival rate similar to the comparator.

Carcinogenicity and mutagenicity

Lifetime studies in animals have not been conducted to evaluate the carcinogenic potential of linezolid. There was no evidence of genotoxicity in tests for gene mutations (bacteria and Chinese hamster ovary cells), chromosomal changes (human lymphocytes *in vitro* and mouse micronucleus assay *in vivo*) and DNA damage (unscheduled DNA synthesis *in vitro*).

Use in paediatric patients

The clearance of linezolid is most rapid in the youngest age groups (excluding neonates less than 1 week old), resulting in a shorter half-life. As children mature, the clearance of linezolid gradually decreases and by adolescence the clearance values approach those observed for the adult population. While drug clearance in adolescents (ages 12 through 17 years) is usually similar to the clearance in adults, there is wider inter-subject variation in this age group compared with adults (refer **section 5.2**). Results of clinical studies showed similar efficacy in adult and adolescent patients. Given the wider inter subject variation in adolescents, the slight possibility that high clearance may result in decreased efficacy in some adolescent patients should be considered. The dosage for paediatric patients younger than 12 years of age should be 10 mg/kg every 8 hours, while children 12 years and older should receive the same dose as adult patients, 600 mg every 12 hours (refer **section 4.2**).

In limited clinical experience, 5 out of 6 (83%) paediatric patients with infections due to Gram-positive pathogens with MICs of 4 µg/mL treated with linezolid had clinical cures. However, paediatric patients with a sub-optimal clinical response, particularly those with pathogens with MIC of 4 µg/mL, lower systemic exposure, site and severity of infection and the underlying medical condition should be considered when assessing clinical response (see **section 4.2** and **section 5.2**).

4.5 Interactions with other medicines and other forms of interaction

Linezolid is not detectably metabolised by the cytochrome P450 (CYP) enzyme system and it does not induce or inhibit the activities of clinically significant human CYP isoforms (1A2, 2C9, 2C19, 2D6, 2E1, 3A4). Therefore, no CYP450-induced drug interactions are expected with linezolid. Drugs such as warfarin and phenytoin, which are CYP2C9 substrates, may be given with linezolid without changes in dosage regimen.

No interactions have been observed in pharmacokinetic studies with either aztreonam or gentamicin.

The effect of rifampin on the pharmacokinetics of linezolid was studied in sixteen healthy adult male volunteers administered linezolid 600 mg twice daily for 2.5 days with and without rifampin 600 mg once daily for 8 days. Rifampin decreased the linezolid C_{max} and AUC by a mean 21% [90% CI, 15, 27] and a mean 32% [90% CI, 27, 37], respectively. The mechanism of this interaction and its clinical significance are unknown (see **section 4.4**).

Linezolid is a reversible, non-selective inhibitor of monoamine oxidase (MAOI). Limited clinical studies have shown that coadministration of linezolid with either pseudoephedrine or phenylpropanolamine resulted in mild, reversible enhancement of the pressor responses in normotensive patients. Similar studies in hypertensive subjects have not been conducted. The potential for interaction with sympathomimetic and adrenergic agents should be considered. Initial doses of potent vasopressors, such as dopamine and adrenaline, should be reduced and carefully titrated to achieve the desired response when coadministered with linezolid.

No significant pressor response was observed in subjects receiving both linezolid and less than 100 mg tyramine. This suggests that it is only necessary to avoid ingesting excessive amounts of food and beverages with a high tyramine content (e.g., mature cheese, yeast extracts, undistilled alcoholic beverages and fermented soya bean products such as soya sauce).

Linezolid has the potential for interaction with serotonergic agents. Limited clinical studies have shown that coadministration of linezolid with dextromethorphan was not associated with serotonin syndrome effects (e.g., confusion, delirium, restlessness, tremors, blushing, diaphoresis and

hyperpyrexia). The effects of other serotonin uptake inhibitors have not been studied. Physicians should be alert to the possible signs and symptoms of serotonergic syndrome in patients receiving concomitant linezolid and serotonergic agents.

Very rare spontaneous reports of serotonin syndrome with co-administration of linezolid and serotonergic agents have been reported. If the concomitant serotonergic agent is withdrawn, discontinuation symptoms can be observed.

4.6 Fertility, pregnancy and lactation

Use in Fertility

Whilst linezolid did not affect female rat fertility or reproductive performance, it reversibly decreased the fertility of adult male rats at oral doses of 50 mg/kg/day with exposure levels approximately equal to those expected in humans.

Epithelial hypertrophy of the epididymis may have contributed to the decreased fertility by affecting sperm maturation. However, an effect on spermatogenesis cannot be excluded as delayed spermiation in the testes occurred at 100 mg/kg/day (twice the clinical exposure). Sperm counts in the testis were unaffected but sperm counts in the cauda epididymis were decreased and sperm from the vas deferens had decreased motility. Most sperm from the epididymis in rats treated with 100 mg/kg/day had detached head/neck from the tail.

Epithelial hypertrophy was not observed in beagle dogs which suggests that the above effects are species specific to rats.

Slightly decreased fertility occurred in juvenile male rats treated orally with linezolid at 50 mg/kg/day from 7 days to 42 days old and at 100 mg/kg/day from 43 days to 55 days old.

Delayed spermatid development was observed in juvenile rats treated with linezolid at 63 mg/kg/day and single-cell spermatocyte/spermatid degeneration or necrosis (apoptosis) was observed in juvenile rats treated with linezolid at 100 mg/kg/day (all reversible).

Use in pregnancy

There are no adequate data from the use of linezolid in pregnant women. Studies in animals have shown reproductive effects (see below). The potential risk for humans is unknown.

Linezolid should not be used during pregnancy unless clearly necessary *i.e.*, only if the potential benefit outweighs the potential risk.

Linezolid and/or its metabolites crossed the placenta in rats. Linezolid was not teratogenic in mice or rats at exposure levels 4 times (mice) or equivalent to (rats) the expected human exposure level, based on AUCs.

Embryofetal effects were observed in mice at 450 mg/kg/day (4 times the clinical exposure based on AUC) and in rats at 15 mg/kg/day (0.14 times the clinical exposure based on AUC). Decreased fetal weights and delayed ossification occurred in rats without maternal toxicity. In mice, increased embryo death including total litter loss, decreased fetal body weights and an exacerbation of the normal genetic predisposition to sternal variations in the strain of mice used were seen at doses causing maternal toxicity (clinical signs and decreased body weight gain).

Use in lactation

Animal data suggest that linezolid is likely to pass into breast milk. Breastfeeding should be discontinued prior to administration.

Linezolid and its metabolites were excreted into the milk of rats. The concentration of total drug-related materials in milk was similar to or greater than that in maternal plasma. The development of pups from rats treated orally with 50 mg/kg/day linezolid during gestation and lactation (0.6 times the clinical exposure based on AUC) was slightly delayed, manifest as decreased body weight gain, delayed pinna detachment and balanopreputial separation and decreased negative geotaxis response. These pups when allowed to mature showed slightly decreased fertility, increased implantation loss and decreased epididymides and testes weights.

4.7 Effects on ability to drive and use machines

The effect of linezolid on the ability to drive or operate machinery has not been systematically evaluated.

4.8 Undesirable effects

Clinical Trials

The information provided is based on data generated from clinical studies in adults and paediatric patients.

Adults

More than 2,000 adult patients received the recommended linezolid doses for up to 28 days. In these studies, the majority of adverse reactions to linezolid were of mild to moderate intensity, of limited duration and did not require discontinuation of treatment. The adverse reactions were not dose dependent.

Approximately 22% of patients experienced adverse reactions; those most commonly reported were headache, diarrhoea, nausea and candidiasis (particularly oral and vaginal candidiasis; see table below). The most commonly reported drug-related adverse events which led to discontinuation of treatment were headache, diarrhoea, nausea and vomiting. Table 3 shows the incidence of adverse reactions reported in at least 1% of patients in these trials.

Table 3. Incidence (%) of adverse reactions reported in $\geq 1\%$ of patients in comparator-controlled clinical trials with linezolid and 600 mg bid patients in the VRE dose-response study

Event	Linezolid (n=2125)	All comparators* (n=2001)
Gastrointestinal disorders		
Diarrhoea	4.2	3.2
Nausea	3.3	2.3
Vomiting	1.2	0.4
Abnormal liver function tests	1.0	0.3
General body Headache	2.1	1.3
Special senses		
Taste perversion	1.1	0.7
Urogenital		
Vaginal candidiasis	1.1	0.6

*Comparators included cefpodoxime proxetil, ceftriaxone, clarithromycin, dicloxacillin, oxacillin and vancomycin

Changes observed in laboratory parameters (without regard to drug relationship) generally reflected resolution of the infection, were not clinically significant, did not lead to discontinuation of therapy and were reversible. The incidence of patients with at least one substantially abnormal haematologic or serum chemistry value is presented in Table 4.

Table 4. Percent of patients who experienced at least one substantially abnormal* haematology or chemistry laboratory value in comparator-controlled clinical trials with linezolid

Laboratory assay	Linezolid	All comparators ^a
Haemoglobin	5.4	4.8
Platelet count	2.4	1.5
Leucocytes	1.6	1.1
Neutrophils	0.8	0.9
AST	4.1	5.3
ALT	7.4	7.2
LDH	1.4	1.1
Alkaline phosphatase	2.6	2.3
Lipase	3.9	3.7
Amylase	1.8	1.5
Total bilirubin	0.7	0.8
BUN	1.6	1.1
Creatinine	0.2	0.5

*Haematology:

< 75% (< 50% for neutrophils) of Lower Limit of Normal (LLN) for values normal at baseline

< 75% (< 50% for neutrophils) of LLN and of baseline for values abnormal at baseline

Chemistry:

> 2 Upper Limit of Normal (ULN) for values normal at baseline > 2 ULN and > 2 x baseline for values

abnormal at baseline ^aComparators included clarithromycin, cefpodoxime proxetil, ceftriaxone,

dicloxacillin, oxacillin and vancomycin

Paediatric patients

The safety of linezolid formulations was evaluated in 215 paediatric patients ranging in age from birth through 11 years and in 248 paediatric patients aged 5 through 17 years (146 of these, 248 were age 5 through 11 and 102 were age 12 to 17). These patients were enrolled in two phase 3 comparator-controlled clinical trials and were treated for up to 28 days. In these studies 83% and 99% respectively, of the adverse events reported with linezolid were described as mild to moderate in intensity. In the study of hospitalised paediatric patients (birth through 11 years) with Gram-positive infections, who were randomised 2 to 1 (linezolid: vancomycin), mortality was 6.0% (13/215). In the linezolid arm and 3.0% (3/101) in the vancomycin arm. However, given the severe underlying illness in the patient population, no causality could be established. Table 5 shows the incidence of drug-related adverse events reported in more than 1% of paediatric patients (and more than 1 patient) in either treatment group in the comparator controlled phase 3 trials.

Table 5. Incidence (%) of drug related adverse events occurring in >1% of paediatric patients (and >1 patient) in either treatment group in comparator-controlled clinical trials

Event	Uncomplicated skin and skin structure infections [†]		All other indications [‡]	
	Linezolid (n= 248)	Cefadroxil (n= 251)	Linezolid (n= 215)	Vancomycin (n=101)
% of patients with 1 drug related adverse event	19.2	14.1	18.8	34.3
% of patients discontinuing due to a drug-related adverse event.	1.6	2.4	0.9	6.1
Diarrhoea	5.7	5.2	3.8	6.1
Nausea	3.3	2.0	1.4	0
Headache	2.4	0.8	0	0
Loose stools	1.2	0.8	1.9	0
Thrombocytopenia	0	0	1.9	0
Vomiting	1.2	2.4	1.9	1.0
Generalised abdominal pain	1.6	1.2	0	0
Localised abdominal pain	1.6	1.2	0	0
Anaemia	0	0	1.4	1.0
Eosinophilia	0.4	0.4	1.4	0
Rash	0.4	1.2	1.4	7.1
Vertigo	1.2	0.4	0	0
Oral moniliasis	0	0	0.9	4.0
Fever	0	0	0.5	3.0
Pruritus non-application site	0.4	0	0	2.0
Anaphalaxis	0	0	0	10.1*

[†]Patients 5 through 11 years of age receive linezolid 10mg/kg PO q12h or cefadroxil 15 mg/kg PO q12h.

[‡]Patients 12 years or older received linezolid 600 mg PO Q12h or cefadroxil 500 mg PO q12h.

[‡]Patients from birth through 11 years received linezolid 10 mg/kg IV/PO or vancomycin 10 to 15 mg/kg IV q6-24, depending on age and renal clearance.

* These reports were of “red-man syndrome”, which were coded as anaphylaxis.

In a study of severely ill, hospitalised paediatric patients ranging in age from birth through 11 years, the percentage of patients who developed a substantially low platelet count was 12.9% with linezolid and 13.4% with vancomycin. In an outpatient study of paediatric patients aged from 5 through 17 years, the percentage of patients who developed a substantially low platelet count was 0% with

linezolid and 0.4% with cefadroxil. Other changes observed in laboratory parameters, were not clinically significant, did not lead to discontinuation of therapy and were reversible. The incidence of paediatric patients with at least one substantially abnormal haematologic or serum chemistry value is presented in Table 6.

Table 6. Percent of paediatric patients who experienced at least one substantially abnormal* haematology or serum chemistry laboratory value in comparator controlled clinical trials with linezolid

Laboratory Assay	Uncomplicated skin and skin structure infections [†]		All other indications [‡]	
	Linezolid	Cefadroxil	Linezolid	Vancomycin
Haemoglobin (g/dL)	0.0	0.0	15.7	12.4
Platelet count (x 10 ³ /mm ³)	0.0	0.4	12.9	13.4
WBC (x 10 ³ /mm ³)	0.8	0.8	12.4	10.3
Neutrophils (x 10 ³ /mm ³)	1.2	0.8	5.9	4.3
ALT (U/L)	0.0	0.0	10.1	12.5
Lipase (U/L)	0.4	1.2	---	---
Amylase (U/L)	---	---	0.6	1.3
Total bilirubin (mg/dL)	---	---	6.3	5.2
Creatinine (mg/dL)	0.4	0.0	2.4	1.0

*Haematology:

< 75% (< 50% for neutrophils) of Lower Limit of Normal (LLN) for values normal at baseline

< 75% (< 50% for neutrophils, <90% for haemoglobin) of LLN and of baseline for values abnormal at baseline

Serum chemistry:

> 2 Upper Limit of Normal (ULN) for values normal at baseline

> 2 ULN and > 2 x baseline for values abnormal at baseline

Dosage

[†]Patients 5 through 11 years of age receive linezolid 10mg/kg PO q12h or cefadroxil 15 mg/kg PO q12h.

[†]Patients 12 years or older received Linezolid 600 mg PO Q12h or cefadroxil 500 mg PO q12h.

[‡]Patients from birth through 11 years received linezolid 10 mg/kg IV/PO or vancomycin 10 to 15 mg/kg IV q6-24, depending on age and renal clearance.

Post-marketing surveillance

Myelosuppression (including anaemia, leucopenia, pancytopenia and thrombocytopenia) has been reported.

Peripheral neuropathy, and optic neuropathy sometimes progressing to loss of vision, have been reported in patients treated with linezolid. These reports have primarily been in patients treated for longer than the maximum recommended duration of 28 days (see **section 4.4**).

Lactic acidosis (see Precautions), rash, convulsions, angioedema and anaphylaxis have been reported.

Very rare reports of bullous skin disorders such as those described as Stevens Johnson syndrome have been received.

Gastrointestinal Disorders: Tongue discoloration. Superficial tooth discoloration has been reported very rarely with the use of linezolid. The discoloration was removable with professional dental cleaning (manual descaling) in cases with known outcome.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare professionals are asked to report any suspected adverse reactions <https://nzphvc.otago.ac.nz/reporting/>

4.9 Overdosage

No cases of overdose have been reported. Symptomatic and supportive care is advised together with maintenance of glomerular filtration. Approximately 30% of a linezolid dose is removed during 3 hours of haemodialysis. No data are available for the removal of linezolid by peritoneal dialysis or haemoperfusion.

For advice on the management of overdose please contact the National Poisons Centre on 0800 POISON (0800 764766)

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

General Properties

Linezolid is a synthetic, antibacterial agent belonging to a new class of antibiotics, the oxazolidinones, with *in vitro* activity against Gram positive aerobic bacteria, some Gram positive anaerobic bacteria and certain Gram negative bacteria. It selectively inhibits bacterial protein synthesis via a mechanism of action different from that of other antibacterial agents. Linezolid binds to the 23S ribosomal RNA of the 50S subunit of the bacterial ribosome and prevents the formation of a functional 70S initiation complex which is an essential component of the bacterial translation process. The results of time-kill studies have shown linezolid to be bacteriostatic against enterococci and staphylococci. For streptococci, linezolid was found to be bactericidal for the majority of strains.

Breakpoints

The MIC breakpoints in Table 7 separate susceptible from non-susceptible isolates.

Table 7. MIC breakpoints for linezolid

Pathogen	Susceptibility interpretive criteria					
	MIC in micrograms/mL			Disk diffusion (zone diameters in mm)		
	S	I	R	S	I	R
<i>Enterococcus</i> species	≤ 2	4	≥ 8	≥ 23	21-22	≤ 20
<i>Staphylococcus</i> species	≤ 4	---*	---	≥ 21	---	---*
<i>Streptococcus pneumoniae</i>	≤ 2	---	---	≥ 21	---	---*
<i>Streptococcus</i> species other than <i>S. pneumoniae</i>	≤ 2	---	---	≥ 21	---	---*

*The current absence of data on resistant strains precludes defining categories other than “susceptible”. Strains yielding results suggestive of a “non-susceptible” category should be re-tested and, if confirmed, the isolate should be submitted to a reference laboratory for further testing.

S = susceptible

I = intermediately susceptible

R = resistant

The studies used to define the above breakpoints employed standard NCCLS (National Committee for Clinical Laboratory Standards) microdilution and agar diffusion methods.

Susceptibility

Prevalence of resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. Therefore, the following information gives only an approximate guidance on the probabilities as to whether or not microorganisms will be susceptible to linezolid. Only microorganisms relevant to the given clinical indications are presented here. An asterisk indicates that clinical efficacy has been demonstrated for susceptible isolates in approved clinical indications.

Susceptible organisms

Gram positive aerobes

Corynebacterium jeikeium

*Enterococcus faecalis** (including glycopeptide resistant strains)

*Enterococcus faecium** (including glycopeptide resistant strains)

Enterococcus casseliflavus

Enterococcus gallinarum

Listeria monocytogenes

Staphylococcus aureus (including methicillin resistant strains)*

Staphylococcus aureus (including glycopeptide intermediate resistant strains)

Staphylococcus epidermidis (including methicillin resistant strains)*

Staphylococcus haemolyticus

Staphylococcus lugdunensis

*Streptococcus agalactiae**

Streptococcus intermedius

Streptococcus pneumoniae (including penicillin intermediate and resistant strains)*

*Streptococcus pyogenes**

Viridans group streptococci

Group C streptococci

Group G streptococci

Gram negative aerobes

Pasteurella canis

Pasteurella multocida

Gram positive anaerobes

Clostridium perfringens

Peptostreptococcus anaerobius

Peptostreptococcus species

Gram negative anaerobes

Bacteroides fragilis

Prevotella species

Other

Chlamydia pneumoniae

Intermediately susceptible organisms

Legionella species

Moraxella catarrhalis
Resistant organisms
Haemophilus influenzae
Neisseria species
Enterobacteriaceae
Pseudomonas aeruginosa

Resistance

The mechanism of action of linezolid differs from that of other classes of antibiotics and cross-resistance between linezolid and other classes of antibiotics is unlikely.

Resistance to linezolid developed under selective pressure *in vitro* and was associated with point mutations in the 23S ribosomal RNA. Spontaneous resistance occurs at frequencies of less than 10⁻⁹ *in vitro*. In clinical trials, resistance to linezolid developed in 6 patients infected with *E. faecium* (4 patients received 200 mg twice daily, lower than the recommended dose, and 2 patients received 600 mg twice daily). In a compassionate use program, resistance to linezolid developed in 8 patients with *E. faecium* and in 1 patient with *E. faecalis*. All patients had either unremoved prosthetic devices or undrained abscesses.

Clinical efficacy and safety

Adult

There are no data from comparator controlled clinical trials on the use of linezolid in the treatment of endocarditis, central nervous system infections and osteomyelitis.

Nosocomial pneumonia

Adult patients with clinically and radiologically documented nosocomial pneumonia participated in a randomised, multi-centre, double-blind clinical trial. Patients were treated for 7 to 21 days. One group (no. enrolled = 205) received linezolid injection 600 mg twice daily (bid), and another group (no. enrolled = 197) received vancomycin 1 g bid intravenously (IV). Both groups received concomitant aztreonam (1 to 2 g every 8 hours IV). Linezolid demonstrated efficacy equivalent to vancomycin in the treatment of patients with nosocomial pneumonia in all outcome measurements. The overall clinical cure rates in the ITT population was 53% in the linezolid group and 52% in the vancomycin group. These cure rates do not include patients with missing or indeterminate outcomes. The clinical cure rate for microbiologically evaluable patients is presented in Table 8.

Table 8. Clinical cure rates at the test-of-cure visit for microbiologically evaluable patients with nosocomial pneumonia (subjects with indeterminate or missing outcomes excluded)

Pathogen	Cured	
	Linezolid n/N (%)	Vancomycin n/N (%)
<i>Staphylococcus aureus</i>	25/41 (61)	14/23 (61)
<i>Streptococcus pneumoniae</i>	9/9 (100)	9/9 (100)

Community-acquired pneumonia

Adult patients with clinically and radiologically documented community-acquired pneumonia participated in two randomised, comparator-controlled, multi-centre trials.

One of these trials was an open-label study in which hospitalised patients received study medications administered IV followed by medications administered orally for a total of 7 to 14 days of treatment. One group of patients (no. enrolled = 389) received linezolid injection (600 mg bid) followed by linezolid tablets (600 mg bid), and another group (no. enrolled = 370) received ceftriaxone (1 g bid IV) followed by cefpodoxime proxetil tablets (200 mg bid orally).

The second study was an investigator-blinded trial in outpatients with community acquired pneumonia who were treated for 10 – 14 days. One group of patients received linezolid tablets 600 mg bid (no. enrolled = 278) and another group received cefpodoxime proxetil tablets 200 mg bid (no. enrolled = 270).

In these trials, linezolid demonstrated efficacy equivalent to ceftriaxone or cefpodoxime proxetil by all outcome measurements. The overall clinical cure rates in the ITT population in linezolid and comparator groups were 83% vs 76% and 82% vs 86% in respective studies. These cure rates do not include patients with missing or indeterminate outcomes. Table 9 shows the clinical cure rates for microbiologically evaluable patients in these studies.

Table 9. Clinical cure rates at the test-of-cure visit for microbiologically evaluable patients with community-acquired pneumonia (subjects with indeterminate or missing outcomes excluded)

Pathogen	Cured	
	Linezolid n/N (%)	ceftriaxone and cefpodoxime proxetil n/N (%)
<i>Streptococcus pneumoniae</i>	88/98 (90)	81/90 (90)
<i>Staphylococcus aureus</i>	29/32 (91)	22/29 (76)
<i>Haemophilus influenzae</i>	13/14 (93)*	23/26 (88)

Excluding patients who received concomitant treatment with aztreonam

Complicated skin and skin structure infections

Adult patients with clinically documented complicated skin and skin structure infections participated in a randomised, multi-centre, double-blind trial comparing study medications administered IV followed by medications given orally for a total of 10 to 21 days of treatment. One group of patients (no. enrolled = 403) received linezolid injection (600 mg bid) followed by linezolid tablets (600 mg bid); another group (no. enrolled = 423) received oxacillin 2 g every 6 hours (q6h) IV followed by dicloxacillin 500 mg q6h orally. Linezolid demonstrated equivalent efficacy to oxacillin and dicloxacillin against a variety of common pathogens by all outcome measurements. The overall clinical cure rates in the ITT population was 85% in the linezolid group and 77% in the oxacillin group, respectively. These cure rates do not include patients with missing or indeterminate outcomes. The clinical cure rates for microbiologically evaluable patients are presented in Table 10.

Table 10. Clinical cure rates at the test-of-cure visit for microbiologically evaluable patients with complicated skin and skin structure infections (subjects with indeterminate or missing outcomes excluded)

Pathogen	Cured	
	Linezolid n/N (%)	oxacillin and dicloxacillin n/N (%)
<i>Staphylococcus aureus</i>	83/93 (89)	88/103 (85)
<i>Staphylococcus epidermidis</i>	19/19 (100)	10/12 (83)
<i>Streptococcus pyogenes</i>	23/29 (79)	27/32 (84)
<i>Streptococcus agalactiae</i>	7/7 (100)	4/6 (67)

Methicillin-resistant Staphylococcus aureus (MRSA) infections

Adult patients with documented MRSA infections participated in a randomised, multi-centre, open-label trial. One group of patients (no. enrolled = 243) received linezolid injection 600 mg bid followed by linezolid tablets 600 mg bid. Another group of patients (no. enrolled = 225) received vancomycin 1 g bid IV. Both groups were treated for 7 to 28 days. Linezolid was comparable to vancomycin in the treatment of patients with MRSA pneumonia and skin and soft tissue infections. The overall clinical cure rates in the ITT population was 57% in the linezolid group and 55% in the comparator groups respectively. These cure rates do not include patients with missing or indeterminate outcomes. The clinical cure rates for microbiologically evaluable patients with MRSA are presented in Table 11.

Table 11. Clinical cure rates at the test-of-cure visit for microbiologically evaluable patients with MRSA infections (subjects with indeterminate or missing outcomes excluded)

Infection	Cured	
	Linezolid n/N (%)	vancomycin n/N (%)
MRSA pneumonia	9/12 (75)	12/16 (75)
MRSA skin and soft tissue infection	27/34 (79)	22/30 (73)

Vancomycin-resistant enterococcus (VRE) infections

Adult patients with documented or suspected VRE infections participated in a randomised, multi-centre, double-blind trial comparing a high dose (600 mg bid IV or orally) with a low dose of linezolid (200 mg bid IV or orally) for 7 to 28 days. 79 patients were enrolled in the high dose group and 66 enrolled in the low dose group.

Patients with VRE infections were also treated with linezolid 600 mg bid IV or orally in an open label, non-comparative, compassionate-use trial. These patients were treated for up to 21 days.

144 patients with VRE infections were enrolled in this trial.

The overall clinical cure rates in the ITT populations were 67% and 54% in the high dose compared to low dose group in the controlled study and 90% (evaluable population) in the compassionate use trial. These cure rates do not include patients with missing or indeterminate outcomes. The clinical cure rates for clinically evaluable patients are presented in Table 12 by source of infection.

Table 12. Clinical cure rates at the test-of-cure visit for clinically evaluable patients with suspected or proven VRE infections (subjects with indeterminate or missing outcomes excluded)

Source of Infection	Cured		
	Linezolid 600 mg bid n/N (%)		Linezolid 600 mg bid n/N (%)
	VRE Patients in Compassionate Use Study	Dose- Comparator Study	Dose- Comparator Study
Bacteraemia of unknown origin	10/12 (83)	6/9 (67)	2/2 (100)
Other	33/35 (94)	11/11 (100)	7/11 (64)
Peritonitis *	11/12 (92)	1/1 (100)	3/6 (50)
Intra-abdominal *	11/12 (92)	4/4 (100)	2/2 (100)
Catheter-related *	9/9 (100)	3/3 (100)	1/1 (100)
Not classified *†	2/2 (100)	3/3 (100)	1/2 (50)
Pneumonia	1/1 (100)	2/2 (100)	---
Skin and soft tissue	7/9 (78)	8/9 (89)	6/6 (100)
Urinary tract	1/1 (100)	12/13 (92)	13/19 (68)

* Data for these sources of infections are subset of 'Other'

† Includes sources of infection such as hepatic abscess, biliary sepsis, necrotic gall bladder, pericolic abscess, and pancreatitis

Paediatric patients

Uncomplicated Skin and Skin Structure Infections

Paediatric patients (5 through 17 years of age) with clinically documented uncomplicated skin and skin structure infections were enrolled in a randomised, multi-centre, double-blind trial comparing linezolid versus cefadroxil for a total of 10 to 21 days of treatment. Patients aged 5 through 11 years received linezolid for Oral Suspension 10 mg/kg every 12 hours or cefadroxil oral suspension 15 mg/kg every 12 hours. Patients aged 12 years or older received linezolid tablets 600 mg every 12 hours or cefadroxil capsules 500 mg every 12 hours. There were 248 linezolid-treated and 251 cefadroxil-treated patients enrolled in the study. Two hundred twentyfour (90.3%) linezolid-treated patients and 216 (86.1%) cefadroxil-treated patients were clinically evaluable. The cure rates in clinically evaluable patients were 91% in linezolid-treated patients and 90% in cefadroxil-treated patients. The cure rates by pathogen for microbiologically evaluable patients are presented in Table 13.

Table 13. Cure rates at the test-of-cure visit for microbiologically evaluable paediatric patients with uncomplicated skin and skin structure infections

Pathogen	Cured	
	Linezolid n/N (%)	Cefadroxil n/N (%)
<i>Staphylococcus aureus</i>	120/133(90)	111/125 (89)
<i>Methicillin-resistant S. aureus</i>	12/13 (92)	6/7 (86)
<i>Streptococcus agalactiae</i>	1/1 (100)	2/2 (100)
<i>Streptococcus pyogenes</i>	32/33 (97)	26/27 (96)

Infections Due to Resistant Gram-positive Organisms

A safety and efficacy study provided experience on the use of linezolid in paediatric patients for the treatment of hospital acquired pneumonia, complicated skin and skin structure infections, catheter-related bacteremia, bacteremia of unidentified source, and other infections due to resistant gram-positive bacterial pathogens, including methicillin-resistant *Staphylococcus aureus* (MRSA), methicillin-resistant *Staphylococcus epidermidis* (MRSE), penicillin-resistant *Streptococcus pneumoniae* (PRSP) and vancomycin-resistant *Enterococcus faecium* (VRE). Paediatric patients ranging in age from birth through 11 years with infections caused by the documented or suspected above organisms were enrolled in a randomised, open-label, comparator-controlled trial. One group of patients received linezolid IV injection 10 mg/kg every 8 hours followed by linezolid for Oral Suspension 10 mg/kg every 8 hours. A second group received vancomycin 10 to 15 mg/kg IV every 6 to 24 hours, depending on age and renal clearance. Patients who had confirmed VRE infections were placed in a third arm of the study and received linezolid 10 mg/kg every 8 hours IV and/or orally. All patients were treated for a total of 10 to 28 days and could receive concomitant gram-negative antibiotics if clinically indicated. There were 215 linezolid treated and 101 vancomycin-treated patients enrolled in the study. One hundred and fifty-one (70.2%) linezolid-treated patients and 73 (72.3%) vancomycin-treated patients were clinically evaluable. The cure rates in clinically evaluable patients were 89% in linezolid treated patients and 85% in vancomycin-treated patients. The cure rates for clinically and microbiologically evaluable patients are presented in Table 14.

Table 14. Cure rates at the test-of-cure visit for microbiologically evaluable paediatric patients with infections due to Gram-positive pathogens

Pathogen	Cured	
	Linezolid n/N (%)	Vancomycin n/N (%)
<i>Enterococcus faecalis</i>	7/10 (70)	3/4 (75)
<i>Enterococcus faecium</i>	5/5 (100)	0/0
<i>Staphylococcus aureus</i>	37/39 (95)	24/26 (92)
<i>Staphylococcus epidermidis</i>	23/29 (79)	11/13 (85)
All coagulase-negative <i>Staphylococci</i> *	32/38 (84)	12/15 (80)
<i>Streptococcus pneumoniae</i>	3/3 (100)	1/1 (100)
<i>Streptococcus pyogenes</i>	2/2 (100)	1/2 (50)

*Coagulase-negative staphylococci were considered pathogens in catheter-related bacteraemia and in neonates.

5.2 Pharmacokinetic properties

The mean pharmacokinetic parameters (standard deviation) of linezolid following single and multiple (*i.e.*, twice daily administration to steady-state) intravenous (IV) is given in Table 15.

Table 15. Mean (standard deviation) pharmacokinetic parameters of linezolid in adults derived from plasma concentrations

Healthy adult volunteers						
Linezolid dosage regimen	C _{max} mcg/mL (SD)	C _{min} mcg/mL (SD)	T _{max} hrs (SD)	AUC* mcg.h/mL (SD)	t _{1/2} hrs (SD)	CL mL/min (SD)
600 mg injection‡ single dose	12.90 (1.60)	-----	0.50 (0.10)	80.20 (33.30)	4.40 (2.40)	138 (39)
	bid dose	15.10 (2.52)	3.68 (2.36)	0.51 (0.03)	89.70 (31.00)	4.80 (1.70)

* AUC for single dose = AUC_{0-∞}
 * AUC for multiple doses = AUC_{0-τ}
 ‡ Data normalised from 625 mg dose
 C_{max} = Maximum plasma concentration

C_{min} = Minimum plasma concentration
 T_{max} = Time to C_{max}
 AUC = Area under concentration-time curve
 t_{1/2} = Elimination half life
 CL = Systemic clearance

As can be seen from the above table, average C_{min} values achieved in plasma using the 600 mg twice daily dosage regimen approximate to the highest MIC₉₀ (4 micrograms/mL) for the least susceptible microorganisms.

Absorption

Linezolid is rapidly and extensively absorbed following oral dosing. Maximum plasma concentrations are reached within 2 hours of dosing and the absolute bioavailability is approximately 100%. Absorption from the oral suspension is similar to that achieved with the film coated tablets. Steady-state conditions are achieved by the second or third day of dosing.

Linezolid may be administered without regard to the timing of meals. The time to reach the maximum concentration is delayed from 1.5 hours to 2.2 hours and C_{max} is decreased by about 17% when high fat food is given with linezolid. However, the total exposure measured as AUC_{0-∞} values is similar under both conditions.

Distribution

Linezolid is readily distributed to well perfused tissues. Its volume of distribution at steady-state averages at about 40-50 litres in healthy adults and approximates to total body water. Plasma protein binding is about 31% and is not concentration dependent.

Linezolid concentrations have been determined in various fluids from a limited number of subjects in volunteer studies following multiple dosing. The ratio of linezolid in saliva and sweat relative to plasma was 1.2:1.0 and 0.55:1.0 respectively. The ratio for epithelial lining fluid and alveolar cells of the lung was 4.5:1.0 and 0.15:1.0, when measured at steady-state C_{max} , respectively. In a small study of subjects with ventricular-peritoneal shunts and essentially noninflamed meninges, the ratio of linezolid in cerebrospinal fluid to plasma at C_{max} was 0.7:1.0 after multiple linezolid dosing.

Metabolism

Linezolid is not detectably metabolised by cytochrome P450 (CYP) isoenzymes *in vitro* and it does not inhibit the activities of clinically significant human CYP isoforms (1A2, 2C9, 2C19, 2D6, 2E1, 3A4). Linezolid does not significantly induce major cytochrome P450 isoenzymes in rats and does not induce human CYP2C9. Metabolic oxidation of the morpholine ring results primarily in two inactive open-ring carboxylic acid derivatives. The hydroxyethyl glycine metabolite (A) is the predominant human metabolite and is formed by a non-enzymatic process. The amino ethoxy acetic acid metabolite (B) is less abundant. Other minor, inactive metabolites have been characterised.

Elimination

In patients with normal renal function or mild to moderate renal insufficiency, linezolid is primarily excreted as metabolite A (40%), parent drug (30-35%) and metabolite B (10%) in the urine. Virtually no parent drug is found in the faeces whilst approximately 6% and 3% of each dose appears as metabolites A and B, respectively. The elimination half-life averages at about 57 hours.

Non-renal clearance accounts for approximately 65% of the total clearance of linezolid. A small degree of non-linearity in clearance is observed with increasing doses of linezolid. This appears to be due to lower renal and non-renal clearance at higher linezolid concentrations. However, the difference in clearance is small and is not reflected in the apparent elimination half-life.

Special Populations

Paediatric. The pharmacokinetics of linezolid following a single IV dose were investigated in healthy adolescent subjects, ranging in age from 12 through 17 years, and in paediatric patients, ranging in age from 1 week through 12 years. The pharmacokinetic parameters of linezolid are summarised in Table 16 for the paediatric populations studied and healthy adults subjects after administration of single IV dose.

The C_{max} and the volume of distribution (V_{ss}) of linezolid are similar regardless of age in paediatric patients. However, clearance of linezolid varies as a function of age. With the exclusion of pre-term neonates less than one week of age, clearance is most rapid in the youngest age groups ranging from >1 week old to 11 years, resulting in lower single-dose systemic exposure (AUC) and shorter half-life as compared with adults. As age of paediatric patients increases, the clearance of linezolid gradually decreases, and by adolescence mean clearance values approach those observed for the adult population. There is a wider inter-subject variability in linezolid clearance and systemic drug exposure (AUC) across all paediatric age groups as compared with adults.

Similar mean daily AUC values were observed in paediatric patients from birth to 11 years of age dosed every 8 hours relative to adolescents or adults dosed every 12 hours. Therefore, the dosage for paediatric patients up to 11 years of age should be 10 mg/kg every 8 hours. Paediatric patients 12 years and older should receive 600 mg every 12 hours (see **section 4.2**).

Recommendations for the dosage regimen for pre-term neonates less than 7 days of age (gestational age less than 34 weeks) are based on pharmacokinetic data from 9 pre-term neonates. Most of these pre-term neonates have lower systemic linezolid clearance values and larger AUC values than many full-term neonates and older infants. Therefore, these pre-term neonates should be initiated with a dosing regimen of 10 mg/kg every 12 hours. Consideration may be given to the use of a 10 mg/kg every 8 hours regimen in neonates with a sub-optimal clinical response. All neonatal patients should receive 10 mg/kg every 8 hours by 7 days of life (see **section 4.2**).

Table 16. Pharmacokinetic parameters of linezolid in paediatric and adult patients following a single intravenous infusion of 10 mg/kg or 600 mg linezolid (Mean:(% CV); Min., Max. values)

Age group	C _{max} µg/mL (SD)	V _{ss} L/kg (SD)	AUC* µg h/mL (SD)	t _{1/2} hrs (SD)	CL mL/min/kg (SD)
<u>Neonatal patients</u> <u>Pre - Term **</u> <u>< 1 week (N = 9)</u>	12.7 (30%) (9.6, 22.2)	0.81 (24%) (0.43, 1.05)	108 (47%) (41, 191)	5.6 (46%) (2.4, 9.8)	2.0 (52%) (0.9, 4.0)
<u>Full – Term***</u> <u>< 1 week (N = 10) †</u>	11.5 (24%) (8.0, 18.3)	0.78 (20%) (0.45, 0.96)	55 (47%) (19, 103)	3.0 (55%) (1.3, 6.1)	3.8 (55%) (1.5, 8.8)
Full – Term *** ≥1 week to ≤ 28 days (N = 10)	12.9 (28%) (7.7, 21.6)	0.66 (29%) (0.35, 1.06)	34 (21%) (23, 50)	1.5 (17%) (1.2, 1.9)	5.1 (22%) (3.3, 7.2)
<u>Infant Patients ≥</u> <u>28 days to < 3</u> <u>months (N = 12) †</u>	11.0 (27%) (7.2, 18.0)	0.79 (26%) (0.42, 1.08)	33 (26%) (17, 48)	1.8 (28%) (1.2, 2.8)	5.3 (34%) (3.5, 9.9)
Paediatric Patients 3 months through 11 years † (N = 59)	15.1 (30%) (6.8, 36.7)	0.69 (28%) (0.31, 1.50)	58 (54%) (19, 153)	2.9 (53%) (0.9, 8.0)	3.8 (53%) (1.0, 8.5)
Adolescents 12 through 17 years ‡ (N = 18)	16.7 (24%) (9.9, 28.9)	0.61 (15%) (0.44, 0.79)	95.0 (44%) (32, 178)	4.1 (46%) (1.3, 8.1)	2.1 (53%) (0.9, 5.2)
Adults§ (N = 29)	12.5 (21%) (8.2, 19.3)	0.65 (16%) (0.45, 0.84)	91 (33%) (53, 155)	4.9 (35%) (1.8, 8.3)	1.7 (34%) (0.9, 3.3%)

AUC = Area Under the Curve

* AUC = single dose AUC_{0-∞}

** In this data set “pre-term” is defined as <34 weeks gestational age (Note: only 1 patient enrolled was pre-term with a postnatal age between 1 week and 28 days)

*** In this data set “full-term” is defined as ≥ 34 weeks of gestational age.

† Dose of 10 mg/kg

‡ Dose of 10 mg/kg up to a maximum of 600 mg

§ Dose normalized to 600 mg

C_{max} = maximum plasma concentration

V_{ss} = volume of distribution

t_{1/2} = apparent elimination

half-life

CL = systemic clearance normalized for body weight

Geriatric: The pharmacokinetics of linezolid are not significantly altered in elderly patients aged 65 and over.

Gender: Some pharmacokinetic parameters of linezolid differ in female subjects. Females have a slightly lower volume of distribution than males and the mean clearance is reduced by approximately 20% when corrected for body weight. Plasma concentrations are somewhat higher in females and this can partly be attributed to body weight differences. However, because the mean half-life of linezolid is not significantly different in males and females, plasma concentrations in females are not expected to substantially rise above those known to be well tolerated and, therefore, dose adjustments are not required.

Renal insufficiency: No dose adjustment is necessary in patients with either mild, moderate or severe renal insufficiency as total clearance is independent of creatinine clearance. There is evidence that the two primary metabolites of linezolid accumulate in patients with severe renal insufficiency ($CL_{CR} < 30$ mL/min). The clinical significance of this has not been established as limited safety data are currently available. As approximately 30% of a linezolid dose is removed during 3 hours of haemodialysis (beginning 3 hours after administration), linezolid should be given after dialysis in patients receiving such treatment. The primary metabolites of linezolid are also removed by haemodialysis, but the concentrations of these metabolites are still considerably higher following dialysis than those observed in patients with normal renal function or mild to moderate renal insufficiency. No information is available on the effect of peritoneal dialysis on the pharmacokinetics of linezolid.

Hepatic insufficiency. The pharmacokinetics of linezolid are not altered in patients with mild to moderate hepatic insufficiency (*i.e.*, Child-Pugh class A or B). Dose adjustment in such patients is not required. The pharmacokinetics of linezolid in patients with severe hepatic insufficiency (*i.e.*, Child-Pugh class A or B) have not been evaluated. However, as linezolid is metabolised by a non-enzymatic process, impairment of hepatic function would not be expected to significantly alter its metabolism.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Glucose
Sodium citrate
Citric acid anhydrous
Hydrochloric acid/sodium hydroxide
Water for injection

6.2 Incompatibilities

Additives should not be introduced into linezolid injection. If linezolid is to be given concomitantly with other drugs, each drug should be given separately in accordance with its own directions for use. Similarly, if the same intravenous line is to be used for sequential infusion of several drugs, the line should be flushed prior to and following linezolid administration with a compatible infusion solution (5% glucose injection, 0.9% sodium chloride injection or compound sodium lactate injection [Hartmann's solution for injection]).

Linezolid solution for infusion is known to be physically incompatible with the following compounds: amphotericin B, chlorpromazine hydrochloride, diazepam, pentamidine isethionate, erythromycin lactobionate, phenytoin sodium and sulphamethoxazole/trimethoprim. Additionally, it is chemically incompatible with ceftriaxone sodium.

6.3 Shelf life

36 months.

6.4 Special for storage

Store below 25°C. Protect from freezing.
Do not use if the bag leaks as sterility may be impaired.

6.5 Nature and contents of container

Single use, ready-to-use, LDPE infusion bags.

Bags are packed in boxes of 1 or 10 300ml infusion bags.

6.6 Special precautions for disposal

Linezolid injection contains no preservative. The product is for single use in one patient only. Discard any residue. Do not use these bags in series connections. Do not reconnect partially used bags.

Any solutions which are discoloured, hazy or contain visible particulate matter should not be used.

7. MEDICINE SCHEDULE

Prescription Medicine

Further Information

Chemistry


Non-proprietary name: Linezolid

Chemical name: (S)-N-[[3-[3-Fluoro-4-(4-morpholinyl)phenyl]-2-oxo-5-oxazolidinyl]methyl]-acetamide

CAS Number: CAS-165800-03-3

Linezolid injection contain linezolid, which is a synthetic antibacterial agent of the oxazolidinone class. Linezolid is biologically active and is metabolised to form inactive derivatives. The aqueous solubility of linezolid is approximately 3 mg/mL, independent of pH between pH 3 to 9.

The empirical formula of linezolid is C₁₆H₂₀FN₃O₄. Its molecular weight is 337.35 and its chemical structure is represented below:



8. SPONSOR

Neo Health (NZ) Limited
Level 27, PWC Tower
188 Quay Street
Auckland 1010
New Zealand

Contact number: +64 3 4779 669

9. DATE OF FIRST APPROVAL

29 April 2019

10. DATE OF REVISION OF TEXT

29 April 2019

Summary of changes

Section changed	Summary of new information