

NEW ZEALAND DATA SHEET

1 PRODUCT NAME

Metaraminol

10 mg/mL solution for injection.

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 1 mL of solution contains 10mg of metaraminol (as tartrate).

Excipients of known effect:

Sodium chloride

Sodium metabisulfite

Each 1 mL of solution contains 98.3 micromol (2.26 mg) sodium.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Solution for injection

Glass ampoule containing a clear colourless solution with pH of 3.2 to 4.5 and osmolarity of 305 mOsm/L.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Prevention and treatment of the acute hypotensive state occurring with spinal anaesthesia; adjunctive treatment of hypotension due to haemorrhage, reactions to medications, surgical complications, and shock associated with brain damage due to tumour or trauma.

It may also be useful as an adjunct in the treatment of hypotension due to cardiogenic shock or septicaemia.

4.2 Dose and method of administration

Metaraminol may be administered subcutaneously, intra- muscularly, or intravenously (injection or infusion). Because the maximum effect is not immediately apparent, at least 10 minutes should elapse before increasing the dosage. As the effect tapers off when the vasopressor is discontinued the patient should be carefully observed so that therapy can be reinitiated promptly if the blood pressure falls too rapidly. Patients with coexistent shock and acidosis may show a poor response to vasopressors. Established methods of shock management, such as blood or fluid replacement when indicated, and other measures directed to the specific cause of the shock should also be used.

Intramuscular or Subcutaneous Injection (for prevention of hypotension)

The recommended dose is 2 to 10 mg (0.2 to 1 mL). As with other agents used subcutaneously, only the preferred sites of injection, as set forth in standard texts, should be used.

NEW ZEALAND DATA SHEET

Intravenous Infusion (for adjunctive treatment of hypotension)

The recommended dose is 15 to 100 mg (1.5 to 10 mL) in 500 mL of Sodium Chloride Injection or 5% Dextrose Injection, adjusting the rate of infusion to maintain the blood pressure at the desired level.

Higher concentrations of metaraminol (150 to 500 mg per 500 mL of infusion fluid) have been used.

If the patient needs additional saline or dextrose solution at a rate of flow that would provide an excessive dose of the vasopressor, the recommended volume (500 mL) of infusion fluid should be increased accordingly. Conversely, if a smaller volume of infusion fluid is desirable, the required dose of metaraminol may be added to less than 500 mL of diluent.

Direct Intravenous Injection

In severe shock, when time is of great importance, it may be desirable to administer metaraminol by direct intravenous injection. The suggested dose is 0.5 to 5 mg (0.05 to 0.5 mL), followed by an infusion of 15 to 100 mg (1.5 to 10 mL) in 500ml of diluent.

Direct intravenous injection of undiluted solution should be employed only in instances of grave emergency when prompt action is imperative to save life. Extreme care must be exercised to give the proper dosage.

4.3 Contraindications

Use of metaraminol tartrate with cyclopropane or halothane anaesthesia should be avoided, unless clinical circumstances demand such use. Hypersensitivity to any component of this product, including sulphites (refer to section 6.1).

4.4 Special warnings and precautions for use

Caution should be exercised to avoid excessive blood-pressure changes since response to treatment with metaraminol is very variable and the ensuing control of the blood pressure may prove difficult.

Rapidly induced hypertensive responses have been reported to cause acute pulmonary oedema, cardiac arrhythmias and arrest. Metaraminol should be used with caution in patients with cirrhosis; electrolyte levels should be adequately restored if a diuresis ensues. A fatal ventricular arrhythmia was reported in a patient with Laennec's cirrhosis while receiving metaraminol tartrate. In several instances ventricular extrasystoles that appeared during infusion of metaraminol promptly subsided when the rate of flow was reduced.

With the prolonged action of metaraminol, a cumulative effect is possible. An excessive vasopressor response may cause a prolonged elevation of blood pressure, even after discontinuation of therapy. Metaraminol should be used with caution in cases of heart disease, hypertension, thyroid disease or diabetes mellitus because of the vasoconstrictor action.

Sympathomimetic amines may provoke a relapse in patients with a history of malaria.

When vasopressor amines are used for long periods, the resulting vasoconstriction may prevent adequate expansion of circulating volume and may cause perpetuation of the shock state. There is evidence that plasma volume may be reduced in all types of shock, and that the measurement of

NEW ZEALAND DATA SHEET

central venous pressure is useful in assessing the adequacy of the circulating blood volume. Blood, or plasma-volume expanders, should therefore be employed when the principal reason for hypotension of shock is decreased circulating volume.

In choosing the site for injection, it is important to avoid those areas generally recognised as being unsuitable for the use of any pressor agent and to discontinue the infusion immediately if infiltration or thrombosis occurs. Although the urgent nature of the patient's condition may force the choice of an unsuitable injection site, the preferred areas of injection should be used when possible. The larger veins of the antecubital fossa or thigh are preferred to the veins in the ankle or dorsum of the hand, particularly in patients with peripheral vascular disease, diabetes mellitus, Buerger's disease or conditions with coexistent hypercoagulability.

The preservative sodium metabisulfite in Metaraminol 10mg/mL Solution for Injection may cause hypersensitivity. In particular it is associated with circulatory or respiratory collapse, and depression of the CNS in certain susceptible individuals, particularly in those with asthma.

Accidental spillage of Metaraminol 10mg/mL Solution for Injection on the skin can cause dermatitic reactions linked to the presence of the product's preservatives.

4.5 Interaction with other medicines and other forms of interaction

Metaraminol should be used with caution in patients receiving digitalis, since the combination of digitalis and sympathomimetic amines is capable of causing ectopic arrhythmic activity.

Monoamine oxidase inhibitors have been reported to potentiate the action of sympathomimetic amines. The pressor effect of metaraminol is decreased but not reversed by alpha-adrenergic blocking agents.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no well-controlled studies in pregnant women. Metaraminol should be used during pregnancy only if the potential benefit to the mother justifies the potential risk to the foetus.

Breast-feeding

It is not known whether metaraminol is secreted in human milk. Because many medicines are secreted in human milk, caution should be exercised if metaraminol is given to a breastfeeding mother.

Fertility

There are no fertility data available.

4.7 Effects on ability to drive and use machines

None stated.

NEW ZEALAND DATA SHEET

4.8 Undesirable effects

The frequency of adverse events with metaraminol has not been firmly established. Excessive therapeutic effect leading to hypertension, quickly reversible by reducing the rate of infusion, and headaches are very common.

Adverse reactions listed below are classified according to frequency and system organ class (SOC). The frequencies of adverse reactions are ranked according to the following convention: Common ($\geq 1/100$ to $< 1/10$); Uncommon ($\geq 1/1,000$ to $< 1/100$); Rare ($\geq 1/10,000$ to $< 1/1,000$); Very rare ($< 1/10,000$); Not known (cannot be estimated from the available data).

System Organ Class	Undesirable Effect
Nervous system disorders	Very common: Headache
Cardiac disorders	Not known: Palpitations; sinus tachycardia; bradycardia; ventricular tachycardia; other cardiac arrhythmias (especially in patients with myocardial infarction); fatal ventricular arrhythmia reported in Laennec's cirrhosis.
Vascular disorders	Very Common: Hypertension Not known: Peripheral ischaemia;
Skin and Subcutaneous tissue disorders:	Rare: Abscess formation; tissue necrosis; sloughing.
Gastrointestinal disorders	Not known: Nausea.

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare professionals are asked to report any suspected adverse reactions <https://nzphvc.otago.ac.nz/reporting/>

4.9 Overdose

Metaraminol acts rapidly. The major therapeutic effects are complete within an hour of parenteral administration. Overdosage may result in severe hypertension accompanied by headache, constricting sensation in the chest, nausea, vomiting, euphoria, diaphoresis, pulmonary oedema, tachycardia, bradycardia, sinus arrhythmia, atrial or ventricular arrhythmias, myocardial infarction, cardiac arrest or convulsions.

If the medicine has been ingested, induce emesis or perform gastric lavage. If metaraminol has been administered by subcutaneous or intramuscular injection, local ice packs may be applied to delay absorption. Intravenous infusion should be stopped immediately, but reinstated if hypotension occurs.

If needed, alpha-adrenergic blocking agents may also be useful for reducing hypertension and may have a beneficial effect on cardiac arrhythmia, if present. Parenteral diazepam may be given for convulsions.

NEW ZEALAND DATA SHEET

For advice on the management of overdose please contact the National Poisons Centre on 0800 POISON (0800 764766).

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Adrenergic and dopaminergic agent. ATC code: C01CA09.

Metaraminol is a sympathomimetic agent with direct and indirect effects on adrenergic receptors. It has both alpha and beta-adrenergic activity, the former being predominant.

Metaraminol increases the force of myocardial contractions as well as having a peripheral vasoconstrictor action. It increases both systolic and diastolic blood pressures.

The vasoconstrictor action of metaraminol is not affected by depletion of the tissue stores of noradrenaline. Metaraminol is highly effective in displacing and replacing noradrenaline from the stores in adrenergic neurones and competitively inhibits noradrenaline uptake. The metaraminol that is taken up by the adrenergic neurones then acts as a false transmitter.

The overall effects of metaraminol are similar to those of noradrenaline but it is much less potent and has a more prolonged action. It can cause pulmonary vasoconstriction, and pulmonary blood pressure is elevated when cardiac output is reduced.

5.2 Pharmacokinetic properties

The pressor effect of a single dose of metaraminol lasts from about 20 minutes up to one hour. Its onset is around one or two minutes after direct intravenous injection. The vasopressor effects taper off when therapy is stopped.

5.3 Preclinical safety data

No relevant information.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium chloride
Sodium metabisulfite
Water for injections

6.2 Incompatibilities

Metaraminol must not be mixed with the following medicinal products due to their additive incompatibilities:

Amphotericin B

NEW ZEALAND DATA SHEET

Dexamethasone
Prednisolone
Erythromycin
Hydrocortisone
Methicillin
Penicillin G
Thiopental

6.3 Shelf life

2 years.

After dilution:

Stable between 2-8°C for 24-48 hours in an intravenous infusion of Sodium Chloride 0.9% Solution or Glucose 5% Solution.

6.4 Special precautions for storage

Do not store above 25°C.

After dilution, chemical and physical in-use stability has been demonstrated for 48 hours when the diluted product is stored between 2 to 8°C.

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions are the responsibility of the user and would normally not be longer than 48 hours at 2 to 8°C unless opening has taken place in controlled and validated aseptic conditions.

6.5 Nature and contents of container

Glass ampoule containing 1mL of solution for injection or infusion.

Pack size of 10 ampoules in an outer carton.

6.6 Special precautions for disposal and other handling

Dilution instructions:

Metaraminol Injection may be diluted with Sodium Chloride 0.9% Injection or Dextrose 5% Injection. Refer to Section 4.2 for further details.

The product is compatible with:

- PVC infusion bags

Any unused product or waste material should be disposed of in accordance with local requirements.

NEW ZEALAND DATA SHEET

7 MEDICINE SCHEDULE

Prescription Medicine

8 SPONSOR

Max Health Ltd, P O Box 65 231, Mairangi Bay, Auckland 0754

Ph:(09) 815 2664.

9 DATE OF FIRST APPROVAL

20 June 2018

10 DATE OF REVISION OF THE TEXT

20 June 2018

SUMMARY TABLE OF CHANGES

Date of Revision	Section Changed	Summary of new information
20 June 2018	All	<ul style="list-style-type: none">• First approval