

New Zealand Datasheet

1 PRODUCT NAME

Omeprazole Infusion 40 mg

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Omeprazole 40 mg for intravenous infusion

3 PHARMACEUTICAL FORM

Each vial contains a white to off-white lyophilised powder consisting of omeprazole sodium 42.6 mg, equivalent to omeprazole 40 mg, which is intended to be reconstituted with 100 ml 5% glucose (solution not supplied with the dosage form). No other infusion solution should be used. The cap is aluminium with a white coloured plastic flip-off lid.

4.1 Therapeutic indications

Omeprazole Infusion 40 mg is indicated primarily for the treatment of Zollinger-Ellison syndrome, and may also be used for the treatment of gastric ulcer, duodenal ulcer and reflux oesophagitis.

4.2 Dose and method of administration

In patients with duodenal ulcer, gastric ulcer or reflux oesophagitis where oral medication is inappropriate, Omeprazole Infusion 40 mg once daily is recommended. In patients with Zollinger-Ellison syndrome the recommended initial dose of omeprazole given intravenously is 60 mg daily. Higher daily doses may be required and the dose should be adjusted individually. When doses exceed 60 mg daily, the dose should be divided and given twice daily.

Impaired Renal Function

Dose adjustment is not needed in patients with impaired renal function.

Impaired Hepatic Function

As plasma half-life of omeprazole is increased in patients with impaired hepatic function a daily dose of 10 - 20 mg may be sufficient.

Elderly

Dose adjustment is not needed in the elderly.

Children

There is limited experience with omeprazole IV in children.

Method of Administration

Omeprazole Infusion 40 mg infusion should be given as an intravenous infusion (over a period of 20-30 minutes or more). The contents of one vial must be dissolved in 100 ml of 5% dextrose for infusion or 100 ml of 0.9% saline for infusion. Do not use the solution if reconstitution is not complete. The solution should be used within 6 hours when dissolved in 5% dextrose and 12 hours when dissolved in saline. After reconstitution, start the infusion immediately. The constituted solution should not be mixed or co-administered in the same infusion set with any other drug.

4.3 Contraindications

Known hypersensitivity to omeprazole substituted benzimidazoles or any other constituent of the formulation.

4.4 Special warnings and precautions for use

In the presence of any alarm symptom (e.g. significant unintentional weight loss, recurrent vomiting, dysphagia, haematemesis or melena) and when gastric ulcer is suspected or present, the possibility of malignancy should be excluded as treatment may alleviate symptoms and delay diagnosis.

4.5 Interaction with other medicines and other forms of interaction

Effects of omeprazole on the pharmacokinetics of other medicines

Nelfinavir, atazanavir

Omeprazole has been reported to interact with some antiretroviral medicines. The clinical importance and the mechanisms behind these interactions are not always known. Increased gastric pH during omeprazole treatment may change the absorption of the antiretroviral medicine. Other possible interaction mechanisms are via CYP 2C19. For some antiretroviral medicines, such as atazanavir and nelfinavir, decreased serum levels have been reported when given together with omeprazole. Concomitant administration with omeprazole and medicines such as atazanavir and nelfinavir is therefore not recommended.

Citalopram / Escitalopram

Co-administration of omeprazole (20 mg) with citalopram (20 mg single dose) doubles the AUC of the S-isomer of citalopram, but the R-isomer of citalopram is not affected. A reduction in the dose of citalopram may be necessary based on clinical judgement. For patients taking omeprazole, the citalopram dose should not exceed the maximum dose of 20 mg/day.

Co-administration of omeprazole (30 mg) with escitalopram (20 mg single dose) increased the plasma levels (approximately 50%) and terminal half-life (31%) of escitalopram. A reduction in the dose of escitalopram may be necessary based on clinical judgement.

Digoxin

Concomitant treatment with omeprazole (20 mg daily) and digoxin in healthy subjects increased the bioavailability of digoxin by 10% (up to 30% in two out of ten subjects).

Clopidogrel

Results from studies in healthy subjects have shown a pharmacokinetic/pharmacodynamic interaction between clopidogrel (300 mg loading dose/75 mg daily maintenance dose) and omeprazole (80 mg p.o. daily, i.e. four times the recommended dose) resulting in decreased exposure to the active metabolite of clopidogrel by an average of 46%, and resulting in decreased maximum inhibition of (ADP induced) platelet aggregation by an average of 16%.

It is, however, uncertain to what extent this interaction is clinically important. One prospective, randomised (but incomplete) study (in over 3760 patients comparing placebo with omeprazole 20 mg in patients treated with clopidogrel and ASA) and non-randomised, post-hoc analyses of data from large, prospective, randomised clinical outcome studies (in over 47000 patients) did not show any evidence of an increased risk for adverse cardiovascular outcome when clopidogrel and PPIs, including omeprazole, were given concomitantly.

Results from a number of observational studies are inconsistent with regard to increased risk or no increased risk for CV thromboembolic events when clopidogrel is given together with a PPI.

When clopidogrel was given together with a fixed dose combination of esomeprazole 20 mg + ASA 81 mg compared to clopidogrel alone in a study in healthy subjects there was a decreased exposure by almost 40% of the active metabolite of clopidogrel. However, the maximum levels of inhibition of (ADP induced) platelet aggregation in these subjects were the same in the clopidogrel and the clopidogrel + the combined (esomeprazole + ASA) product groups, likely due to the concomitant administration of low dose ASA.

Other active substances

The absorption of erlotinib, ketoconazole and itraconazole is significantly reduced and thus clinical efficacy may be impaired. For posaconazole and erlotinib concomitant use should be avoided.

Active substances metabolised by CYP2C19

Omeprazole inhibits CYP2C19, the major omeprazole metabolising enzyme. Thus, the metabolism of concomitant medicines also metabolised by CYP2C19, such as diazepam, phenytoin, warfarin (R-warfarin) or other vitamin K antagonists and cilostazol, may be delayed.

Monitoring of patients receiving phenytoin is recommended and a reduction of the phenytoin dose may be necessary. However, concomitant treatment with omeprazole capsules 20 mg daily did not change the blood concentration of phenytoin in patients on continuous treatment with this medicine.

In patients receiving warfarin or other vitamin K antagonists, monitoring of INR is recommended and a reduction of the warfarin (or other vitamin K antagonist) dose may be necessary. Concomitant treatment with omeprazole capsules 20 mg daily did, however, not change coagulation time in patients on continuous treatment with warfarin.

Cilostazol

Omeprazole, given in doses of 40 mg to healthy subjects in a cross-over study, increased C_{max} and AUC for cilostazol by 18% and 26% respectively, and one of its active metabolites by 29% and 69% respectively.

Other

Omeprazole is partly metabolised also by CYP3A4, but omeprazole does not inhibit this enzyme. Thus, omeprazole does not affect the metabolism of medicines metabolised by CYP3A4, such as cyclosporin, lidocaine, quinidine, oestradiol, erythromycin, and budesonide. However, omeprazole has been shown to induce CYP1A2-mediated metabolism of clozapine. Close monitoring of plasma clozapine levels is recommended.

Results from a range of interaction studies with omeprazole versus other medicines demonstrate that omeprazole, 20-40 mg daily, has no significant influence on any other CYP enzymes relevant for medicine metabolism, as shown by the lack of metabolic interaction with substrates for CYP1A2 (such as caffeine, theophylline), CYP2C9 (such as S-warfarin, piroxicam, diclofenac, naproxen), CYP2D6 (such as metoprolol, propranolol), CYP2E1 (such as ethanol). However, omeprazole has been shown to induce CYP1A2-mediated metabolism of clozapine. Close monitoring of plasma clozapine levels is recommended.

Unknown mechanism:

Tacrolimus

Concomitant administration of omeprazole has been reported to increase the serum levels of tacrolimus.

Methotrexate

When given together with proton pump inhibitors, methotrexate levels have been reported to increase in some patients. In high-dose methotrexate administration a temporary withdrawal of omeprazole may need to be considered.

Saquinavir

For other antiretroviral medicines, such as saquinavir, elevated serum levels have been reported. There are also some antiretroviral medicines of which unchanged serum levels have been reported when given with omeprazole.

Effects of other medicines on the pharmacokinetics of omeprazole**Inhibitors CYP2C19 and/or CYP3A4**

Since omeprazole is metabolised by CYP2C19 and CYP3A4, medicines known to inhibit CYP 2C19 or CYP 3A4 or both (such as clarithromycin and voriconazole) may lead to increased omeprazole serum levels by decreasing the rate of omeprazole's metabolism.

Concomitant voriconazole treatment resulted in more than doubling of the omeprazole exposure. Since high doses of omeprazole have been well-tolerated, adjustment of the omeprazole dose is not required during temporary concomitant use.

Inducers of CYP2C19 and/or CYP3A4

Medicines known to induce CYP 2C19 or CYP 3A4 or both (such as rifampicin) may lead to decreased omeprazole serum levels by increasing omeprazole's rate of metabolism.

4.6 Fertility, pregnancy and lactation

Pregnancy

Results from three prospective epidemiological studies indicate no adverse effects of omeprazole on pregnancy or on the health of the foetus/newborn child. Omeprazole can be used during pregnancy.

Breast feeding

Omeprazole is excreted in breast milk but is not likely to influence the child when therapeutic doses are used.

4.7 Effects on ability to drive and use machines

Omeprazole is not likely to affect the ability to drive or use machines.

4.8 Undesirable effects

The following adverse reactions have been identified or suspected in the clinical trials programme for omeprazole and post-marketing. None was found to be dose-related. The reactions are classified according to frequency (common >1/100, <1/10; uncommon >1/1000, <1/100; rare >1/10000, <1/1000; very rare <1/10000).

Blood and lymphatic system disorders

Rare: Leukopenia, thrombocytopenia, agranulocytosis, pancytopenia

Immune system disorders

Rare: Hypersensitivity reactions e.g. fever, angioedema and anaphylactic reaction/shock

Metabolism and nutrition disorders

Rare: Hyponatraemia

Very rare: Hypomagnesaemia, severe hypomagnesaemia may result in hypocalcaemia. Hypomagnesaemia may also result in hypokalaemia

Psychiatric disorders

Uncommon: Insomnia

Rare: Agitation, aggression, confusion, depression, hallucinations

Nervous system disorders

Common: Headache

Uncommon: Dizziness, paraesthesia, somnolence

Rare: Taste disturbance

Eye disorders

Rare: Blurred vision

Ear and labyrinth disorders

Uncommon: Vertigo

Respiratory, thoracic and mediastinal disorders

Rare: Bronchospasm

Gastrointestinal disorders

Common: Abdominal pain, constipation, diarrhoea, flatulence, nausea/vomiting

Rare: Dry mouth, stomatitis, gastrointestinal candidiasis, microscopic colitis

Frequency not known: Withdrawal of long-term PPI therapy can lead to aggravation of acid-related symptoms and may result in rebound acid hypersecretion.

Hepatobiliary disorders

Uncommon: Increased liver enzymes

Rare: Hepatitis with or without jaundice, hepatic failure, encephalopathy in patients with pre-existing liver disease

Skin and subcutaneous tissue disorders

Uncommon: Dermatitis, pruritus, rash, urticaria

Rare: Alopecia, photosensitivity, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis (TEN)

Musculoskeletal, connective tissue and bone disorders

Rare: Arthralgia, myalgia, muscular weakness

Renal and urinary disorders

Rare: Interstitial nephritis

Reproductive system and breast disorders

Rare: Gynaecomastia

General disorders and administration site conditions

Uncommon: Malaise

Rare: Increased sweating, peripheral oedema

Irreversible visual impairment has been reported in isolated cases of critically ill patients who have received omeprazole intravenous injection, especially at high doses, but no causal relationship has been established.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare professionals are asked to report any suspected adverse reactions

<https://nzphvc.otago.ac.nz/reporting/>.

4.9 Overdose

Omeprazole IV doses of up to 270 mg on a single day and up to 650 mg over a three-day period have been given in clinical trials without any dose-related adverse reactions.

For advice on the management of overdose please contact the National Poisons Centre on 0800 POISON (0800 764766).

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Proton pump inhibitors, ATC code: A02BC01

Omeprazole is a racemic mixture of two active enantiomers, reduces gastric acid secretion through a highly targeted mechanism of action. It is a specific inhibitor of the acid pump in the parietal cell. It is rapid acting and provides control through reversible inhibition of gastric acid secretion with once daily dosing.

Site and mechanism of action

Omeprazole is a weak base and is concentrated and converted to the active form in the highly acidic environment of the intracellular canaliculi within the parietal cell, where it inhibits the enzyme H⁺, K⁺-ATPase, the acid pump. This effect on the final step of the gastric acid formation process is dose-dependent and provides for highly effective inhibition of both basal acid secretion and stimulated acid secretion, irrespective of the stimulus.

All pharmacodynamic effects observed can be explained by the effect of omeprazole on acid secretion.

Effect on gastric acid secretion

Intravenous omeprazole produces a dose dependent inhibition of gastric acid secretion in humans. In order to immediately achieve a similar reduction of intragastric acidity as after repeated dosing with 20 mg orally, a first dose of 40 mg intravenously is recommended. This results in an immediate decrease in intragastric acidity and a mean decrease over 24 hours of approximately 90%.

The inhibition of acid secretion is related to the area under the plasma concentration-time curve (AUC) of omeprazole and not to the actual plasma concentration at a given time.

No tachyphylaxis has been observed during treatment with omeprazole.

Effect on *Helicobacter pylori*

See Omeprazole capsules datasheet for information.

Other effects related to acid inhibition

During long-term treatment gastric glandular cysts have been reported in a somewhat increased frequency. These changes are a physiological consequence of pronounced inhibition of acid secretion, are benign and appear to be reversible

Decreased gastric acidity due to any means including proton pump inhibitors, increases gastric counts of bacteria normally present in the gastrointestinal tract.

Treatment with acid-reducing drugs may lead to slightly increased risk of gastrointestinal infections such as *Salmonella* and *Campylobacter* and possibly also *Clostridium difficile* in hospitalised patients.

5.2 Pharmacokinetic properties

Distribution

The apparent volume of distribution in healthy subjects is approximately 0.3 L/kg and a similar value is also seen in patients with renal insufficiency. In elderly patients, and in patients with hepatic insufficiency, the volume of distribution is slightly decreased.

The plasma protein binding of omeprazole is about 95%.

Metabolism and excretion

Omeprazole is completely metabolised by the cytochrome P450 system (CYP). The major part of its metabolism is dependent on the polymorphically expressed, specific isoform CYP2C19, responsible for the formation of hydroxyomeprazole, the major metabolite in plasma. The remaining part is dependent on another specific isoform, CYP3A4, responsible for the formation of omeprazole sulphone. As a consequence of high affinity of omeprazole to CYP2C19, there is a potential for competitive inhibition and metabolic drug-drug interactions with other substrates for CYP2C19. However, due to low affinity to CYP3A4, omeprazole has no potential to inhibit the metabolism of other CYP3A4 substrates.

The parameters below reflect mainly the pharmacokinetics in individuals with a functional CYP2C19 enzyme, extensive metabolisers.

Total plasma clearance is about 30-40 L/h after a single dose. The plasma elimination half-life of omeprazole is usually shorter than one hour both after single and repeated oral once daily dosing. The AUC of omeprazole increases with repeated administration. This increase is dose-dependent and results in a non-linear dose-AUC relationship after repeated administration. This time- and dose-dependency is due to a decrease of first pass metabolism and systemic clearance probably caused by an inhibition of the CYP2C19 enzyme by omeprazole and/or its metabolites (eg, the sulphone). Omeprazole is completely eliminated from plasma between doses with no tendency for accumulation during once daily administration.

No metabolite has been found to have any effect on gastric acid secretion. Almost 80% of an intravenously given dose is excreted as metabolites in the urine, and the remainder is found in the faeces, primarily originating from bile secretion.

Poor metabolisers: Approximately 3% of the Caucasian population and 15-20% of Asian populations lack a functional CYP2C19 enzyme and are called poor metabolisers. In such individuals the metabolism of omeprazole is probably mainly catalysed by CYP3A4. After repeated once-daily administration of 20 mg omeprazole, the mean AUC was 5 to 10 times higher in poor metabolisers than in subjects having a functional CYP2C19 enzyme (extensive metabolisers). Mean peak plasma concentrations were also higher, by 3 to 5 times. These findings have no implications for the dosage of Omeprazole.

Special patient populations

Impaired hepatic function: The metabolism of omeprazole in patients with liver dysfunction is impaired, resulting in an increased AUC. Omeprazole has not shown any tendency to accumulate with once daily dosing.

Impaired renal function: The pharmacokinetics of omeprazole, including systemic bioavailability and elimination rate, are unchanged in patients with reduced renal function.

Elderly: The metabolism rate of omeprazole is somewhat reduced in elderly subjects (75-79 years of age).

Children: There is limited experience with Omeprazole for intravenous use in children.

5.3 Preclinical safety data

Gastric ECL-cell hyperplasia and carcinoids, have been observed in life-long studies in rats treated with omeprazole. These changes are the result of sustained hypergastrinaemia secondary to acid inhibition. Similar findings have been made after treatment with H₂-receptor antagonists, proton pump inhibitors and after partial fundectomy. Thus, these changes are not from a direct effect of any individual drug.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Each vial contains disodium edetate 1.0 mg and sodium hydroxide q.s. for pH adjustment.

6.2 Incompatibilities

None known when instructions in section 4.2 are followed.

6.3 Shelf life

Unopened packages: 2 years at a temperature not exceeding 25°C.

6.4 Special precautions for storage

Omeprazole infusion 40 mg must be dissolved in 100 ml 5% dextrose for infusion or 100 ml 0.9% saline for infusion. Chemical and physical in-use stability has been demonstrated for 6 hours after reconstitution with 5% dextrose and 12 hours after reconstitution with saline.

From a microbiological point of view, the product should be used immediately, unless reconstitution has taken place in controlled and validated aseptic conditions. The solution can be handled at normal indoor light without special precaution.

6.5 Nature and contents of container

Omeprazole Infusion 40 mg for Infusion: 1 or 5 vials of lyophilised powder.

6.6 Special precautions for disposal

Any unused product or waste material should be disposed of in accordance with local requirements.

7 MEDICINE SCHEDULE

Prescription Medicine.

8 SPONSOR

Dr Reddy's New Zealand Ltd
82 Totara Crescent
Lower Hutt 5011
WELLINGTON

Tel: 0800 362 733

9 DATE OF FIRST APPROVAL

28 August 2008

10 DATE OF REVISION OF THE TEXT

14 February 2019

SUMMARY TABLE OF CHANGES

Section changed	Summary of new information
4.8	Reporting of adverse events added
4.8	Additional adverse effect for rebound acid hypersecretion
4.9	Poison centre contact details added