

## RAZENE®

---

### 1. Product Name

---

Razene, 10 mg, film coated tablet.

---

### 2. Qualitative and Quantitative Composition

---

Each film coated tablet contains 10 mg of cetirizine hydrochloride.

Razene tablets contain lactose.

For the full list of excipients, see section 6.1.

---

### 3. Pharmaceutical Form

---

White, capsule shaped, film coated tablets marked CZ break line 10 on one side and G on the reverse.

The tablet can be divided into equal doses.

---

### 4. Clinical Particulars

---

#### 4.1 *Therapeutic indications*

Cetirizine is indicated for the relief of symptoms associated with seasonal allergic rhinitis (hay fever) and perennial allergic rhinitis. Symptoms treated effectively include sneezing, rhinorrhea, post-nasal discharge, nasal pruritus, ocular pruritus and tearing.

Cetirizine is also indicated for the treatment of:

- allergic conjunctivitis;
- insect bites and
- the uncomplicated skin manifestations of chronic idiopathic urticaria. It significantly reduces the occurrence, severity and duration of hives and markedly reduces pruritus.

#### 4.2 *Dose and method of administration*

##### **Dose**

##### **Adults**

The recommended initial dose of cetirizine is 5 to 10 mg depending on symptom severity, given as a single daily dose with a glass of liquid. The time of administration may be varied to suit individual patient needs. If sufficient response is not obtained, the dose may be increased to the maximum recommended daily dose of 20 mg.

##### **Special populations**

##### **Elderly**

Data does not suggest that the dose needs to be reduced in elderly subjects provided that the renal function is normal.

---

## Renal impairment

The dosing intervals must be individualised according to renal function. Refer to the following table and adjust the dose as indicated. To use this dosing table, an estimate of the patient's creatinine clearance (CL<sub>cr</sub>) in mL/min is needed. The CL<sub>cr</sub> (mL/min) may be estimated from serum creatinine (mg/dl) determination using the following formula:

$$CL_{cr} = \frac{[140 - \text{age}(\text{years})] \times \text{weight}(\text{kg})}{72 \times \text{serum creatinine}(\text{mg} / \text{dl})} \quad (\times 0.85 \text{ for women})$$

### Dosing adjustments for adult patients with impaired renal function

Group	Creatinine clearance (mL/min)	Dosage and frequency
Normal	≥ 80	10 mg once daily
Mild	50-79	10 mg once daily
Moderate	30-49	5 mg once daily
Severe	<30	5 mg once every 2 days
End-stage renal disease – patients undergoing dialysis	<10	Contraindicated

In paediatric patients suffering from renal impairment, the dose will have to be adjusted on an individual basis taking into account the renal clearance of the patient and body weight.

## Hepatic impairment

No dose adjustment is needed in patients with solely hepatic impairment.

### Children aged 6-12 years

10 mg once daily.

A 5 mg starting dose (half a tablet) may be proposed if this leads to satisfactory control of the symptoms.

### Children aged 2-6 years

The recommended dose of cetirizine is 5 mg (half a tablet) once daily.

## 4.3 Contraindications

Cetirizine is contraindicated in:

- patients with a history of hypersensitivity to any of the constituents of the formulation, to hydroxyzine or to any piperazine derivatives
- patients with end stage renal impairment at less than 10 mL/min creatine clearance.

## 4.4 Special warnings and precautions for use

### Alcohol

At therapeutic doses, no clinically significant interactions have been demonstrated with alcohol (for a blood alcohol level of 0.5 g/L). Nevertheless, precaution is recommended if alcohol is taken concomitantly.

### **Increased risk of urinary retention**

Caution should be taken in patients with predisposition factors of urinary retention (e.g. spinal cord lesion, prostatic hyperplasia) as cetirizine may increase the risk of urinary retention (see section 4.8).

### **Patients at risk of convulsions**

Caution in epileptic patients and patients at risk of convulsions is recommended.

### **Skin reactions**

Pruritus and/or urticaria may occur when cetirizine is stopped, even if those symptoms were not present before treatment initiation (see section 4.8). In some cases, the symptoms may be intense and may require treatment to be restarted. The symptoms should resolve when the treatment is restarted.

### **Use in children**

Cetirizine is not recommended for use in children under 2 years of age.

### **Rebound pruritus**

Pruritus and/or urticarial may occur when cetirizine is stopped, even if those symptoms were not present before treatment initiation. In some cases, the symptoms may be intense and may require treatment to be restarted. The symptoms should resolve when the treatment is restarted.

## **4.5 Interaction with other medicines and other forms of interaction**

### **Lack of interaction**

- Pharmacokinetic interaction studies were conducted with cetirizine and pseudoephedrine, antipyrine, cimetidine, ketoconazole, erythromycin, and azithromycin; no pharmacokinetic interactions were observed.
- In a multiple dose study of theophylline (400 mg once a day) and cetirizine, there was a small (16%) decrease in clearance of cetirizine, while the disposition of theophylline was not altered by concomitant cetirizine administration.
- Studies with cetirizine and cimetidine, glipizide, diazepam, and pseudoephedrine have revealed no evidence of adverse pharmacodynamic interactions.
- Studies with cetirizine and azithromycin, erythromycin, ketoconazole, theophylline, antipyrine, and pseudoephedrine have revealed no evidence of adverse clinical interactions.
- In particular, concomitant administration of cetirizine with macrolides or ketoconazole has never resulted in clinically relevant ECG changes.

### **Ritonavir**

In a multiple dose study of ritonavir (600 mg twice daily) and cetirizine (10 mg daily), the extent of exposure to cetirizine was increased by about 40% while the disposition of ritonavir was slightly altered (-11%) further to concomitant cetirizine administration.

### **Food**

The extent of absorption of cetirizine is not reduced with food, although the rate of absorption is decreased by 1 hour.

### **Allergy skin test**

Allergy skin tests are inhibited by antihistamines and a wash-out period of 3 days is recommended before performing them.

## **4.6 Fertility, pregnancy and lactation**

### **Fertility**

Limited data is available on human fertility but no safety concern has been identified. Animal data shows no safety concern for human reproduction.

### **Pregnancy**

Category B2. Caution should be exercised in pregnant women.

For cetirizine, very rare clinical data on exposed pregnancies are available. Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal development, parturition or postnatal development.

### **Breast-feeding**

Caution should be exercised in lactating women.

Cetirizine is excreted in human milk at concentrations representing 0.25 to 0.90 those measured in plasma, depending on sampling time after administration.

## **4.7 Effects on ability to drive and use machines**

Objective measurements of driving ability, sleep latency and assembly line performance have not demonstrated any clinically relevant effects at the recommended dose of 10 mg.

Patients intending to drive, engage in potentially hazardous activities or operate machinery should not exceed the recommended dose and should take their response to the medicinal product into account.

In sensitive patients, concurrent use with alcohol or other CNS depressants may cause additional reductions in alertness and impairment of performance.

## **4.8 Undesirable effects**

### **Clinical trial data**

Clinical studies have shown that cetirizine at the recommended dosage has minor adverse effects on the CNS, including somnolence, fatigue, dizziness and headache.

In some cases, paradoxical CNS stimulation has been reported.

Although cetirizine is a selective antagonist of peripheral H<sub>1</sub>-receptors and is relatively free of anticholinergic activity, isolated cases of micturition difficulty, eye accommodation disorders and dry mouth have been reported.

Instances of abnormal hepatic function with elevated hepatic enzymes accompanied by elevated bilirubin have been reported. Mostly this resolves upon discontinuation of the drug.

Double blind controlled clinical trials comparing cetirizine to placebo or other antihistamines at the recommended dosage (10 mg daily for cetirizine), of which quantified safety data are available, included more than 3200 subjects exposed to cetirizine.

From this pooling, the following adverse reactions were reported for cetirizine 10 mg in the placebo-controlled trials at rates of 1.0% or greater:

<b>Adverse reactions: (WHO-ART)</b>	<b>Cetirizine (n=3260)</b>	<b>Placebo (n=3061)</b>
<i>Body as a whole- general disorders:</i> Fatigue	1.63%	0.95%
<i>Central and peripheral nervous systems disorders:</i> Dizziness Headache	1.10% 7.42%	0.98% 8.07%
<i>Gastro-intestinal system disorders:</i> Abdominal pain Dry mouth Nausea	0.98% 2.09% 1.07%	1.08% 0.82% 1.14%
<i>Psychiatric disorders:</i> Somnolence	9.63%	5.00%
<i>Respiratory disorders:</i> Pharyngitis	1.29%	1.34%

Although statistically more common than under placebo, somnolence was mild to moderate in the majority of cases.

Objective tests as demonstrated by other studies have demonstrated that usual daily activities are unaffected at the recommended daily dose in healthy young volunteers.

Adverse reactions at rates 1% or greater in children aged from 6 months to 12 years, included in placebo-controlled clinical trials are:

<b>Adverse reactions (WHO-ART)</b>	<b>Cetirizine (n=1656)</b>	<b>Placebo (n=1294)</b>
<i>Gastro-intestinal system disorders:</i> Diarrhoea	1.0%	0.6%
<i>Psychiatric disorders:</i> Somnolence	1.8%	1.4%
<i>Respiratory system disorders:</i> Rhinitis	1.4%	1.1%
<i>Body as a whole- general disorders:</i> Fatigue	1.0%	0.3%

### Post marketing experience

Adverse reactions are ranked under headings of frequency using the following convention:

Very common  $\geq 1/10$

Common  $\geq 1/100$  to  $<1/10$

Uncommon  $\geq 1/1000$  to  $<1/100$

Rare  $\geq 1/10000$  to  $<1/1000$

Very Rare  $<1/10000$

Not known (cannot be estimated from the available data).

<b>Body system</b>	<b>Undesirable effect</b>	<b>Frequency</b>
Blood and lymphatic system disorders	Thrombocytopenia	Very rare
Immune system disorders	Hypersensitivity	Rare
	Anaphylactic shock	Very rare

Metabolism and nutrition disorders	Increased appetite	Not known
Psychiatric disorders	Agitation	Uncommon
	Aggression, confusion, depression, hallucination, insomnia	Rare
	Tic	Very rare
	Suicidal ideation, nightmare	Not known
Nervous system disorder	Paraesthesia	Uncommon
	Convulsions	Rare
	Dysgeusia, dyskinesia, dystonia, syncope, tremor	Very rare
	Amnesia, memory impairment	Not known
Eye disorder	Accommodation disorder, blurred vision, oculogyration	Very rare
Ear and labyrinth disorders	Vertigo	Not known
Cardiac disorder	Tachycardia	Rare
Gastrointestinal disorders	Diarrhoea	Uncommon
Hepatobiliary disorder	Hepatic function abnormal, transaminases increased, blood bilirubin increased, blood alkaline phosphatase increased, Gamma-glutamyltransferase increased	Rare
	Hepatitis	Not known
Skin and subcutaneous tissue disorders	Pruritis, rash	Uncommon
	Urticaria	Rare
	Angioedema, drug eruption	Very rare
	Acute generalized exanthematous pustulosis (AGEP)	Not known
Musculoskeletal and connective tissue disorders	Arthralgia	Not known
Renal and urinary disorders	Dysuria, enuresis	Very rare
	Urinary retention (See Section 4.4)	Not known

General disorders and administration site conditions	Asthenia, malaise Oedema	Uncommon Rare
Investigations	Weight gain	Rare

### **Skin reactions occurring after discontinuation of cetirizine**

After discontinuation of cetirizine, pruritus (intense itching) and/or urticaria have been reported (see Section 4.4).

### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare professionals are asked to report any suspected adverse reactions <https://nzphvc.otago.ac.nz/reporting/>.

## **4.9 Overdose**

### **Symptoms and signs**

Symptoms observed after an overdose of cetirizine are mainly associated with CNS effects or with effects that could suggest an anticholinergic effect.

Adverse events reported after an intake of at least five times the recommended daily dose are: confusion, diarrhoea, dizziness, fatigue, headache, malaise, mydriasis, pruritus, restlessness, sedation, somnolence, stupor, tachycardia, tremor, and urinary retention.

### **Treatment**

There is no known specific antidote to cetirizine.

Should overdose occur, symptomatic or supportive treatment is recommended.

Cetirizine is not effectively removed by dialysis.

For further advice on management of overdose please contact the National Poisons Information Centre (0800 POISON or 0800 764 766).

## **5. Pharmacological Properties**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: antihistamines for systemic use, piperazine derivatives.

ATC code: R06AE07

### **Mechanism of action**

Cetirizine, a human metabolite of hydroxyzine, is a potent and selective antagonist of peripheral H<sub>1</sub>-receptors. *In vitro* receptor binding studies have shown no measurable affinity for receptors other than H<sub>1</sub>-receptors.

*Ex vivo* experiments in mice have shown that systemically administered cetirizine does not significantly occupy the cerebral H<sub>1</sub>-receptors.

In addition to its anti-H<sub>1</sub> effect, cetirizine was shown to display anti-allergic activities: at a dose of 10 mg once or twice daily, it inhibits the late phase recruitment of inflammatory cells, notably eosinophils, in the skin and conjunctiva of atopic subjects submitted to antigen challenge, and the dose of 30 mg/day inhibits the influx of eosinophils in the bronchoalveolar lavage fluid during a late-

phase bronchial constriction induced by allergen inhalation in asthmatic subjects. Moreover, cetirizine inhibits the late-phase inflammatory reaction induced in chronic urticarial patients by intradermal administration of kallikrein. It also down-regulates the expression of adhesion molecules, such as ICAM-1 and VCAM-1, which are markers of allergic inflammation.

### **Onset and duration of action**

Studies in healthy volunteers show that cetirizine, at doses of 5 and 10 mg strongly inhibits the wheal and flare reactions induced by very high concentrations of histamine into the skin. The onset of activity after a single 10 mg dose occurs within 20 minutes in 50% of the subjects and within one hour in 95%. This activity persists for at least 24 hours after a single administration. In a 35-day study in children aged 5 to 12, no tolerance to the antihistaminic effect (suppression of wheal and flare) of cetirizine was found. When a treatment with cetirizine is stopped after repeated administration, the skin recovers its normal reactivity to histamine within 3 days.

## **5.2 Pharmacokinetic properties**

Cetirizine exhibits linear kinetics over the range of 5 to 60 mg. The terminal half-life is approximately 10 hours and the apparent volume of distribution is 0.50 L/kg.

No accumulation is observed for cetirizine following daily doses of 10 mg for 10 days. The steady state maximum plasma concentration is approximately 300 ng/mL and is achieved within  $1.0 \pm 0.5$  h.

Plasma protein binding is  $93 \pm 0.3\%$ .

Cetirizine does not modify the protein binding of warfarin.

Cetirizine does not undergo extensive first pass metabolism. About two third of the dose are excreted unchanged in urine. The distribution of pharmacokinetic parameters as peak level and area under curve, is unimodal in human volunteers and no differences were observed in the kinetics of cetirizine between white and black adult males. The extent of absorption of cetirizine is not reduced with food, although the rate of absorption is decreased. The extent of bioavailability is similar when cetirizine is given as solutions, capsules or tablets.

### **Special populations**

#### ***Children***

The half-life of cetirizine was about 6 hours in children 6-12 years and 5 hours in children 2-6 years. In infants and toddlers aged 6 to 24 months, it is reduced to 3.1 hours.

#### ***Elderly***

Following a single 10 mg oral dose, half-life increased by about 50% and clearance decreased by 40% in 16 elderly subjects compared to the normal subjects. The decrease in cetirizine clearance in these elderly volunteers appeared to be related to their decreased renal function.

#### ***Renal impairment***

The pharmacokinetics of the drug was similar in patients with mild impairment (creatinine clearance higher than 40 mL/min) and normal volunteers.

Moderately renally impaired patients had a 3-fold increase in half-life and 70% decrease in clearance compared to normal volunteers.

Patients on haemodialysis (creatinine clearance less than 7 mL/min) given a single oral 10 mg dose of cetirizine had a 3-fold increase in half-life and a 70% decrease in clearance compared to normal. Cetirizine was poorly cleared by haemodialysis. Dosing adjustment is necessary in patients with moderate or severe renal impairment.



### ***Hepatic impairment***

Patients with chronic liver diseases (hepatocellular, cholestatic, and biliary cirrhosis) given 10 or 20 mg of cetirizine as a single dose had a 50% increase in half-life along with a 40% decrease in clearance compared to healthy subjects.

Dosing adjustment is only necessary in hepatically impaired patients if concomitant renal impairment is present.

### ***5.3 Preclinical safety data***

Non-clinical data reveals no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential and toxicity to reproduction.

---

## **6. Pharmaceutical Particulars**

---

### ***6.1 List of excipients***

Each Razene tablet also contains:

Tablet core:

- lactose
- pregelatinised maize starch
- povidone
- magnesium stearate.

Film coat:

- hydroxypropylmethyl cellulose
- titanium dioxide
- macrogol.

Razene tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, (the Lapp lactase deficiency or glucose-galactose malabsorption) should not take this medicine.

### ***6.2 Incompatibilities***

Not applicable.

### ***6.3 Shelf life***

Clear blister: 3 years.

Green blister: 2 years.

### ***6.4 Special precautions for storage***

Store at or below 25°C. Protect from light.

### ***6.5 Nature and contents of container***

Al/PVC/PVdC blister packs containing 10, 30, 50, 90, 100 or 120 film coated tablets.

Not all pack types and sizes may be marketed.

## **6.6 Special precautions for disposal**

Not applicable.

---

## **7. Medicines Schedule**

---

Pharmacy Only Medicine.

---

## **8. Sponsor Details**

---

Viatris Ltd  
PO Box 11-183  
Ellerslie  
AUCKLAND  
[www.viatris.co.nz](http://www.viatris.co.nz)  
Telephone 0800 168 169

---

## **9. Date of First Approval**

---

4 October 2001

---

## **10. Date of Revision of the Text**

---

19 September 2022

Section changed	Summary of new information
Section 6.3	Removed bottle pack.
Section 6.5	Removed bottle pack.

Razene® is a Viatris company trade mark.