

DATA SHEET

Name of Medicine

IPCA-Enalapril Tablets 5 mg, 10 mg and 20 mg

Enalapril 5 mg, 10 mg and 20 mg Tablets

PRESENTATION

IPCA-Enalapril Tablets 5 mg are white to off-white, round, flat faced beveled edge tablets with breakline on one side and '5' debossed on the other side. Each tablet contains 5 mg of enalapril maleate (as the active ingredient)

IPCA-Enalapril Tablets 10 mg are white to off white, round, flat faced, beveled edge tablets with breakline on one side and '10' debossed on the other side. Each tablet contains 10 mg of enalapril maleate (as the active ingredient)

IPCA-Enalapril Tablets 20 mg are white to off white, round, flat faced, beveled edge tablets with breakline embossed on one side and '20' debossed on the other side. Each tablet contains 20 mg of enalapril maleate (as the active ingredient)

IPCA-Enalapril Tablets is the maleate salt of enalapril, a derivative of two amino acids, L-alanine and L-prolin. Following oral administration, enalapril is rapidly absorbed and then hydrolysed to enalaprilat, which is a highly specific, long acting, non-sulphydryl angiotensin converting enzyme inhibitor.

Indications

Treatment of:

- All grades of essential hypertension.
- Renovascular hypertension.
- All degrees of heart failure. In patients with symptomatic heart failure, IPCA-Enalapril Tablets is also indicated to:
 - Improve survival
 - Retard the progression of heart failure
 - Reduce hospitalisation for heart failure
- Prevention of symptomatic heart failure. In asymptomatic patients with left ventricular dysfunction, IPCA-Enalapril Tablets is indicated to:
 - Retard the development of symptomatic heart failure
 - Reduce hospitalisation for heart failure
- Prevention of coronary ischaemic events in patients with left ventricular dysfunction. IPCA-Enalapril Tablets is indicated to:
 - Reduce the incidence of myocardial infarction
 - Reduce hospitalisation for unstable angina pectoris

Dosage and Administration

Since absorption of IPCA-Enalapril tablets is not affected by food, the tablets may be administered before, during, or after meals.

Essential Hypertension

The initial dose is 5 mg and is given once daily. The usual maintenance dose is one 20 mg tablet taken once daily. The dosage should be adjusted according to the needs of the patient to a maximum of 40 mg daily.

Renovascular Hypertension

Since blood pressure and renal function in such patients may be particularly sensitive to ACE inhibition, therapy should be initiated with a lower starting dose (2.5 - 5 mg). The dosage should then be adjusted according to the needs of the patient. Most patients may be expected to respond to one 20 mg tablet taken once daily. For patients with hypertension who have been treated recently with diuretics, caution is recommended (see next paragraph).

Concomitant Diuretic Therapy in Hypertension

Symptomatic hypotension may occur following the initial dose of IPCA-Enalapril ; this is more likely in patients who are being treated currently with diuretics. Caution is recommended, therefore, since these patients may be volume- or salt-depleted. The diuretic therapy should be discontinued for 2-3 days prior to initiation of therapy with IPCA-Enalapril . If this is not possible, the initial dose of IPCA-Enalapril should be low (2.5 mg) to determine the initial effect on the blood pressure. Dosage should then be adjusted according to the needs of the patient.

Dosage in Renal Insufficiency

Generally, the intervals between the administration of enalapril should be prolonged and/or the dosage reduced.

Renal Status	Creatinine Clearance mL/min	Initial Dose mg/day
Mild Impairment	<80 >30 mL/min	5 mg
Moderate Impairment	<30 >10 mL/min	2.5 - 5 mg
Severe Impairment. Normally, these patients will be on dialysis [#]	<10 mL/min	2.5 mg on dialysis days ^{##}

Heart Failure/Asymptomatic Left Ventricular Dysfunction

The initial dose of IPCA-Enalapril in patients with symptomatic heart failure or asymptomatic left ventricular dysfunction is 2.5 mg, and it should be administered under close medical supervision to determine the initial effect on the blood pressure. IPCA-Enalapril may be used in the management of symptomatic heart failure usually with diuretics and, where appropriate, digitalis. In the absence of, or after effective management of, symptomatic hypotension following initiation of therapy with IPCA-

Enalapril in heart failure, the dose should be increased gradually to the usual maintenance dose of 20 mg, given in a single dose or two divided doses, as tolerated by the patient. This dose titration may be performed over a 2 to 4 week period, or more rapidly if indicated by the presence of residual signs and symptoms of heart failure. In patients with symptomatic heart failure this dosage regimen was effective in reducing mortality.

Blood pressure and renal function should be monitored closely both before and after starting treatment with IPCA-Enalapril (see Warnings and Precautions) because hypotension and (more rarely) consequent renal failure have been reported. In patients treated with diuretics the dose should be reduced, if possible, before beginning treatment with IPCA-Enalapril. The appearance of hypotension after the initial dose of IPCA-Enalapril does not imply that hypotension will recur during chronic therapy with IPCA-Enalapril and does not preclude continued use of the medicine. Serum potassium also should be monitored (see Interactions).

Contraindications

IPCA-Enalapril is contraindicated in patients who are hypersensitive to any component of this product and in patients with a history of angioneurotic oedema relating to previous treatment with an angiotensin-converting enzyme inhibitor and in patients with hereditary or idiopathic angioedema. All angiotensin converting enzyme inhibitors, including IPCA-Enalapril, are contraindicated in pregnancy because of the potential risk of foetotoxicity. IPCA-Enalapril should not be administered with aliskiren in patients with diabetes (see Interactions).

Warnings and Precautions

Symptomatic Hypotension

Symptomatic hypotension was seen rarely in uncomplicated hypertensive patients. In hypertensive patients receiving IPCA-Enalapril, hypotension is more likely to occur if the patient has been volume-depleted, e.g. by diuretic therapy, dietary salt restriction, dialysis, diarrhoea or vomiting (see Adverse Effects and Interactions). In patients with heart failure, with or without associated renal insufficiency, symptomatic hypotension has been observed. This is most likely to occur in those patients with more severe degrees of heart failure, as reflected by the use of high doses of loop diuretics, hyponatraemia or functional renal impairment. In these patients, therapy should be started under medical supervision and the patients should be followed closely whenever the dose of IPCA-Enalapril and/or diuretic is adjusted. Similar considerations may apply to patients with ischaemic heart or cerebrovascular disease in whom an excessive fall in blood pressure could result in a myocardial infarction or cerebrovascular accident. If hypotension occurs, the patient should be placed in the supine position and, if necessary, should receive an intravenous infusion of normal saline. A transient hypotensive response is not a contraindication to further doses, which can be given usually without difficulty once the blood pressure has increased after volume expansion. In some patients with heart failure, who have normal or low blood pressure, additional lowering of systemic blood pressure may occur with IPCA-Enalapril. This effect is anticipated, and usually is not a reason to

discontinue treatment. If hypotension becomes symptomatic, a reduction of dose and/or discontinuation of the diuretic and/or IPCA- Enalapril may be necessary.

Aortic Stenosis/Hypertrophic Cardiomyopathy

As with all vasodilators, ACE inhibitors should be given with caution to patients with obstruction in the outflow tract of the left ventricle.

Renal Function Impairment

In some patients with heart failure, hypotension following the initiation of therapy with IPCA- Enalapril may lead to some further impairment in renal function. Acute renal failure, usually reversible, has been reported in this situation. Patients with renal insufficiency may require reduced and/or less frequent doses of IPCA- Enalapril (see Dosage and Administration). In some patients, with bilateral renal artery stenosis or stenosis of the artery to a solitary kidney, increases of blood urea and serum creatinine, usually reversible upon discontinuation of therapy, have been seen. This is especially likely in patients with renal insufficiency. Some patients, with no apparent pre-existing renal disease, have developed minor and usually transient increases in blood urea and serum creatinine when IPCA- Enalapril has been given concomitantly with a diuretic. Dosage reduction and/or discontinuation of the diuretic and/or IPCA- Enalapril may be required.

Hypersensitivity/Angioneurotic Oedema

Angioneurotic oedema of the face, extremities, lips, tongue, glottis and/or larynx has been reported rarely in patients treated with angiotensin converting enzyme inhibitors, including IPCA- Enalapril. This may occur at any time during treatment. In such cases, IPCA- Enalapril should be discontinued promptly and appropriate monitoring should be instituted to ensure complete resolution of symptoms prior to dismissing the patient. Even in those instances where swelling of only the tongue is involved, without respiratory distress, patients may require prolonged observation since treatment with antihistamines and corticosteroids may not be sufficient. Very rarely, fatalities have been reported due to angioedema associated with laryngeal oedema or tongue oedema. Patients with involvement of the tongue, glottis or larynx are likely to experience airway obstruction, especially those with a history of airway surgery. Where there is involvement of the tongue, glottis or larynx, likely to cause airway obstruction, appropriate therapy, which may include subcutaneous epinephrine solution 1:1000 (0.3 mL to 0.5 mL) and/or measures to ensure a patent airway, should be administered promptly. Black patients receiving ACE inhibitors have been reported to have a higher incidence of angioedema compared to non-blacks. Patients with a history of angioedema unrelated to ACE inhibitor therapy may be at increased risk of angioedema while receiving an ACE inhibitor (also see Contraindications).

Anaphylactoid Reactions During Hymenoptera Desensitisation

Rarely, patients receiving ACE inhibitors during desensitisation with hymenoptera venom have experienced life-threatening anaphylactoid reactions. These reactions were avoided by temporarily withholding ACE inhibitor therapy prior to each desensitisation.

Haemodialysis Patients

Anaphylactoid reactions have been reported in patients dialysed with high flux membranes (e.g. AN 69*) and treated concomitantly with an ACE inhibitor. In these patients consideration should be given to using a different type of dialysis membrane or a different class of anti-hypertensive agent.

Anaphylactoid Reactions During LDL Apheresis

Rarely, patients receiving ACE inhibitors during low density lipoprotein (LDL)-apheresis with dextran sulfate have experienced life-threatening anaphylactoid reactions. These reactions were avoided by temporarily withholding ACE inhibitor therapy prior to each apheresis.

Cough

Cough has been reported with the use of ACE inhibitors. Characteristically, the cough is non-productive, persistent and resolves after discontinuation of therapy. ACE inhibitor-induced cough should be considered part of the differential diagnosis of cough.

Surgery Anaesthesia

In patients undergoing major surgery or during anaesthesia with agents that produce hypotension, enalapril blocks angiotensin II formation, secondary to compensatory renin release. If hypotension occurs and is considered to be due to this mechanism, it can be corrected by volume expansion.

Hyperkalaemia

Risk factors for the development of hyperkalaemia include renal insufficiency, diabetes mellitus, and concomitant use of potassium-sparing diuretics (e.g., spironolactone, eplerenone, triamterene, or amiloride), potassium supplements, or potassium-containing salt substitutes. The use of potassium supplements, potassium-sparing diuretics, or potassium-containing salt substitutes particularly in patients with impaired renal function may lead to a significant increase in serum potassium. Hyperkalaemia can cause serious, sometimes fatal, arrhythmias. If concomitant use of IPCA- Enalapril and any of the above-mentioned agents is deemed appropriate, they should be used with caution and with frequent monitoring of serum potassium. (See Interactions, *Serum Potassium*.)

Hypoglycaemia

Diabetic patients treated with oral anti-diabetic agents or insulin starting an ACE inhibitor should be told to closely monitor for hypoglycaemia, especially during the first month of combined use. (See Interactions.)

Use in Pregnancy

The use of IPCA- Enalapril during pregnancy is contraindicated. When pregnancy is detected, IPCA- Enalapril should be discontinued as soon as possible. In a published retrospective epidemiological study, infants whose mothers had taken an ACE inhibitor medicine during the first trimester of pregnancy appeared to have an increased risk of major congenital malformations compared with infants whose mothers had not undergone first trimester exposure to ACE inhibitor medicines. The number of cases of birth defects

is small and the findings of this study have not yet been repeated. ACE inhibitors can cause foetal and neonatal morbidity and mortality when administered to pregnant women during the second and third trimesters. Use of ACE inhibitors during this period has been associated with foetal and neonatal injury including hypotension, renal failure, hyperkalaemia, and/or skull hypoplasia in the newborn. Maternal oligohydramnios, presumably representing decreased foetal renal function, has occurred and may result in limb contractures, craniofacial deformations and hypoplastic lung development. These adverse effects to the embryo and foetus do not appear to have resulted from intrauterine ACE-inhibitor exposure limited to the first trimester. Infants whose mothers have taken IPCA- Enalapril should be closely observed for hypotension, oliguria and hyperkalaemia. Enalapril, which crosses the placenta, has been removed from the neonatal circulation by peritoneal dialysis with some clinical benefit, and theoretically may be removed by exchange transfusion.

Nursing Mothers

Enalapril and enalaprilat are secreted in human milk in trace amounts. ACE inhibitors and angiotensin II receptor antagonists should not be used by breastfeeding mothers in the first few weeks after delivery because of possible profound neonatal hypotension; preterm babies may be at particular risk. In mothers who are breastfeeding older infants, the use of captopril, enalapril, or quinapril may be considered.

Paediatric Use

The safety and effectiveness of IPCA- Enalapril tablets have been established in hypertensive paediatric patients age 1 month to 16 years. Use of IPCA- Enalapril in these age groups is supported by evidence from adequate and well-controlled studies of IPCA- Enalapril in paediatric and adult patients as well as by published literature in paediatric patients. In a multiple dose pharmacokinetic study in 40 hypertensive paediatric patients, excluding neonates IPCA-Enalapril tablets was generally well tolerated. Pharmacokinetics following oral administration of enalapril are similar in these patients and comparable to historical values in adults. In a clinical study involving 110 hypertensive paediatric patients 6 to 16 years of age, patients who weighed <50 kg received either 0.625, 2.5 or 20 mg of enalapril daily and patients who weighed ≥50 kg received either 1.25, 5 or 40 mg of enalapril daily. Enalapril administration once daily lowered trough blood pressure in a dose-dependent manner. The dose-dependent anti-hypertensive efficacy of enalapril was consistent across all subgroups (age, Tanner stage, gender, race). However, the lowest doses studied, 0.625 mg and 1.25 mg, corresponding to an average of 0.02 mg/kg once daily, did not appear to offer consistent anti-hypertensive efficacy. The maximum dose studied was 0.58 mg/kg (up to 40 mg) once daily. In this study, IPCA- Enalapril was generally well tolerated. The adverse experience profile for paediatric patients is not different from that seen in adult patients. IPCA- Enalapril is not recommended in neonates and in paediatric patients with glomerular filtration rate <30 mL/min/1.73 m², as no data are available.

Pancreatitis

Pancreatitis may occur with angiotensin converting enzyme inhibitors and patients with abdominal pain on ACE inhibitors should be tested accordingly.

Effect on Ability to Use and Drive Machinery

Individual responses to medication may vary. Certain adverse effects that have been reported with IPCA- Enalapril may affect some patient's ability to drive or operate machinery (see Adverse Effects).

Animal Toxicology

Studies were performed to assess the teratogenic potential of enalapril in rats and rabbits and its effect on reproduction and postnatal development in rats. Enalapril given to pregnant rats at doses up to 1200 mg/kg/day (2000 times the maximum human dose) from Day 6 through Day 17 of gestation did not reveal any evidence of embryo lethality or teratogenicity. Decreased average foetal weight occurred at 1200 mg/kg/day, but did not occur at this dosage level if the pregnant animals were given physiological saline for drinking instead of tap water during the dosing period. Average foetal weights were not affected in unsupplemented rats given up to 120 mg/kg/day. Decreased maternal weight gain during the dosing period occurred at doses as low as 12 mg/kg/day, but did not occur in saline-supplemented rats given 1200 mg/kg/day. Saline supplementation in rats given 1200 mg/kg/day also prevented increases in serum urea nitrogen which occurred at doses as low as 100 mg/kg/day in unsupplemented rats (lowest dose level examined in pregnant rats), but only partially inhibited increases in serum potassium. In supplemented rats serum potassium was elevated in rats given 200 mg/kg/day, but not 100 mg/kg/day. Enalapril was not teratogenic to saline-supplemented rabbits given doses up to 30 mg/kg/day (50 times the maximum human dose) from Day 6 through Day 18 of gestation. At 30 mg/kg/day (50 times the maximum human dose), enalapril produced maternal and foetal toxicity. Doses of 3 and 10 mg/kg/day were without maternotoxic or foetotoxic effects in saline-supplemented rabbits. There were no adverse effects on reproductive performance in male and female rats treated with 10 to 90 mg/kg/day of enalapril. An *in vitro* Coombs' test of enalapril and its active metabolite (enalaprilat) did not show a positive Coombs' reaction within the range of concentrations tested (which did not induce direct haemolysis). Neither enalapril nor enalaprilat was mutagenic in the Ames microbial mutagen test with or without metabolic activation. Enalapril was also negative in the following genotoxicity studies: Rec-Assay, reverse mutation assay with *E.coli*, sister chromatid exchange with cultured mammalian cells, and the micro-nucleus test with mice, as well as an *in vivo* cytogenic study using mouse bone marrow. There was no evidence of a carcinogenic effect when enalapril was administered for 106 weeks to rats at a dose up to 90 mg/kg/day (150 times the maximum daily human dose). Enalapril has also been administered for 94 weeks to male and female mice at doses up to 90 and 180 mg/kg/day, respectively (150 and 300 times the maximum daily dose for humans) and showed no evidence of carcinogenicity.

Dual blockade of the renin-angiotensin-aldosterone system

It has been reported in the literature that in patients with established atherosclerotic disease, heart failure, or with diabetes with end organ damage, dual blockade of the renin-angiotensin-aldosterone system, is associated with a higher frequency of hypotension, syncope, hyperkalaemia, and changes in renal function (including acute renal failure) as compared to use of a single renin-angiotensin-aldosterone system agent. Dual blockade (e.g, by adding an ACE-inhibitor to an angiotensin II receptor antagonist) should be limited to individually defined cases with close monitoring of renal function.

Adverse Effects

IPCA- Enalapril has been demonstrated to be generally well tolerated. For the most part, adverse experiences have been mild and transient in nature, and have not required discontinuation of therapy. The following adverse effects have been associated with the use of IPCA- Enalapril tablets: Dizziness and headache were the more commonly reported adverse effects. Fatigue and asthenia were reported in 2-3% of patients. Other adverse effects occurred in less than 2% of patients and included hypotension, orthostatic hypotension, syncope, nausea, diarrhoea, muscle cramps, and cough. Skin rash was reported in 1.2% of patients and taste disturbances in 0.5% of patients. Less frequently renal dysfunction, renal failure and oliguria have been reported.

Hypersensitivity/Angioneurotic Oedema

Angioneurotic oedema of the face, extremities, lips, tongue, glottis and/or larynx has been reported rarely (see Warnings and Precautions). In very rare cases, intestinal angioedema has been reported with angiotensin converting enzyme inhibitors including enalapril.

Adverse effects which occurred very rarely, either during controlled clinical trials or after the medicine was marketed, include

Cardiovascular: Myocardial infarction or cerebrovascular accident, possibly secondary to excessive hypotension in high risk patients (see Warnings and Precautions), chest pain, palpitations, rhythm disturbances, angina pectoris, Raynaud's phenomenon
Endocrine: Syndrome of inappropriate anti-diuretic hormone secretion (SIADH)
Gastrointestinal: Ileus pancreatitis (see Warnings and Precautions), hepatic failure, hepatitis - either hepatocellular or cholestatic, jaundice, abdominal pain, vomiting, dyspepsia, constipation, anorexia, stomatitis
Metabolic: Cases of hypoglycaemia in diabetic patients on oral anti-diabetic agents or insulin have been reported (see Interactions)
Nervous System/Psychiatric: Depression, confusion, somnolence, insomnia, nervousness, paresthesia, vertigo, dream abnormality
Respiratory: Pulmonary infiltrates, bronchospasm/asthma, dyspnoea, rhinorrhoea, sore throat and hoarseness
Skin: Diaphoresis, erythema multiforme, exfoliative dermatitis, Stevens-Johnson syndrome, toxic epidermal necrolysis, pemphigus, pruritus, urticaria, alopecia
Other: Impotence, flushing, taste alteration, tinnitus, glossitis, blurred vision
A symptom complex has been reported which may include some or all of the following: fever, serositis, vasculitis, myalgia/myositis, arthralgia/arthritis, a positive ANA, elevated ESR, eosinophilia, and leukocytosis. Rash, photosensitivity or other dermatologic manifestations may occur.

Laboratory Test Findings

Clinically important changes in standard laboratory parameters were rarely associated with administration of IPCA- Enalapril. Increases in blood urea and serum creatinine, and elevations of liver enzymes and/or serum bilirubin have been seen. These are usually reversible upon discontinuation of IPCA- Enalapril. Hyperkalaemia and hyponatraemia have occurred. Decreases in haemoglobin and haematocrit have been reported. Since the medicine was marketed a small number of cases of neutropaenia, thrombocytopaenia, bone marrow depression, and agranulocytosis have been reported in which a causal relationship to therapy with IPCA- Enalapril could not be excluded.

Interactions

Anti-hypertensive Therapy

The combination of IPCA- Enalapril with other anti-hypertensive medicines may increase the anti-hypertensive effect, especially in combination with diuretics. The combination of IPCA- Enalapril with beta-adrenergic blocking agents, methyldopa, or calcium entry blockers has been shown to improve the efficacy of lowering the blood pressure. Ganglionic blocking agents or adrenergic blocking agents, combined with IPCA- Enalapril, should only be administered under careful observation of the patient. A possible drop in serum-potassium due to thiazide-containing diuretics may be reduced by simultaneous administration of IPCA- Enalapril. There are no clinically significant pharmacokinetic medicine interactions between enalapril maleate and the following compounds: hydrochlorothiazide, furosemide, digoxin, timolol, methyldopa, warfarin, indomethacin and sulindac. Propranolol co-administered with enalapril maleate reduces serum enalaprilat concentrations, but this does not appear to be of any clinical significance. Since cimetidine does not interact with enalapril maleate in animals, it is not anticipated that a medicine interaction will occur in humans.

Serum Potassium

In clinical trials, serum potassium usually remained within normal limits. In hypertensive patients treated with IPCA- Enalapril alone for up to 48 weeks, mean increases in serum potassium of approximately 0.2 mEq/L were observed. In patients treated with IPCA- Enalapril plus a thiazide diuretic, the potassium-losing effect of the diuretic was attenuated usually by the effect of enalapril. If IPCA- Enalapril is given with a potassium-losing diuretic, diuretic-induced hypokalaemia may be ameliorated. Risk factors for the development of hyperkalaemia include renal insufficiency, diabetes mellitus, and concomitant use of potassium-sparing diuretics (e.g. spironolactone, eplerenone, triamterene or amiloride) potassium supplements, or potassium-containing salt substitutes. The use of potassium supplements, potassium-sparing diuretics or potassium containing salt substitutes particularly in patients with impaired renal function may lead to a significant increase in serum potassium. If concomitant use of IPCA- Enalapril and the above mentioned agents is deemed appropriate, they should be used with caution and with frequent monitoring of serum potassium. (See Warnings and Precautions, *Hyperkalaemia*.)

Anti-diabetics

Epidemiological studies have suggested that concomitant administration of ACE inhibitors and anti-diabetic medicines (insulins, oral hypoglycaemic agents) may cause an increased blood-glucose-lowering effect with risk of hypoglycaemia. This phenomenon appeared to be more likely to occur during the first weeks of combined treatment and in patients with renal impairment. In diabetic patients treated with oral anti-diabetic agents or insulin, glycaemic control should be closely monitored for hypoglycaemia, especially during the first month of treatment with an ACE inhibitor.

Serum Lithium

As with other medicines which eliminate sodium, lithium clearance may be reduced. Therefore, the serum lithium levels should be monitored carefully if lithium salts are to be administered.

Non-Steroidal Anti-Inflammatory Medicines including Selective Cyclooxygenase-2 Inhibitors

Non-steroidal anti-inflammatory medicines (NSAIDs) including selective cyclooxygenase-2 inhibitors (COX-2 inhibitors) may reduce the effect of diuretics and other anti-hypertensive medicines. Therefore, the anti-hypertensive effect of angiotensin II receptor antagonists or ACE inhibitors may be attenuated by NSAIDs including selective COX-2 inhibitors. In some patients with compromised renal function (e.g., elderly patients or patients who are volume-depleted, including those on diuretic therapy) who are being treated with non-steroidal anti-inflammatory medicines, including selective cyclooxygenase-2 inhibitors, the co-administration of angiotensin II receptor antagonists or ACE inhibitors may result in a further deterioration of renal function, including possible acute renal failure. These effects are usually reversible. Therefore, the combination should be administered with caution in patients with compromised renal function.

Gold

Nitritoid reactions (symptoms include facial flushing, nausea, vomiting and hypotension) have been reported rarely in patients on therapy with injectable gold (sodium aurothiomalate) and concomitant ACE inhibitor therapy including enalapril.

Dual blockade of the renin-angiotensin-aldosterone system

Dual blockade of the renin-angiotensin-aldosterone system, (RAAS) with angiotensin receptor blockers, ACE inhibitors, or direct renin inhibitors (such as aliskiren) is associated with increased risks of hypotension, syncope, hyperkalaemia, and changes in renal function (including acute renal failure) compared to monotherapy. Closely monitor blood pressure, renal function, and electrolytes in patients on IPCA-Enalapril and other agents that affect the RAAS. Do not coadminister aliskiren with IPCA-Enalapril in patients with diabetes. Avoid use of aliskiren with IPCA-Enalapril in patients with renal impairment (GFR <60 ml/min).

Overdosage

Limited data are available for overdosage in humans. The most prominent feature of overdosage reported to date is marked hypotension, beginning some six hours after ingestion of tablets, concomitant with blockade of the renin-angiotensin system, and stupor. Serum enalaprilat levels 100 times and 200 times higher than usually seen after therapeutic doses have been reported after ingestion of 300 mg of 440 mg of enalapril, respectively. The recommended treatment of overdosage is intravenous infusion of normal saline solution. If available, angiotensin II infusion may be beneficial. If ingestion is recent, induce emesis. Enalaprilat may be removed from the general circulation by haemodialysis. (See Warnings and Precautions, *Haemodialysis Patients*.)

Actions

Mechanism of Action

Angiotensin converting enzyme (ACE) is a peptidyl dipeptidase which catalyses the conversion of angiotensin I to the pressor substance angiotensin II. After absorption, enalapril is hydrolysed to enalaprilat, which inhibits ACE. Inhibition of ACE results in decreased plasma angiotensin II, which leads to increased plasma renin activity (due to removal of negative feedback of renin release), and decreased aldosterone secretion. ACE is identical to kininase II. Thus IPCA-Enalapril may also block the degradation of bradykinin, a potent vasodepressor peptide. However, the role that this plays in the therapeutic effects of IPCA- Enalapril remains to be elucidated. While the mechanism through which IPCA-Enalapril lowers blood pressure is believed to be primarily suppression of the renin-angiotensin-aldosterone system, which plays a major role in the regulation of blood pressure, IPCA- Enalapril is anti-hypertensive even in patients with low-renin hypertension. The onset of action of oral IPCA- Enalapril is gradual and smooth; it begins within one hour and its effects usually continue for 24 hours. Consequently, IPCA- Enalapril may be administered on a once-daily basis, with the advantages this brings in convenience and compliance. The Studies of (patients with) Left Ventricular Dysfunction (SOLVD) was a multicentre, placebo controlled, double blind study of 6797 patients assessed as having Left Ventricular Dysfunction. All patients had a Left Ventricular Ejection Fraction of <35% and were classified as New York Heart Association Class I - IV (NYHA). The 2569 symptomatic patients (primarily NYHA Class II & III) were randomised into a Treatment arm, whilst the 4228 asymptomatic patients (NYHA Class I) were randomised into the Prevention arm. The combined results demonstrated an overall reduced risk for the development of major ischaemic events. IPCA- Enalapril decreased the incidence of myocardial infarction and reduced the number of hospitalisations for unstable angina pectoris in patients with left ventricular dysfunction. In the Prevention arm, IPCA- Enalapril significantly prevented the development of symptomatic heart failure and reduced the number of hospitalisations for heart failure. IPCA- Enalapril in the Treatment arm, as an adjunct to conventional therapy, significantly reduced overall mortality and hospitalisation for heart failure and improved NYHA functional class. In CONSENSUS, a similar study involving 253 patients with severe heart failure (NYHA Class IV), IPCA- Enalapril was shown to improve symptoms and reduce mortality significantly. The cardio-protective properties of IPCA- Enalapril were demonstrated in these studies by the beneficial effects on survival and retardation of the progression of heart failure in patients with symptomatic heart

failure; retardation of the development of symptomatic heart failure in asymptomatic patients with left ventricular dysfunction; and prevention of coronary ischaemic events in patients with left ventricular dysfunction, specifically reduction in the incidence of myocardial infarction and reduction in hospitalisation for unstable angina pectoris.

Pharmacokinetics

Oral enalapril is rapidly absorbed, with peak serum concentrations of enalapril occurring within one hour. Based on urinary recovery, the extent of absorption of enalapril from oral enalapril tablet is approximately 60%. Following absorption, oral enalapril is rapidly and extensively hydrolysed to enalaprilat, a potent angiotensin converting enzyme inhibitor. Similar peak serum concentrations of enalaprilat occur about four hours after an oral dose of enalapril. Excretion of enalaprilat is primarily renal. The principal components in urine are enalaprilat, accounting for about 40% of the dose, and intact enalapril. Except for conversion to enalaprilat, there is no evidence for significant metabolism of enalapril. The serum concentration profile of enalaprilat exhibits a prolonged terminal phase, apparently associated with binding to ACE. In subjects with normal renal function, steady state serum concentrations of enalaprilat were achieved by the fourth day of administration of oral enalapril. The effective half-life for accumulation of enalaprilat following multiple doses of oral enalapril is 11 hours. The absorption of oral IPCA- Enalapril is not influenced by the presence of food in the gastrointestinal tract. The extent of absorption and hydrolysis of enalapril are similar for the various doses in the recommended therapeutic range. Studies in dogs indicate that enalapril crosses the blood-brain barrier poorly, if at all; enalaprilat does not enter the brain. Multiple doses of oral enalapril in rats do not result in accumulation in any tissues. Milk of lactating rats contains radioactivity following administration of ¹⁴C enalapril maleate. Radioactivity was found to cross the placenta following administration of ¹⁴C enalapril maleate to pregnant hamsters. There is no significant change in the plasma half-life of IPCA- Enalapril in elderly patients. No pharmacokinetic data is available on the effect of IPCA- Enalapril in patients with hepatic dysfunction.

Pharmacodynamics

Administration of IPCA- Enalapril to patients with hypertension results in a reduction of both supine and standing blood pressure without a significant increase in heart rate. Symptomatic postural hypotension is infrequent. In some patients the development of optimal blood pressure reduction may require several weeks of therapy. Abrupt withdrawal of IPCA- Enalapril has not been associated with rapid increase in blood pressure. Effective inhibition of ACE activity usually occurs 2 to 4 hours after oral administration of an individual dose of enalapril. Onset of anti-hypertensive activity was usually seen at one hour, with peak reduction of blood pressure achieved by 4-6 hours after administration. The duration of effect is dose related. However, at recommended doses, anti-hypertensive and haemodynamic effects have been shown to be maintained for at least 24 hours. Anti-hypertensive treatment with IPCA- Enalapril leads to a significant regression of left ventricular hypertrophy with preservation of left ventricular systolic performance. In haemodynamic studies in patients with essential hypertension, blood pressure reduction was accompanied by a reduction in peripheral arterial resistance with an increase in cardiac output and little or no change in heart rate. Following

administration of IPCA- Enalapril there was an increase in renal blood flow; glomerular filtration rate was unchanged. There was no evidence of sodium or water retention. However, in patients with low pre-treatment glomerular filtration rates, the rates were usually increased. Chronic administration of IPCA- Enalapril to patients with essential hypertension and renal insufficiency may be associated with improvements in renal function, evidenced by increased glomerular filtration rate. In short term clinical studies in diabetic and non-diabetic patients with renal disease, decreases in albuminuria and urinary excretion of IgG and total urinary protein were seen after the administration of enalapril. Treatment with IPCA- Enalapril has been associated with favourable effects on plasma lipoprotein fractions and favourable or no effect on total cholesterol levels. In patients with mild to moderate heart failure, enalapril retarded progressive cardiac dilatation/enlargement and failure, as evidenced by reduced left ventricular end diastolic and systolic volumes and improved ejection fraction. Clinical data have shown that enalapril reduced the frequency of ventricular arrhythmias in patients with heart failure, although the underlying mechanisms and clinical significance are not known.

Pharmaceutical Precautions

Store below 25°C in a dry place. Protect from light.

Medicine Classification

Prescription Medicine.

Package Quantities

IPCA- Enalapril Tablets 5 mg: Tablets are packed in blister pack of 10 tablets which are further placed in an outer carton to give packs of 30 and 90 tablets.

IPCA- Enalapril Tablets 10 mg: Tablets are packed in blister pack of 10 tablets which are further placed in an outer carton to give packs of 30 and 90 tablets.

IPCA- Enalapril Tablets 20 mg: Tablets are packed in blister pack of 10 tablets which are further placed in an outer carton to give packs of 30 and 90 tablets.

Name and Address of the sponsor

IpcA Pharma (NZ) Pty. Ltd
3-A, St. Ostwalds Road
Greenlane, Auckland 1061

Distributed By

Miro Healthcare Ltd.,
Auckland, New Zealand

Date of Preparation

12th August 2025

Version : 03